


This article was downloaded by: [190.151.168.196]

On: 10 January 2015, At: 17:39

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Avian Pathology

Publication details, including instructions for authors and subscription information:
<http://www.tandfonline.com/loi/cavp20>

A comparison of the efficacy of cleaning and disinfection methods in eliminating *Salmonella* spp. from commercial egg laying houses

Juan J. Carrique-Mas^a, Clara Marín^b, Mark Breslin^a, Ian McLaren^a & Rob Davies^a

^a Department of Food and Environmental Safety, Veterinary Laboratories Agency, New Haw, Addlestone, Surrey, KT15 3NB, UK

^b Centro de Tecnología Animal, Instituto Valenciano de Investigaciones Agrarias, Polígono La Esperanza, 12400, Segorbe, Spain

Published online: 18 Sep 2009.

To cite this article: Juan J. Carrique-Mas, Clara Marín, Mark Breslin, Ian McLaren & Rob Davies (2009) A comparison of the efficacy of cleaning and disinfection methods in eliminating *Salmonella* spp. from commercial egg laying houses, *Avian Pathology*, 38:5, 419-424, DOI: [10.1080/03079450903193768](https://doi.org/10.1080/03079450903193768)

To link to this article: <http://dx.doi.org/10.1080/03079450903193768>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

A comparison of the efficacy of cleaning and disinfection methods in eliminating *Salmonella* spp. from commercial egg laying houses

Juan J. Carrique-Mas^{1*}, Clara Marín², Mark Breslin¹, Ian McLaren¹ and Rob Davies¹

¹Department of Food and Environmental Safety, Veterinary Laboratories Agency, New Haw, Addlestone, Surrey KT15 3NB, UK, and ²Centro de Tecnología Animal, Instituto Valenciano de Investigaciones Agrarias, Polígono La Esperanza, 12400 Segorbe, Spain

Effective terminal cleaning and disinfection (C&D) is regarded as a necessary step for the elimination of *Salmonella* spp. from laying houses. A total of 60 commercial laying houses that had housed laying flocks infected with *Salmonella enterica* serovar Enteritidis or *Salmonella enterica* serovar Typhimurium that were representative of all production systems (cage, barn, free-range) were intensively sampled immediately after C&D as well as in the follow-on flock. The procedures investigated were: (1) a compound disinfectant consisting of a mixture of formaldehyde, glutaraldehyde and quarternary ammonium applied at the recommended concentration; (2) a 10% (vol/vol) dilution of the standard 37% commercial formalin, applied by a contractor; and (3) other disinfection procedures selected and applied by the farmer. The recovery of *Salmonella* in the cleaned and disinfected houses was variable, with samples from floor and dropping boards/belts (cage houses) and scratching areas (non-cage houses) being the most likely to remain contaminated. In cage houses, the use of the 10% formalin dilution led to a statistically greater reduction in the sample prevalence than using any of the other C&D methods. A negative post-C&D result predicted clearance of *Salmonella* in 52% of cases, although the isolation of *Salmonella* from the houses immediately after C&D was not a perfect predictor of carry-over of infection.

Introduction

Salmonella infection is one of the most common causes of infectious gastroenteritis in humans worldwide (World Health Organization, 2005). In the European Union (EU), *Salmonella enterica* serovar Enteritidis is the agent most frequently involved in human salmonellosis and it is also the serovar most widespread in egg production, followed at a distance by *S. enterica* serovar Typhimurium (Anonymous, 2006).

Eggs are regarded as the main source of infection of *Salmonella* Enteritidis for humans (Gillespie *et al.*, 2005; de Jong & Ekdahl, 2006). Over the past few years, the egg industry has stepped up its efforts to control *Salmonella* in laying houses. These have mostly involved an upgrade of biosecurity and hygiene procedures. In the UK and other countries, vaccination with attenuated, live or killed vaccines is also now widespread among commercial laying flocks.

Despite this, effective interventions to control *Salmonella* infection during the life of the flocks are limited, and effective terminal cleaning and disinfection (C&D) is regarded as a crucial step to reduce the risk of infection to new flocks placed in houses that have held infected birds (van de Giessen *et al.*, 1994).

In the UK, carry-over of *S. Enteritidis* between consecutive laying flocks has been shown to be a common

occurrence (Carrique-Mas *et al.*, 2008b, 2009; Wales *et al.*, 2006b), which was concomitant with a generally low standard of C&D, although the presence of rodents has also been described to be a major factor contributing to carry-over (Carrique-Mas *et al.*, 2009). C&D is a costly and laborious task, and its success in eliminating *Salmonella* from the houses depends on the attention to detail, as well as on the choice and correct application of disinfectants (Davies & Wray, 1995; Wales *et al.*, 2006a). There is a wide range of disinfectant formulations available on the market, but there are marked differences in their efficacy (McDonnell & Russell, 1999). Furthermore the efficacy of a disinfectant in the field is highly dependent on the level of residual organic matter remaining on the house surfaces. Because of their design, C&D of cage laying houses is known to be particularly problematic (Davies & Breslin, 2003; Wales *et al.*, 2006a). Aldehydes are generally more effective than other disinfectants in poultry houses (Rose *et al.*, 2000; Wales *et al.*, 2006a), and where formaldehyde has been investigated it has performed better than any other product, both in laboratory models using artificially inoculated surfaces (Gradel *et al.*, 2004) and in field conditions (Davies & Wray, 1995). Historically C&D has typically been carried out in the same manner over subsequent production

*To whom correspondence should be addressed. Tel: +44 1932 357814, ext. 2814. Fax: +44 1932 357595. E-mail: j.carrique-mas@vla.defra.gsi.gov.uk
Received 2 April 2009

cycles, irrespective of the *Salmonella* status of the laying houses. However the introduction in 2009 of *Salmonella* legislation across the EU (enacted by National Control Programmes) banning the sale of fresh eggs from *S. Enteritidis*-positive or *S. Typhimurium*-positive flocks (EC No. 1237/2007) is likely to alter this situation, and more effective disinfection procedures will be sought by farmers with *Salmonella*-positive flocks.

The aim of the present study was to determine the comparative effectiveness of disinfection programmes in *Salmonella*-positive cage and non-cage (i.e. free-range, barn) houses in the field; in particular, an intensive method using 10% formalin applied with a pressure washer by a specialist contractor.

Materials and Methods

Farms, laying houses and disinfection procedures. A total of 60 post-C&D visits to houses where either *S. Enteritidis* or *S. Typhimurium* had been isolated were carried out on 47 different laying houses in 23 different farms (sites) between February 2003 and September 2008. Prior to disinfection, houses were washed using a pressure washer and allowed to dry. The disinfection procedures investigated were: (1) a 10% (vol/vol) formalin dilution of the standard 37% commercial dilution, applied by a specialist contractor (Elebert Pestforce, Lymm, UK) using a high-pressure washer to run-off point; (2) a formaldehyde, glutaraldehyde quaternary ammonium (FGQ) compound (Superkill®; AFS Animal Care, Thetford, UK) applied at the “General Orders” (GO) rate (1:22) (Department for Environment, Food and Rural Affairs [DEFRA], 2009) (the GO rate is the minimum concentration recommended by DEFRA, based on an *in vitro* suspension approval test), with the disinfectant applied by the farmer by power or pressure washer to run-off point; and (3) “in house” procedures, other disinfection procedures that were routinely used by the producers (Table 1). In all cases the disinfectants were applied to all surfaces of the house and anteroom (where present). Houses were restocked 1 to 3 days after the disinfectants had dried out from the house surfaces.

Sampling for *Salmonella* to assess the effectiveness of C&D. The effectiveness of the C&D procedure was assessed by the culture of hand-held gauze swabs impregnated with buffered peptone water (BPW) that were used to wipe a range of surfaces in the house, before they were placed back into jars containing 225 ml BPW (Davies & Breslin, 2003; Wales *et al.*, 2006a; Carrique-Mas *et al.*, 2008b). For cage houses, 10 samples from each of the following areas were collected: cage interiors (eight cages per swab); drinker cups/troughs (eight per swab); feed troughs (0.5 m² per swab); droppings boards/belts/flaps (1 m² per swab); house floor (0.5 m² per swab); and egg belts (0.5 m² per swab). From non-cage houses, samples were collected from: drinkers (four bell drinkers per swab); feeders (0.5 m² per swab); scratching area (if present) (1 m² per swab); slats (0.5 m² per swab); and nest box interiors (five per swab). In addition, in free-range houses the soil from the paddocks was sampled by scraping off 25 g topsoil, which was added to the 225 ml BPW jars (Davies & Breslin, 2003). For each sampling visit and sample type, a prevalence of positive samples was calculated.

Sampling of flocks in lay. For each house investigated, laying flocks before and after terminal C&D were sampled in an identical way. Dust and faeces were collected from the occupied houses from flocks in lay placed before disinfection and from the first flock after restocking. For this, 10 pooled faeces (25 g) and 10 dust samples (15 g) were collected from each flock directly into 225 ml BPW pots using large gauze swabs. This is regarded as a very sensitive sampling method (Carrique-Mas *et al.*, 2008a).

Bacteriological methods. Samples in BPW were cultured following a simplified protocol of ISO 6579:2002 (Annex D), consisting of pre-enrichment in BPW followed by enrichment in modified semi-solid Rappaport Vassiliadis medium and plating onto Rambach agar (Wales *et al.*, 2006a). Suspect *Salmonella* colonies were confirmed by serotyping using the Kauffmann – White typing scheme.

Statistical analyses. The level of contamination of flocks before and after C&D was estimated by calculating an average (weighted) percentage of faeces and dust samples that were positive. Chi-square tests were used to assess differences in proportions and the (non-parametric) Wilcoxon test statistic was used to compare the sample prevalence before and after the disinfection. All analyses were carried out using S-Plus (Insight, USA).

Results

Post C&D sampling visit. All houses sampled after C&D had previously contained flocks positive for *S. Enteritidis*, except one case where *S. Typhimurium* was present. A total of 1080, 1140, and 660 samples (combined data) were collected for houses treated with the “in house”, FGQ (G.O rate), and 10% formalin C&D programmes, respectively. The total prevalence of positive samples was 31.5%, 9.0%, and 3.3%, respectively. For cage houses, the sample prevalence was (from highest to lowest): floors (24.8%), dropping boards/belts/flaps (22.7%), drinkers (15.2%), feeders (13.8%), cage interiors (11.4%) and egg belts (9.4%). A comparable proportion of positive samples were recovered from floor and dropping board samples ($\chi^2=0.35$; $P=0.55$), and the proportion of positive samples from both were significantly greater than that obtained from any other type of sample ($P < 0.05$). In non-cage houses, the highest sample prevalence was in scratching areas (32.6%), drinkers (7.5%), feeders (5.7%), nest boxes (3.3%) and slats (2.7%). The proportion of positive samples from scratching areas was greater than any other type of sample ($P < 0.01$). A total of 35.7% soil samples from the paddocks tested positive. The mean sample prevalence by disinfection type is presented in Table 2, and the median prevalence of positive samples by sample type and disinfection method is shown in Figures 1 and 2. Results for non-cage houses treated using the formalin 10% method are not shown given that only two houses were examined.

Table 1. Number of laying houses investigated by type of production and disinfection procedure

	Cage	Non-cage	All
“In-house” method	18	6	24
FGQ disinfectant (applied at a lower dilution than the GO rate)	7	2	9
Peroxygen-based disinfectant	6	2	8
Non-DEFRA-approved disinfectant	2	0	2
No disinfectant used	3	2	5
FGQ (at the GO rate)	19	4	23
Formalin 10%	11	2	13
Total	48	12	60

Table 2. *Salmonella*-positive sample prevalence by sample type

	"In-house"		FGQ (GO rate)		Formalin 10%	
	% positive	95% confidence interval	% positive	95% confidence interval	% positive	95% confidence interval
Cage houses						
Cage interior	22.9	16.8 to 29.0	4.8	1.7 to 8.0	1.8	0.0 to 4.3
Drinkers	26.0	19.6 to 32.3	12.5	7.7 to 17.4	1.8	0.0 to 4.3
Feeders	26.6	20.0 to 33.3	8.1	4.2 to 12.0	3.6	0.1 to 7.1
Dropping boards	44.9	37.2 to 52.6	10.3	5.4 to 15.2	6.4	1.8 to 10.9
Egg belts	21.4	15.2 to 27.6	3.9	1.2 to 6.7	1.8	0.0 to 4.3
Floor	47.0	39.3 to 55.3	15.5	10.1 to 21.0	3.6	0.1 to 7.1
Non-cage houses						
Nest boxes	1.7	0.0 to 4.9	0.0	N.C.	15	0.0 to 30.6
Drinkers	5.0	0.0 to 10.5	0.0	N.C.	30	9.9 to 50.1
Feeders	9.5	2.0 to 16.9	0.0	N.C.	0.0	N.C.
Slats	3.3	0.0 to 7.9	3.3	0 to 10.0	0.0	N.C.
Scratching	43.5	30.1 to 56.0	0.0	N.C.	0.0	N.C.
Soil (range) ^a	52.5	37.2 to 67.1	2.5	0 to 6.7	45	N.C.

N.C., not calculated. ^aNote that soil prevalence is presented as mean (range).

Level of infection in flocks before treatment. Flocks in cage houses sampled in lay before the C&D had a median (dust and faeces, weighted) sample prevalence of 39.7% (interquartile range, 15.3 to 65.2%), and in non-cage houses the median (dust and faeces weighted) sample prevalence was 48.4% (interquartile range, 28.6 to 71.2%). These differences were not statistically significant (Wilcoxon $Z = -0.75$; $P = 0.454$). Flocks in houses that were later treated using the "in-house" method had a significantly higher level of infection (63.2%; interquartile range, 41.4 to 81.2%) than those treated with formalin

(44.3%; interquartile range, 20.7 to 65.9%) (Wilcoxon $Z = 2.0$; $P = 0.045$) or FGQ (GO) (25.9%; interquartile range, 13.7 to 43.9) (Wilcoxon $Z = 3.54$; $P < 0.001$). The percentage of cases where the follow-on flock was *Salmonella* free (i.e. *Salmonella* was eliminated from the building) was 7/13 (54%), 8/23 (35%) and 5/24 (21%) for houses treated with formalin 10%, FGQ (GO) and the in-house method, respectively.

Comparison of the efficacy of each treatment on the levels of infection of follow-on flock. The level of infection of the




Figure 1. Results of the C&D procedure of cage houses using three disinfection methods: an "in-house" method ($n = 18$), formaldehyde glutaraldehydequarternary ammonium (FGQ) at the GO rate ($n = 19$) and using 10% formalin dilution ($n = 11$).



Figure 2. Results of the cleaning and disinfection of non-cage houses using an “in-house” disinfection method (n = 6), and formaldehyde glutaraldehyde/quarternary ammonium (FGQ) at the GO rate (n = 4).

follow-on flocks, as judged from the percentage of samples that were positive, increased in 6/18 (33.3%), 3/19 (15.8%) and 0/11 (0%) of cases using the “in-house” method, FGQ (GO), and formalin 10%, respectively. In all remaining cases there was a reduction of the levels of infection of the follow-on flock. This included all non-cage houses regardless of the treatment (Figure 3). The variation in the prevalence of positive samples from the laying flocks before and after C&D is presented in Table 3.

The median observed reductions were -71.0% (interquartile range, -95.9 to -14.2) and -97.3% (interquartile range, -100 to -79.4) for cage and non-cage houses, respectively ($Z = 2.10$, $P = 0.035$). Overall, a significantly greater reduction was observed only with formalin 10%, compared with the in-house programme ($Z = -2.64$, $P = 0.008$). In cage houses, the 10% formalin dilution method performed better than the FGQ ($Z = -2.13$, $P = 0.033$) and the “in-house” method




Figure 3. Change in the percentage of positive samples collected from the flocks before and after disinfection using each of the three methods.

Table 3. Change in the sample prevalence between Flock 1 (prior to treatment) and Flock 2 (after treatment)

Type of house	Method	Median difference	75% interquartile range
Cage	“In-house”	-47.2	-72.6 to 3.92
	FGQ	-70.0	-98.9 to -14.1
	Formalin	-92.7	-100 to -85.4
Non-cage	“In-house”	-91.9	-98.6 to -59.8
	FGQ	-100	-100 to -100
	Formalin	-74.0	-53.7 to -33.4

($Z = -3.41$, $P < 0.001$). In non-cage houses, the FGQ method performed better than the “in-house” method, but the difference was only borderline significant ($Z = -1.80$, $P = 0.07$). No statistical comparisons with the 10% formalin dilution method were attempted in non-cage houses due to the small number of houses treated with formalin ($n = 2$).

Significance of post-C&D sampling results. A negative post-C&D sampling result resulted in a *Salmonella*-negative follow-on flock in 57% of cases (Table 4). There was a linear trend towards an increased risk in carry-over for houses with higher levels of *Salmonella* after cleaning and disinfection (χ^2 for linear trend 8.57; $P = 0.0031$), although houses with a negative C&D result still had a 43% probability of carry-over.

In the case of free-range houses, there was evidence of contamination in the soil of the paddocks at the time of the post C&D visit in 6/9 cases. None of the follow-on flocks placed in the three houses with negative paddocks tested positive, compared with 4/6 of those with evidence of contamination of the paddocks.

Discussion

The perpetuation of *S. Enteritidis* infection in laying flocks is commonly related to the presence of a contaminated house environment (van de Giessen *et al.*, 1994; Wales *et al.*, 2007; Carrique-Mas *et al.*, 2008b). Therefore adequate C&D is regarded as a crucial step in eliminating infection from the houses. The present study demonstrated the variable impact of different C&D regimes on *Salmonella*-positive laying houses. An ideal study design to investigate this in the field would be a randomized controlled trial with assessment of compliance but this was not possible because of cost issues. Therefore, we need to accept the limitations and possible biases of this descriptive study.

Table 4. Relationship between post C&D sampling results and follow-on flock

% positive samples	Follow-on flock <i>Salmonella</i> status		
	Positive	Total	% <i>Salmonella</i> positive
Negative	6	14	42.9
>0 to 5%	8	14	57.1
>5 to 11.3%	8	9	88.9
>11.3 to 25%	11	12	91.7
>25%	10	11	90.9
Total	43	60	

The best results were achieved using 10% formalin applied by a specialist contractor, and the second best results were obtained using a blended disinfectant that also contained formaldehyde, in combination with glutaraldehyde and quarternary ammonium compounds. Other disinfection methods did not perform as well, although the diversity of treatments does not permit conclusions regarding any particular product to be made. The efficacy of the application of the 10% formalin treatment by spray was reflected not only in the lower recovery of *Salmonella* from the treated surfaces in the house, but in lower rate of infected follow-on flocks and a reduction in the levels of infection in houses where an infected flock was present. It is important to emphasize that this is different from the application of formalin by fogging, a less intensive treatment that is common in the poultry industry. Fogging is regarded as not very efficient on its own due to the considerable shadowing effect associated with the complicated structures of the house. For health and safety reasons, it is better that formalin is applied by a specialist contractor. This is likely to represent an additional expense, which may largely be compensated by the improved performance of this disinfection method and improved health and production in the subsequent flocks.

There is potentially a source of bias in the study resulting from the uneven distribution of the level of contamination in the houses investigated by type of treatment, since the allocation of C&D treatments was not under our control. The houses that were treated with the “in-house” method had also contained flocks with the overall highest levels of infection. However, flocks that were treated with formalin had a greater initial level of infection than those treated with the FGQ product, and, even so, the disinfection effect was greater using the 10% formalin dilution. In one of the two non-cage houses (a barn), the formalin treatment did not succeed. This was the first house to be treated in the study. The reason for this failure was a very poor standard of cleaning before the application of the disinfectant, being particularly bad in nest boxes and drinker cups, which were largely missed by the cleaning team. In addition the house was quickly re-contaminated by mouse faeces.

Laying houses are notoriously difficult to clean thoroughly because of their intrinsically complicated structures, which are even more complex in the case of cage laying houses (Wales *et al.*, 2006a). Access to cage interiors, feeders, egg belts, and so forth is very difficult unless a great deal of effort and time is invested. It seems that in these circumstances a large amount of residual organic matter is expected after a standard disinfection procedure. Our results are consistent with previous results where formalin has also been shown to perform better than any other disinfectant. This has been confirmed using laboratory models using *Salmonella*-spiked surfaces (Berchieri & Barrow, 1996; Gradel *et al.*, 2004) but also in field studies (Davies & Wray, 1995; Rose *et al.*, 2000).

The general trend in the houses investigated was a reduction of the levels of infection of the follow-on flocks regardless of the treatment used, although there was a poor predictive value of a negative C&D result. Only in 42% of cases where a negative post-C&D result was obtained was there evidence of clearance of infection of the follow-on flock. This is likely to be a consequence of the presence of other major sources of infection to the

flocks, notably rodents (Carrique-Mas *et al.*, 2009). Likewise, a positive result after C&D, particularly at a low level, does not necessarily lead to infection of the new flock, as has been reported previously (Davies & Wray, 1996). It is possible in some cases that our involvement in the study may also have lead to a more pro-active attitude by the farmer in terms of upgraded rodent control and biosecurity measures. In houses where relatively small numbers of rodents were present, an upgrade of baiting procedures may have been sufficient to reduce the challenge to the newly placed flocks.

An important observation in the present study is the widespread lack of knowledge by the farmers of the appropriate concentrations of disinfectants. In the UK, activity against *Salmonella* is evaluated and the efficacy reported in the GO concentration. The main confusion arising from the use of disinfectants appears to be related to their reconstitution at a lower than optimal concentration, as is recommended for other uses. It would be preferable if a suitable concentration for disinfection of *Salmonella* in biofilms on soiled surfaces could be defined for all products so that operators could obtain clear guidance on appropriate disinfection.

Paddocks adjacent to free-range houses are rarely included in the C&D programme. Although in many cases paddocks have been reported to carry *Salmonella*, they are considered to be less of a risk for birds than residual contamination in the house, since carry-over has been rare provided that the house has been correctly disinfected (Davies & Breslin, 2003).

The current results are particularly relevant in the context of the enforcement of restriction on the sale of fresh eggs from flocks infected with *S. Enteritidis* and/or *S. Typhimurium*. The present study provides evidence that the use of the 10% formalin dilution is particularly useful in the decontamination of infected cage laying flocks, although in most cases this intervention has to be concomitant with rodent control to reduce the chances of carry-over of infection.

Acknowledgements

The authors wish to thank all participating farmers for their support. This work was funded through DEFRA of the Government of the United Kingdom (Project OZ0325). The authors are also grateful to Dr Sarah Evans for comments and improvements to the original manuscript.

References

- Anonymous. (2006). Reducing *Salmonella* in European egg-laying hens: EU targets now set. *Eurosurveillance*, 11, E060810 3.
- Berchieri, A., Jr. & Barrow, P.A. (1996). The antibacterial effects for *Salmonella* Enteritidis phage type 4 of different chemical disinfectants and cleaning agents tested under different conditions. *Avian Pathology*, 25, 663–673.
- Carrique-Mas, J.J., Breslin, M., Sayers, A.R., McLaren, I., Arnold, M. & Davies, R. (2008a). Comparison of environmental sampling methods for detecting *Salmonella* in commercial laying flocks in the UK. *Letters in Applied Microbiology*, 47, 514–519.
- Carrique-Mas, J.J., Breslin, M., Snow, L., Arnold, M.E., Wales, A., McLaren, I. & Davies, R.H. (2008b). Observations related to the *Salmonella* EU layer baseline survey in the UK: Follow-up of positive flocks and sensitivity issues. *Epidemiology and Infection*, 47, 514–519.
- Carrique-Mas, J.J., Breslin, M., Snow, L., McLaren, I., Sayers, A.R. & Davies, R.H. (2009). Persistence and clearance of different *Salmonella* serovars in buildings housing laying hens. *Epidemiology and Infection*, 137, 837–846.
- Davies, R. & Breslin, M. (2003). Observations on *Salmonella* contamination of commercial laying farms before and after cleaning and disinfection. *Veterinary Record*, 152, 283–287.
- Davies, R.H. & Wray, C. (1995). Observations on disinfection regimens used on *Salmonella enteritidis* infected poultry units. *Poultry Science*, 74, 638–647.
- Davies, R.H. & Wray, C. (1996). Persistence of *Salmonella enteritidis* in poultry units and poultry food. *British Poultry Science*, 37, 589–596.
- de Jong, B. & Ekdahl, K. (2006). Human salmonellosis in travellers is highly correlated to the prevalence of *Salmonella* in laying hen flocks. *Eurosurveillance*, 11, E060706 1.
- Department for Environment, Food and Rural Affairs. (2009). *Disinfectants Approved for use in England and Wales*. Available online at http://disinfectants.defra.gov.uk/Default.aspx?Location=None&module=ApprovalsList_SI (accessed 9 October 2008).
- Gillespie, I.A., O'Brien, S.J., Adak, G.K., Ward, L.R. & Smith, H.R. (2005). Foodborne general outbreaks of *Salmonella* Enteritidis phage type 4 infection, England and Wales, 1992–2002: where are the risks? *Epidemiology and Infection*, 133, 795–801.
- Gradel, K.O., Sayers, A.R. & Davies, R.H. (2004). Surface disinfection tests with *Salmonella* and a putative indicator bacterium, mimicking worst-case scenarios in poultry houses. *Poultry Science*, 83, 1636–1643.
- McDonnell, G. & Russell, A.D. (1999). Antiseptics and disinfectants: activity, action, and resistance. *Clinical Microbiology Reviews*, 12, 147–179.
- Rose, N., Beaudreau, F., Drouin, P., Toux, J.Y., Rose, V. & Colin, P. (2000). Risk factors for *Salmonella* persistence after cleansing and disinfection in French broiler-chicken houses. *Preventive Veterinary Medicine*, 44, 9–20.
- van de Giessen, A.W., Ament, A.J. & Notermans, S.H. (1994). Intervention strategies for *Salmonella enteritidis* in poultry flocks: a basic approach. *International Journal of Food Microbiology*, 21, 145–154.
- Wales, A., Breslin, M. & Davies, R. (2006a). Assessment of cleaning and disinfection in *Salmonella*-contaminated poultry layer houses using qualitative and semi-quantitative culture techniques. *Veterinary Microbiology*, 116, 283–293.
- Wales, A., Breslin, M. & Davies, R. (2006b). Semiquantitative assessment of the distribution of *Salmonella* in the environment of caged layer flocks. *Journal of Applied Microbiology*, 101, 309–318.
- Wales, A., Breslin, M., Carter, B., Sayers, R. & Davies, R. (2007). A longitudinal study of environmental *Salmonella* contamination in caged and free-range layer flocks. *Avian Pathology*, 36, 187–197.
- World Health Organization. (2005) *Drug-resistant Salmonella Factsheet No. 139*. Available online at <http://www.who.int/mediacentre/factsheets/fs139/en/> (accessed 5 November 2008).