



Effect of gamma irradiation and storage time on microbial growth and physicochemical characteristics of pumpkin (*Cucurbita Moschata* Duchesne ex Poiret) puree

María F Gliemmo^{1,2}, María E Latorre^{1,2}, Patricia Narvaiz³,
Carmen A Campos^{1,2} and Lía N Gerschenson^{1,2}

Abstract

The effect of gamma irradiation (0–2 kGy) and storage time (0–28 days) on microbial growth and physicochemical characteristics of a packed pumpkin puree was studied. For that purpose, a factorial design was applied. The puree contained potassium sorbate, glucose and vanillin was stored at 25 °C. Gamma irradiation diminished and storage time increased microbial growth. A synergistic effect between both variables on microbial growth was observed. Storage time decreased pH and color of purees. Sorbate content decreased with storage time and gamma irradiation. Mathematical models of microbial growth generated by the factorial design allowed estimating that a puree absorbing 1.63 kGy would have a shelf-life of 4 days. In order to improve this time, some changes in the applied hurdles were assayed. These included a thermal treatment before irradiation, a reduction of irradiation dose to 0.75 kGy and a decrease in storage temperature at 20 °C. As a result, the shelf-life of purees increased to 28 days.

Keywords

Gamma irradiation, pumpkin, microbial spoilage, color stability

Date received: 1 June 2012; revised: 29 November 2012

INTRODUCTION

Pumpkin (*Cucurbita moschata* Duchesne ex Poiret) is a vegetable that comes from tropical and subtropical zones like Mexico and South America. Its production and consumption have significantly grown in the last decades in Argentina. It is cheap, available throughout the year and it is a good source of nutrients, such as carotenoids, potassium ion and vitamins. It has low calories and is also an important source of fiber. Pumpkin has numerous culinary uses either alone or as an ingredient in pies, soups, stews and breads. Its consumption as a puree is highly usual, especially for

kids and immunosuppressed people (de Escalada Pla et al., 2007; González et al., 2001).

Pumpkin puree is very sensitive to microbial spoilage due to its high water activity and its pH close to neutrality. In addition, physiological and biochemical changes take place even under refrigerated conditions. Its shelf-life is likely to be improved by applying

¹Departamento de Industrias, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Ciudad Universitaria, Buenos Aires, Argentina

²Consejo Nacional de Investigaciones Científicas y Técnicas de la República, Argentina

³Food Irradiation Section, Radiation Technology and Applications, Ezeiza Atomic Center, National Atomic Energy Commission, Argentina

Corresponding author:

Carmen A Campos, Departamento de Industrias, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Ciudad Universitaria, Int. Güiraldes s/n (1428), Buenos Aires, Argentina.

Email: carmen@di.fcen.uba.ar

minimal preservation processes based on the 'hurdle' concept. The combination of different preservation factors such as water activity reduction through solutes addition, pH depression, addition of lipophilic weak acids and the application of thermal and nonthermal technologies for microbial inactivation like gamma irradiation, have been used to increase the shelf-life of many minimally processed vegetables, such as grated carrots (Lacroix and Lafortune, 2004), pumpkin puree (Dutta et al., 2006), mango puree and pulp and orange juice (Noomhorm et al., 1998; Thakur and Arya, 1993; Youssef et al., 2002).

Gamma irradiation is a safe and effective emerging technology for food preservation. It can penetrate into the food and interact with microorganisms, disrupting its genetic material. The use of irradiation was proposed as the safest method for reducing the risk of foodborne pathogens, such as *E. coli* in fresh fruits and vegetables (Eyre, 2008). However, the interaction between irradiation and food components could cause the development of undesirable sensory and chemical changes in some foods (Thakur and Singh, 1995). Moreover, the resistance of a microbial cell to irradiation depends on its characteristics, process parameters and the extracellular environmental, i.e. temperature, water activity, pH, presence of oxygen and sorbates (Lacroix and Lafortune, 2004; Mañas and Pagan, 2005). The addition of sorbates before irradiation prevents off-odor development, improves the sensory quality and extends food shelf-life (Thakur and Arya, 1993; Thakur and Singh, 1995). However, sorbic acid can suffer radiolytic degradation depending on the composition of the system (Thakur et al., 1990; Singh et al., 1991).

The color of pumpkin puree is a parameter involved in its quality. β -Carotene is responsible for the orange color of pumpkin (González et al., 2001) and it has beneficial effects on health. However, its stability can be affected by isomerization and oxidation reactions during processing and storage promoting nutritional losses and undesirable changes in sensory properties (Ahmed et al., 2002; Dutta et al., 2006; Melendez-Martinez et al., 2004).

The effect of gamma irradiation and storage on the quality of purees and pulps has only been reported for tropical fruits such as mango and papaya, and no information is available on pumpkin puree. The objective of this work was to evaluate the effect of gamma irradiation and storage on the microbial growth and physicochemical characteristics of pumpkin puree.

MATERIALS AND METHODS

To perform the study, in a first step, a two-level, full factorial design was applied. The evolution of

indigenous flora, pH, color and potassium sorbate (KS) content were measured in a packed pumpkin puree with a pH 5.00, stored at 25 °C. Taking into account the results that were obtained, some changes in the applied hurdles were assayed in a second step: the inclusion of a thermal treatment before irradiation, the reduction of the irradiation dose and the decrease in storage temperature.

Preparation of pumpkin purees

Fresh pumpkins (*Cucurbita moschata* Duchesne ex Poiret) were purchased from a local market (Buenos Aires, Argentina). They were washed thoroughly with soap and tap water and rinsed with chlorinated water (0.05 g/kg of NaOCl). The pumpkin's mesocarp was cut, in perpendicular direction to the axis, in 2-cm-thick slices. These slices were peeled off and 3-cm-diameter cylinders (\approx 12.5 g) were cut from each slice. They were exposed to steam at atmospheric pressure for 12 min, cooled by immersion in chlorinated water for 3 min and dried with tissue paper to remove excess water. After that, the cylinders were placed into a 500 mL beaker containing glucose (10 g/100 g of system); KS (0.12 g/100 g of system) and vanillin (0.050 g/100 g of system) and the ingredients were homogenized with a Sorvall Omni Mixer (OMNI Corporation, Connecticut, USA). The addition of glucose and vanillin was done following the results from a previous study on the preservation of pumpkin puree by hurdle technology (Gliemmo et al., 2010). The pH was adjusted to 5.00 by the addition of citric acid. The puree (50 g) was packed in polyvinyl chloride-polyvinylidene chloride copolymer bags (BB4L, Cryovac Inc., Sealed Air Corporation; thickness: 59 μ m; oxygen permeability at 23 °C: 30.4 cm³/m².d.atm). The choice of depressing the pH to 5.0 by the addition of citric acid and the use of bags with low permeability to oxygen was based on previous studies suggesting that these conditions protect the color of puree (Gliemmo et al., 2009). Once bags were heat sealed, two treatments were applied: (i) irradiation and (ii) thermal treatment + irradiation.

The concentration of all the components were within the level admitted by the Argentine Food Code (2007) for low sugar products. Every ingredient used was reagent grade: KS and vanillin were from Sigma (St. Louis, Mo., USA); glucose and citric acid, from Anedra (Buenos Aires, Argentina).

Puree treatments

Irradiation and experimental design. The bags, placed in boxes, were exposed to gamma irradiation from a 550,000 Ci Cobalt 60 source (Semi-industrial facility,

Table 1. Two-level full factorial design used to evaluate the effects of gamma irradiation and storage time on microbial growth and physicochemical characteristics of pumpkin puree

Run	Independent variable			
	Gamma irradiation dose		Storage time	
	Uncoded value (kGy)	Coded value	Uncoded value (d)	Coded value
1	0.00	-1	0	-1
2	2.00	1	0	-1
3	0.00	-1	28	1
4	2.00	1	28	1

Ezeiza Atomic Centre, Argentina) at 25 °C. The bags received a dose of 2.00 kGy that was confirmed by Ag₂Cr₂O₇ dosimetry under the standards of the American Society for Testing and Materials (ASTM EL1401-91e1, 1995). The dose rate was 8.61 kGy/h.

The boxes were rotated 180° during the process to ensure dose uniformity (maximum dose/minimum dose). The uniformity attained was 1.08. After irradiation, the bags were stored at 25 ± 1 °C in a forced convection constant temperature chamber. The puree was sampled at selected times for microbiological and physicochemical analyses.

To evaluate the effect of the applied gamma irradiation doses (0–2.00 kGy) and storage time (0–28 days) on the shelf-life of the puree, a 2² full factorial design was performed with both factors at the indicated levels. The complete design consisted of 4 experimental treatments (Table 1) with 3 replicates each, resulting in a total of 12 experimental units. Results were analyzed using Statgraphics Plus for Windows, version 5.1 (Manugistics, Inc., Rockville, Maryland, USA).

Thermal treatment and irradiation. Nine sealed bags were heated in a bath at 50 ± 1 °C for 15 min before the irradiation treatment. Immediately after that, six bags received a dose of 0.75 kGy with a dose rate of 9.72 kGy/h using the above mentioned methodology. The dose uniformity was 1.06 kGy. Three nonirradiated bags were used as control.

All thermal treated bags were stored at 20 ± 1 °C in a forced convection constant temperature chamber. The puree was sampled at the initial time and after 28 days of storage, in triplicate. Microbiological and physicochemical analyses were performed.

Microbiological analysis

The puree samples (5 g) were aseptically taken from each bag and placed in a sterile bag containing 45 mL of 0.1% w/v peptone water. The content of the bags

were blended for 1 min using a stomacher device. Then serial dilutions were prepared in 0.1% w/v of peptone water to determine the level of indigenous flora. The aerobic and anaerobic mesophilic bacteria were investigated in PCA (plate count agar). The anaerobic bacteria incubation was performed in an anaerobiosis jar with an anaerobic atmosphere generator (BioMérieux® S.A., France). Lactic acid and coliform bacteria were investigated in MRS (deMan, Rogosa and Sharpe) and VRBL (violet red bile agar), respectively. The plates were incubated at 30 °C for 48 h. The yeasts and molds were determined by surface plate count on Sabouraud agar after 5 days of incubation at 25 °C. All the culture media used were from Biokar (Biokar Diagnostics, Beauvais, France).

Physicochemical analysis

The initial soluble solids content was determined for each sample with a hand-held sugar refractometer (Westover, USA) and results were expressed in °Brix.

The pH of each sample was determined by inserting an electrode (Solution Analyzer 5800-05, Cole-Parmer, Chicago, Ill., USA) into the puree sample.

The potassium sorbate content was measured in duplicate according to the AOAC (1995) oxidation method. In a previous study, this AOAC method had a 3.4% variance coefficient (Campos et al., 1991).

The color measurement was performed using a colorimeter (Minolta Co. Ltd., Osaka, Japan) with illuminant D65 and an observer at 10°. For that purpose, each puree pouch was placed onto a white tile and the three-color coordinates *L** (lightness, 100 for white and 0 for black), *a** (greenness to redness, -80 for green and 100 for red) and *b** (blueness to yellowness, -80 for blue and 70 for yellow) were read. The color measurements were taken in triplicate samples. Each bag was measured in three different places and average values were calculated. Pumpkin puree color degradation was also expressed by the

total color difference (ΔE^*). This parameter is expressed by the following equation

$$\Delta E^* = \sqrt{(L_0^* - L^*)^2 + (a_0^* - a^*)^2 + (b_0^* - b^*)^2} \quad (1)$$

where L_0^* , a_0^* and b_0^* represent the averages of zero time readings, and L^* , a^* and b^* , represent the instantaneous individual readings during storage. For thermal treated purees, zero time readings corresponded to non-irradiated samples after thermal treatment.

Sensory evaluation of color difference

The Difference from Control Test was applied to establish from which ΔE^* value an observer can detect color differences from a sensory viewpoint. The evaluation of sensory color differences along storage was done using nonirradiated purees and 2 kGy irradiated purees. Control puree was a frozen puree sample, stored at -18.0°C , thawed previous to evaluation (Meilgaard et al., 1991). Previous analysis revealed that freezing did not alter the color of the control sample. A group of six trained assessors with normal color vision and experience in visual assessment of color took part in the study. The samples were placed on coded watch glasses (10 g). The assessors received a puree sample and the known control. They were asked to evaluate the relative color difference using a numerical scale (1 = no difference from the control; 7 = large difference from the control). They also evaluated a blind control placed as an unknown sample. The samples and the blind control were distributed in randomized order together with a known control. The evaluations were made in three sessions. In parallel, the color of samples was measured with a colorimeter as it was previously mentioned.

DATA ANALYSIS

Experimental data obtained from the full factorial design was subjected to a multiple regression analysis to fit the following first-order regression model

$$Y = \alpha_0 + \alpha_1 x_1 + \alpha_2 x_2 + \alpha_3 x_1 x_2 + \varepsilon \quad (2)$$

where Y are log microorganism populations, pH values and residual KS content; $\alpha_{0,1,2,3}$ are the regression coefficients for the intercept, linear and interaction coefficients, respectively; $x_{1,2}$ are the independent variables (x_1 , irradiation dose and x_2 , time) in coded units and ε is the error term.

The adequacy of the regression model generated by the factorial design was examined by analysis of variance (ANOVA) at 5% significance level, correlation coefficients (R^2) and the absolute average deviation

(AAD) (Baş and Boyaci, 2007). Also, ANOVA and p -value were used to evaluate the significance of the linear and interaction terms of each model.

Significant differences of the three-color coordinates and ΔE^* for nonthermally treated purees, and quality parameters of thermally treated purees were evaluated through an ANOVA and the least significant difference (LSD) test.

To analyze the data from sensory evaluation of color, an ANOVA for a randomized (complete) block design of two factors without interaction was used (Meilgaard et al., 1991). The factors were the subjects and treatments (none and 2 kGy irradiation treatments).

In all cases, statistical significance was evaluated at a 5% level ($\alpha=0.05$) and the analyses were performed using Statgraphics Plus for Windows, version 5.1 (Manugistics, Inc., Rockville, Maryland, USA).

RESULTS AND DISCUSSION

Since the total soluble solids content of pumpkin puree after steam treatment was 10.5 ± 0.5 °Brix, water activity was 0.994 and the pH value was 6.10, microbial spoilage was prone to occur. Thus, processing should be a must in order to extend shelf-life of the product.

IRRADIATION TREATMENT

Regression analysis

The first-order regression model for each response was fitted using the experimental data. The correlation coefficients, AAD and p -values are shown in Table 2. All R^2 values were within the range 0.80–0.99 indicating a good correlation between the observed and the predicted values of responses. This fact together with the small AAD values indicates that the model gives a reasonably good estimate of responses in the studied range.

Microbiological analysis

Coliform bacteria growth was not detected in any sample during storage suggesting that processing conditions were hygienic.

The probability (p) values of terms of regression model indicate that gamma irradiation and storage time significantly affected microbial growth at a 95% confidence level (Table 2). Gamma irradiation diminished whereas time increased microbial growth as it was demonstrated by the signs of regression coefficients in Table 2. An increase in approximately 2.5 log CFU/g was determined for all microorganisms in the nonirradiated purees after 28 days of storage (Figure 1). The application of gamma irradiation produced a decrease

Table 2. Probability (p) values, correlation coefficients (R^2), absolute average deviation (AAD) and coefficients of the regression models for microbial growth, potassium sorbate (KS) content and pH of pumpkin puree. Independent variables, x_1 : irradiation dose and x_2 : time. Coefficients, α_1 : for irradiation dose; α_2 : for time; α_3 : for interaction

Quality parameter	p -value			R^2	AAD	Coefficients of regression models			
	x_1	x_2	x_1x_2			α_0	α_1	α_2	α_3
Log CFU/g of yeasts and molds	0.0001	<0.0001	<0.0001	0.99	0.94	5.02	-0.49	1.94	0.63
Log CFU/g of aerobic bacteria	0.0010	<0.0001	0.0004	0.98	3.2	5.09	-0.53	2.00	0.61
Log CFU/g of anaerobic bacteria	0.0003	<0.0001	<0.0001	0.99	8.6	5.15	0.44	2.07	0.68
Log CFU/g of lactic acid bacteria	0.0001	<0.0001	<0.0001	0.99	3.3	5.44	-0.55	1.81	0.61
pH	0.8145	<0.0001	0.1280	0.99	0.32	4.685	0.002	-0.395	0.011
KS content (% w/w)	0.0225	0.0023	0.9009	0.80	2.82	0.1070	-0.0032	-0.0049	-0.0001




Figure 1. Estimated response surface graphs for the combined effect of time and the radiation dose in coded units on log CFU/g of indigenous flora of pumpkin purees.

in 2.5 log cycles of populations at initial time and did not have effect after 28 days of storage (Figure 1).

A significant interaction between irradiation and storage time on microbial growth was observed. Positive interaction coefficients suggest a synergistic effect (Table 2). This implies that the highest increase in population (5 log CFU/g) was observed in irradiated puree after 28 days of storage (Figure 1).

It must be remarked that after 5 days of storage, nonirradiated bags were inflated due to microbial metabolism whereas the swelling appearance of irradiated bags was delayed to 15 days.

Visual examination of Sabouraud plates showed that the microbial population was mostly constituted by

yeasts, suggesting that they are more radioresistant than molds. Similar results were found by Noomhorm et al. (1998), Prakash et al. (2000), Jiang et al. (2010) and Rivera et al. (2011).

It is known that only the undissociated form of sorbic acid has preservative action (Sofos, 2000). Since the pKa of sorbic acid is 4.75 (Mihyar et al., 1997) at pH 5.00, only 29% of the preservative is in the effective form, which would correspond to 0.039% w/w of sorbic acid in the studied purees. Probably this concentration was not enough to inhibit microbial growth since levels greater than 0.050% w/w of sorbic acid were reported as the minimum inhibitory concentrations for some spoilage yeasts according to

Praphailong and Fleet (1997). The mentioned facts stressed the need to use additional hurdles to minimize microbial growth and extend the shelf-life of the purees.

The resistance of a microbial cell to irradiation depends on several factors such as microbial characteristics, environmental factors and absorbed irradiation dose (Mañas and Pagán, 2005). The existence of a shoulder in typical radiation dose–survival curves is reported and it is related to repairable cellular damage in the low dose range (van Gerwen et al., 1999; Mañas and Pagán, 2005). In the present study, although absorbed irradiation doses diminished microbial population, they were not high enough to diminish growth to acceptable levels during storage.

The Argentine Food Code (2007) establishes counts of aerobic mesophilic bacteria $\leq 5 \times 10^4$ CFU/g, and of yeasts and molds $\leq 10^3$ CFU/g for ready to use dietetic foods. In order to avoid exceeding these values, irradiation dose (D) and storage time (t) were calculated using the following regression models expressed in coded units, and taking into account only the significant coefficients (Table 2):

$$\begin{aligned} \text{Log CFU/g aerobic bacteria} = & 5.09 - 0.53[D] \\ & + 2.00[t] + 0.61[D][t] \end{aligned} \quad (3)$$

$$\begin{aligned} \text{Log CFU/g yeasts and molds} = & 5.02 - 0.49[D] \\ & + 1.94[t] + 0.63[D][t] \end{aligned} \quad (4)$$

Results indicated that a population of 5×10^4 CFU/g of aerobic mesophilic bacteria would be found after 12 days of storage in samples irradiated at 1.14 kGy, and 10^3 CFU/g of yeasts and molds would be found after 4 days of storage in a puree irradiated at 1.63 kGy. Therefore, from the microbiological point of view, the puree would be acceptable with an absorbed dose of 1.63 kGy up to the 4th storage day.

It is important to emphasize that the studied purees were safe from *Clostridium botulinum* growth and its toxin production because they were packed in aerobic atmosphere and contained KS, an effective preservative to inhibit *C. botulinum* growth (Lund et al., 1987; Seward et al., 1982; Thakur and Singh, 1995).

Physicochemical analysis: pH evolution

The storage time significantly decreased the pH of the purees whereas the effect of irradiation was not significant (Table 2). The minimum value reached was 4.28 and it corresponded to the nonirradiated system after 28 days of storage (data not shown). The observed pH

decrease could be related to lactic acid bacteria development (Figure 1, panel D). Similar behaviour was observed by Noomhorm et al. (1998) in mango puree stored at room temperature.

Physicochemical analysis: Potassium sorbate stability

Sorbate content significantly decreased with storage time and gamma irradiation as indicated by the negative sign of regression coefficients in Table 2. No significant interaction was observed between both variables on KS stability. A KS level of 0.11% w/w was reached in nonirradiated puree after 28 days of storage, in samples irradiated at initial time, while the level was 0.10% w/w in samples irradiated after storage.

Sorbic acid is prone to degrade through reactions of autoxidation and radiolysis. It is more susceptible to radiolytic degradation at acidic pH, and the degradation increases with the irradiation dose. The results obtained here suggest that KS degradation occurred by means of both reactions.

The rate of radiolytic degradation is controlled by diffusivity of the primary products formed by the radiolysis of water which are hydrated electrons, hydroxyl radicals and hydrogen atoms (Thakur et al., 1990). In moist systems, hydroxyl radicals are most reactive towards unsaturated compounds with conjugated polyene chains, but ionized carboxyl groups are relatively unreactive (Thakur et al., 1990). Although sorbic acid structure is reactive towards hydroxyl radicals, in the studied purees the percentage of its undissociated form is low, as explained above. Also, KS could be protected from radiolytic degradation by compounds such as pectins, polysaccharides and vanillin due to their competition to react with radiolytic products of water (Thakur et al., 1990). These facts would explain that the greatest loss of KS was of only 16.6%, corresponding to 0.10 g/100 g of residual KS, and it was found in the irradiated puree after 28 days of storage.

It must be stressed that KS degradation occurs via oxidative mechanisms and could generate brown and red pigments, which would therefore interfere in the color evaluation (Gliemmo et al., 2009).

Physicochemical analysis: Color stability

The signs of a^* and b^* values kept positive during the experiment showing a correlation with red and yellow colors, respectively (Table 3). Total color difference (ΔE^*) increased significantly with time, independently of irradiation. This increase is related to the decrease in L^* observed after 28 days of storage (Table 3).

Table 3. Three color coordinates (a^* , b^* and L^*) and total color difference (ΔE^*) of nonthermally treated purees

Pumpkin puree					
Gamma irradiation dose (kGy)	Storage time (d)	a^*	b^*	L^*	ΔE^*
0.00	0	25.2 ± 0.2 a	53 ± 1 a	43.9 ± 0.5 a	0.9 ± 0.3 a
0.00	28	24.1 ± 0.5 a,b	52 ± 4 a	40.3 ± 1.2 b	5.2 ± 1.6 b,c
2.00	0	24.8 ± 0.9 a	54 ± 2 a	43.1 ± 0.8 a	2.1 ± 0.9 a,b
2.00	28	22.7 ± 1.0 b	50 ± 6 a	40.4 ± 0.7 b	6.8 ± 3.8 c

Errors represent standard deviation.

Values in the same column followed by the same letter are not significantly different ($p > 0.05$).

The maximum ΔE^* calculated was 6.8 in irradiated purees at 28 days of storage. The color difference evaluation by sensory analysis allowed determining that from a ΔE^* value of 6.00, differences of color between nonirradiated purees, 2 kGy irradiated purees and control puree ($p = 0.67$) were perceived by the naked eye. This trend was observed at 14 days of storage.

The estimated ΔE^* of irradiated purees after 28 days of storage was coincident with the minimum of redness (Table 3). Loss of redness is related to the degradation of β -carotene (Dutta et al., 2006; Gliemmo et al., 2010; Kidmose et al., 2002). β -carotene may suffer radiolytic degradation through reaction with hydroxyl radicals (Thakur et al., 1990). During puree elaboration, thermal processing and homogenization improved β -carotene bioavailability since the cellulose structure of the plant cell is broken (Dutta et al., 2006). Therefore, the pigment would be more exposed to degradation reactions diminishing the food color during storage.

The decrease in lightness observed may be due to nonenzymatic browning reactions taking place together with oxidation and isomerization of β -carotene during storage time (Dutta et al., 2006). Moreover, the evaluation of redness loss may be influenced by brown and red pigments generated by the oxidative degradation of sorbate. On the other hand, irradiation may degrade soluble sugars of pumpkin puree and produce carbonylic compounds which may react with amino acids increasing browning development during storage (Thakur and Arya, 1993).

THERMAL AND IRRADIATION TREATMENTS

Since an estimated shelf-life of 4 days is not acceptable for commercial purposes and 1.63 kGy seems to be a relatively high dose leading to negative effects on the sensory characteristics (Thakur and Singh, 1995), the irradiation dose was decreased to 0.75 kGy. To compensate, a thermal treatment (50 °C for 15 min) was

applied before irradiation and the storage temperature was decreased to 20 °C.

Microbiological analysis

The counts of coliform, lactic acid bacteria and yeasts and molds were $< 9 \times 10^1$ CFU/g in the irradiated and nonirradiated purees at the initial time and after 28 days of storage.

It must be stressed that prior to thermal treatment puree samples did not contain coliforms. The counts of lactic acid bacteria and yeasts–molds were 3.9×10^3 and 4.2×10^3 CFU/g, respectively. As expected, the thermal treatment helped to reduce microbial flora.

The application of 0.75 kGy diminished aerobic and anaerobic bacteria growth in one log cycle at the initial time. Prior to thermal treatment aerobic bacterial counts diminished to less than one cycle while the anaerobic ones did not change significantly. After 28 days of storage, anaerobic bacteria growth decreased in one log cycle while aerobic bacteria growth did not change (Table 4). The latter values are within the microbiological limits established by the Argentine Food Code (2007) for ready to use dietetic foods.

It is well known that the use of gamma irradiation in combination with heating works synergistically to extend the shelf-life of irradiated foods (Aguirre et al., 2012). Heating sensitizes pathogens such as listeria as well as enzymes to radiation treatment. Furthermore, the irradiation doses required are reduced when the food is heated before irradiation, diminishing the negative effect of radiation on the sensory characteristics of food (Cumming and Blank, 2000; Noomhorm et al., 1998; Patterson, 2001; Parker et al., 2010; Thakur and Singh, 1995).

Physicochemical analysis

Values of pH were constant for 28 days of storage (Table 4). This trend is linked to the microbial stability previously mentioned.

Table 4. Microbial growth, pH, potassium sorbate loss (%) and total color difference (ΔE^*) of thermally treated purees

Quality parameter	Initial time		28 days of storage (at 20 °C)
	0 kGy	0.75 kGy	0.75 kGy
Microbial growth (log CFU/g):			
Aerobic bacteria	3.5 ± 0.2	1.9 ± 0.1 a	2.1 ± 0.3 a
Anaerobic bacteria	3.6 ± 0.1	2.3 ± 0.3	<1.9 ⁺
pH	4.96 ± 0.01 a	4.95 ± 0.01 a	4.93 ± 0.01 a
Potassium sorbate loss (%)	-	2.0 ± 1.0	13.0 ± 2.0
ΔE^*	1.2 ± 0.6	2.4 ± 0.9 a	2.4 ± 0.3 a

Errors represent standard deviation. Values in the same row followed by the same letter are not significantly different ($p > 0.05$).

⁺Estimated value.

In relation to preservative stability, neither thermal treatment nor the application of 0.75 kGy produced significant losses in KS concentration, but after 28 days of storage a loss of 13% was obtained indicating that only time affected KS stability (Table 4). However, this loss did not affect the microbiological stability of the thermally treated purees.

The ΔE^* increased with irradiation at the initial time reaching a value of 2.4. It remained constant in the irradiated puree along storage (Table 4). The mentioned increase was due to a slight enhancement in a^* and b^* values and a decrease in L^* . Since sensory color differences of purees were perceived from a ΔE^* value of 6.00, a color difference of 2.4 (Table 4) would not be perceived by the naked eye.

CONCLUSION

Indigenous flora of the nonthermally treated pumpkin purees increased during storage time and diminished with the application of gamma irradiation. The pH decrease was related to lactic acid bacteria growth. Total color difference increased with time, independently of irradiation. Considering microbial growth, a puree receiving an absorbed dose of 1.63 kGy and stored at 25 °C has a shelf-life of only 4 days.

On the contrary, in thermal treated purees, growth of coliform, lactic acid bacteria, yeasts and molds were not detected. Gamma irradiation diminished the growth of aerobic and anaerobic bacteria and kept counts within the microbiological limits established by the Argentine Food Code after 28 days of storage at 20 °C. The pH values kept constant for this period of time. Potassium sorbate losses were slight and did not affect microbiological stability of purees. The total color differences for these purees were smaller after 28 days of storage and would probably not be perceived

by the naked eye. According to the obtained results, heating before irradiation and storage at 20 °C allowed diminishing the irradiation dose necessary to assure microbial stability and, as a consequence, improved the color of the samples. The shelf-life of the puree could be extended to at least 28 days by the application of the proposed hurdles.

FUNDING

We acknowledge the financial support from Universidad de Buenos Aires, Consejo Nacional de Investigaciones Científicas y Técnicas de la República Argentina and Agencia Nacional de Investigaciones Científicas y Tecnológicas de la República Argentina.

ACKNOWLEDGMENT

Contributions from the Ezeiza Atomic Centre Irradiation Facility and High Radiation Doses Dosimetry Division staff are gratefully acknowledged.

REFERENCES

- Aguirre JS, Ordóñez JA and García de Fernando GD. (2012). A comparison of the effects of E-beam irradiation and heat treatment on the variability of *Bacillus cereus* inactivation and lag phase duration of surviving cells. *International Journal of Food Microbiology* 153(3): 444–452.
- Ahmed J, Shivhare US and Sandhu KS. (2002). Thermal degradation kinetics of carotenoids and visual color of papaya puree. *Journal of Food Science* 67(7): 2692–2695.
- AOAC. (1995). Official methods of analysis. Method 971.15. Sorbic acid in cheese. Oxidation method. In: Cunniff P (ed) *Official Methods of Analysis of AOAC International*, Chapter 47. Virginia: AOAC International, pp. 24–25.
- Argentine Food Code (2007). *Alimentos Dietéticos*. chapter XVII. Articles 1340 and 1371. Buenos Aires, Argentina: Secretaría de Agricultura, Ganadería, Pesca y Alimentos.

- ASTM EL1401-96e1 (1995). *Standard Practice for Use of a Dichromate Dosimetry System*. Philadelphia: American Society for Testing and Materials, pp. 411–416.
- Baş D and Boyaci IH. (2007). Modeling and optimization I: Usability of response surface methodology. *Journal of Food Engineering* 78: 836–845.
- Campos CA, Gerschenson LN, Alzamora SM and Chirife J. (1991). Determination of sorbic acid in raw beef: an improved procedure. *Journal of Food Science* 56(3): 863–866.
- Cumming R and Blank G. (2000). Irradiation. In: Robinson DS and Michael Eskin NA (eds) *Food Shelf Life Stability Chemical, Biochemical, and Microbiological Changes*, chapter 3. Boca Raton: CRC Press.
- de Escalada Pla MF, Ponce NM, Stortz CA, Gerschenson LN and Rojas AM. (2007). Composition and functional properties of enriched fiber products obtained from pumpkin (*Cucurbita moschata* Duchesne ex Poiret). *LWT – Food Science and Technology* 40: 1176–1185.
- Dutta DA, Dutta A, Raychaudhuri U and Chakraborty R. (2006). Rheological characteristics and thermal degradation kinetics of beta-carotene in pumpkin puree. *Journal of Food Engineering* 76: 538–546.
- Eyre C. (2008). Irradiation key to fruit and veg safety, study. Available at: <http://www.foodproductiondaily.com/Quality-Safety/Irradiation-key-to-fruit-and-veg-safety-study> (accessed 1 March 2011).
- Gliemmo MF, Latorre ME, Gerschenson LN and Campos CA. (2009). Color stability of pumpkin (*Cucurbita moschata*, duchesne ex poiret) puree during storage at room temperature: effect of pH, potassium sorbate, ascorbic acid and packaging material. *LWT-Food Science and Technology* 42: 196–201.
- Gliemmo MF, Latorre ME, Gerschenson LN and Campos CA. (2010). Effect of sweet solutes on the quality of a pumpkin puree (*Cucurbita moschata* Duchesne ex Poiret) preserved by the hurdle technology. *Journal of Food Processing and Preservation* 34(6): 1024–1040.
- González E, Montenegro MA, Nazareno MA and López de Mishima BA. (2001). Carotenoid composition and vitamin A value of an Argentinian squash (*Cucurbita moschata*). *Archivos Latinoamericanos de Nutricion* 51(4): 395–399.
- Jiang T, Luo S, Chen Q, Shen L and Ying T. (2010). Effect of integrated application of gamma irradiation and modified atmosphere packaging on physicochemical and microbiological properties of shiitake mushroom (*Lentinula edodes*). *Food Chemistry* 122: 761–767.
- Kidmose U, Edelenbos M, Norbaek R and Christensen LP. (2002). Colour stability in vegetables. In: MacDougall DB (ed.) *Colour in Food. Improving Quality*. Cambridge: Woodhead Publishing Ltd, pp. 179–218.
- Lacroix M and Lafortune R. (2004). Combined effects of gamma irradiation and modified atmosphere packaging on bacterial resistance in grated carrots (*Daucus carota*). *Radiation Physics and Chemistry* 71: 77–80.
- Lund BM, George SM and Franklin JG. (1987). Inhibition of Type A and Type B (proteolytic) *Clostridium botulinum* by sorbic acid. *Applied and Environmental Microbiology* 53(5): 935–941.
- Mañas P and Pagán R. (2005). Microbial inactivation by new technologies of food preservation. *Journal of Applied Microbiology* 98: 1387–1399.
- Meilgaard MC, Civille GV and Carr BT. (1991). *Sensory Evaluation Techniques*. Boca Raton: CRC Press, pp. 67–74.
- Melendez-Martinez AJ, Vicario IM and Heredia F. (2004). Estabilidad de los carotenoides en los alimentos. *Archivos Latinoamericanos de Nutricion* 54(2): 209–215.
- Mihyar GF, Yamani MI and Al-Sa'ed AK. (1997). Resistance of yeast flora of labaneh to potassium sorbate and sodium benzoate. *Journal of Dairy Science* 80(10): 2304–2309.
- Noomhorm A, Ilangantileke SG, Upadhyay IP, Karki DB and Apintanapong M. (1998). Use of irradiation in combination with preservation techniques to extend the shelf-life of tropical fruits and their products. *Combination Processes for Food Irradiation. Proceedings of Final Research Co-ordination meeting, Pretoria, South Africa. March*. Vienna, Austria: International Atomic Energy Agency, pp. 53–75.
- Parker TL, Esagro ST, Miller SA, Myers LE, Meister RA, Toshkov SA, et al. (2010). Development of an optimized papaya pulp nectar using a combination of irradiation and mild heat. *Food Chemistry* 118: 861–869.
- Patterson M. (2001). Tratamientos combinados implicando la irradiación de alimentos. In: Molins R (ed.) *Irradiación de alimentos-Principios y aplicaciones*. Zaragoza, Spain: Acribia, pp. 325–340.
- Prakash A, Guner AR, Caporoso F and Foley DM. (2000). Effects of low-dose gamma irradiation on the shelf life and quality characteristics of cut romaine lettuce packaged and stored under modified atmosphere. *Journal of Food Science* 65(3): 549–553.
- Praphailong W and Fleet GH. (1997). The effect of pH, sodium chloride, sucrose, sorbate and benzoate on the growth of food spoilage yeasts. *Food Microbiology* 14: 459–468.
- Rivera CS, Blanco D, Marco P, Oria R and Venturini ME. (2011). Effects of electron-beam irradiation on the shelf life, microbial populations and sensory characteristics of summer truffles (*Tuber aestivum*) packaged under modified atmospheres. *Food Microbiology* 28: 141–148.
- Seward RA, Deibel RH and Lindsay RC. (1982). Effects of potassium sorbate and other antibotulinal agents on germination and outgrowth of *Clostridium botulinum* Type E spores in microcultures. *Applied and Environmental Microbiology* 44(5): 1212–1221.
- Singh L, Murali HS and Sankaran R. (1991). Extension of shelf life of paneer by sorbic acid and irradiation. *Journal of Food Science of Technology* 28(6): 386–388.
- Sofos JN. (2000). Sorbic acid. In: Naidu AS (ed.) *Natural Food Antimicrobial Systems*. Boca Raton: CRC Press, pp. 637–660.
- Thakur BR and Arya SS. (1993). Effect of sorbic acid on irradiation-induced sensory and chemical changes in sweetened orange juice and mango pulp. *International Journal of Food Science and Technology* 28: 371–376.
- Thakur BR and Singh RK. (1995). Combination processes in food irradiation. *Trends in Food Science and Technology* 6: 7–11.

Thakur BR, Trehan IR and Arya SS. (1990). Radiolysis degradation of sorbic acid in isolated systems. *Journal of Food Science* 55(6): 1699–1710.

van Gerwen SJC, Rombouts FM, Van't Riet K and Zwietering MN. (1999). A data analysis on the irradiation parameter D_{10} for bacteria and spores under various conditions. *Journal of Food Protection* 62: 1024–1032.

Youssef BM, Asker AA, El-Samahy SK and Swailam HM. (2002). Combined effect of steaming and gamma irradiation on the quality of mango pulp stored at refrigerated temperature. *Food Research International* 35: 1–13.