

# Bioactive peptides from meat and their influence on human health

Baltić Ž. Milan<sup>1</sup>, Bošković Marija<sup>1</sup>, Ivanović Jelena<sup>1</sup>, Janjić Jelena<sup>1</sup>, Dokmanović Marija<sup>1</sup>, Marković Radmila<sup>1</sup>, Tatjana Baltić<sup>2</sup>

*Abstract:* Bioactive peptides are functional components, encrypted in the proteins and can be derived from food of plant and animal origin, including meat. After releasing during gastrointestinal digestion or food processing, these peptides exhibit many different effects on human body such as antioxidative, antimicrobial, antihypertensive, antithrombotic, cytomodulatory, immunomodulatory, anticancer, hypocholesterolemic and anti-obesity effects, which mainly depends on their structure and other properties. Considering bioactive activities of these peptides and their beneficial influence on health on one side, and millions of deaths caused by cancer, cardiovascular and other diseases associated with lifestyle on the other side, it is obvious that these peptides can be used for health promotion and disease risk reduction, especially because they have some advantages compared to synthetic drugs.

**Key words:** functional food, ACE inhibitory peptides, muscle proteins, antioxidant and antibacterial activity.

## Introduction

Cardiovascular diseases, cancer, diabetes and obesity are responsible for millions of deaths worldwide annually, and present increasing health and economic problem as well (Murray and Lopez, 1997; CDC, 2005; Ahhmed and Muguruma, 2010; DHHS, 2010; Weiss et al., 2010). These diseases and related conditions are also called chronic lifestyle-related diseases, because they are associated not only with heredity, but also with changes in lifestyle where diet plays important and in some cases crucial role (Murray and Lopez, 1997; Anand et al., 2008; Ahhmed and Muguruma, 2010; Decker and Park, 2010; Cam and de Mejia, 2012). This fact implies that food also may be used in the prevention, control or in some cases treatment of these diseases and this approach may present preventive health care strategy (Decker and Park, 2010). Consequently, as a response to this challenge food industry presented a new class of foods, so-called “functional foods”, and in Europe, these new food products have been labeled as “novel” foods and food ingredients (Diplock et al., 1999; Weiss et al., 2010; Olmedilla-Alonso et al., 2013). This food contains

components which exhibit a beneficial physiological effects on human health (Diplock et al., 1999; Weiss et al., 2010; Olmedilla-Alonso et al., 2013). New trend of promoting human health by using bioactive compounds is particularly interesting and presents a great challenge but at the same time opportunity for the meat industry, to improve the quality and image of meat (Jiménez Colmenero et al., 2010; Jiménez-Colmenero et al., 2012; Olmedilla-Alonso et al., 2013). Meat is important in human diet and had a great role in human evolution, especially in brain and intellectual development (Higgs, 2000; Baltić et al., 2002; Pereira and Vicente, 2013). Also, meat presents a valuable source of proteins, conjugated linoleic acid, antioxidants, vitamins such as riboflavin, niacin, vitamin B<sub>6</sub>, pantothenic and folic acid and numbers of minerals including iron, zinc, selenium and phosphorus (Jimenez-Colmenero et al., 2001; Baltić et al., 2002; Chan, 2004; Mulvihill, 2004; Biesalski, 2005; Arihara and Ohata, 2006; Descalzo and Sancho, 2008; Ahhmed and Muguruma, 2010; Decker and Park, 2010; Marković et al., 2010; Weiss et al., 2010; Toldra and Reig, 2011; Baltić et al., 2013; Pereira and Vicente, 2013). Meat proteins are not only important source of essential amino

**Acknowledgments:** This paper was supported by the Ministry of Education, Science and Technological Development of the Republic of Serbia, Project TR 31034.

<sup>1</sup>University of Belgrade, Faculty of Veterinary Medicine, Bulevar Oslobođenja 18, 11000 Belgrade, Republic of Serbia.

<sup>2</sup>Institute of Meat Hygiene and Technology, Kačanskog 13, 11000 Belgrade, Republic of Serbia.

**Corresponding author:** Baltić Ž. Milan, [baltic@vet.bg.ac.rs](mailto:baltic@vet.bg.ac.rs)

acids, but of bioactive peptides as well, and number of studies are focusing on the development of functional biopeptides from this source (Udenigwe and Ashton, 2013; Weiss et al., 2010; Young et al., 2013).

## Peptides - sources and production

Bioactive peptides are short, approximately 2–20 (in some cases this range can be extended) amino acids sequences with molecular masses of less than 6 kDa (Möller et al., 2008; Shahidi and Zhong, 2008; Di Bernardini et al., 2011). They are food derived components, and can be obtained from different plant and animal sources (Ryan et al., 2011). A great number of bioactive peptides are derived from plants such as soy, pulses (lentil, chickpea, pea and beans), oat, wheat, rice, maize, sunflower, hemp seed, pumpkin, canola, flaxseed and many others including mushrooms (Hartmann and Meisel, 2007; Möller et al., 2008; Rutherford-Markwick, 2012; Udenigwe and Aluko, 2012; Saavedra et al., 2013). Although most peptides derived from animal sources are generated from milk and milk-based products proteins, peptides also were isolated from eggs, bovine blood, collagen, gelatin, various fish species including tuna, sardine, herring, salmon, bonito and from marine organisms (Möller et al., 2008; Shahidi and Zhong, 2008; Ryan et al., 2011; Di Bernardini et al., 2011; Najafian and Babji, 2012; Ngo et al., 2012; Rutherford-Markwick, 2012; Udenigwe and Aluko, 2012; Saavedra et al., 2013). Being a major source of high quality proteins, meat presents one of the most investigated sources for isolation of bioactive peptides in recent number of years (Ryan et al., 2011). In addition, it's not only myosin and actin which are used for peptides generation, but other proteins from thick and thin filaments, and connective tissue proteins like fibrillar collagen, as well (Udenigwe and Ashton, 2013).

Bioactive peptides can be generated from protein precursors by different methods including digestive proteolysis in the gastrointestinal tract, chemical or enzymatic hydrolysis *in vitro* during food processing, and microbial fermentation (Korhonen and Pihlanto, 2006; Möller et al., 2008; Shahidi and Zhong, 2008; Ryan et al., 2011; Agyei and Danquah, 2012; Rutherford-Markwick, 2012). In recent years a new method based on molecular genetic engineering, has been reported and it is also possible to synthesize the peptide by chemical or enzymatic synthesis if amino acid sequence is known (Korhonen and Pihlanto, 2006; Shahidi and Zhong, 2008; Hernández-Ledesma et al., 2011; Agyei and Danquah, 2012). Each of these

methods has some advantages or disadvantages, which should be considered when selecting one or several combined methods for a certain purpose (Shahidi and Zhong, 2008). Use of acid hydrolysis in order to release some peptides is economic, relatively simple to perform, but at the same time difficult to control and can damage certain amino acids. Moreover, selectivity and specificity of this chemical hydrolysis is low, which is why this method is rarely used (Shahidi and Zhong, 2008; Rutherford-Markwick, 2012). Methods based on enzymatic hydrolysis have an advantage because they are more predictable with respect to the end products, and the process conditions can be controlled. Because of that, these are commonly used methods for peptide production in laboratories and industry. Enzymes which are used in this technique can be obtained from plants, microorganisms or animals, and can be used alone or in combination with other enzymes, in order to simulate the fate of a protein in *in vitro* condition (Shahidi and Zhong, 2008; Agyei and Danquah, 2012). There are a number of proteinases including trypsin, subtilisin, chymotrypsin, thermolysin, pepsin, proteinase K, papain, alcalase, pronase, papain, carboxypeptidase A, pancreatin and commercial products such as Alcalase Flavourzyme and Neutrase which are used for peptide preparation (Korhonen and Pihlanto, 2006; Shahidi and Zhong, 2008; Agyei and Danquah, 2012; Udenigwe and Ashton, 2013). It is important that isolation of peptides by enzymatic hydrolysis is performed under respective optimal conditions of enzyme (temperature, pH, time course, etc.) for better results (Shahidi and Zhong, 2008; Agyei and Danquah, 2012).

Enzymatic hydrolysis of protein is the technique mostly used for isolation of peptides from meat sources, and the digestive enzymes which are most commonly used are pepsin, trypsin and chymotrypsin (Ryan et al., 2011). Although bacterial fermentation presents valuable method for isolation of bioactive peptides from milk proteins, it wasn't successful in obtaining peptides from meat source, probably because of poor proteolytic activity of the *Lactobacillus spp.* used in meat fermentations (Ryan et al., 2011).

Hydrolysate obtained after the application of one of the previously mentioned methods, presents a mixture mainly composed of peptides and amino acids. Several methods can be used for separation of bioactive peptides from hydrolysate (Ryan et al., 2011; Agyei and Danquah, 2012; Najafian and Babji, 2012). Ultrafiltration membrane system is a method which can be used in order to fractionate hydrolysates based on peptide size and obtained

peptides with desired molecular weights (Ryan et al., 2011; Najafian and Babji, 2012). More precise method is nanofiltration (Najafian and Babji, 2012). For the same purpose, ion exchange, gel filtration technologies, liquid chromatography (HPLC), reversed-phase liquid chromatography (RP-HPLC), and gel permeation chromatography could be used (Pedroche et al., 2007; Chabeaud et al., 2009; Agyei and Danquah, 2012). For strongly charged biomolecules electro-membrane filtration (EMF) can be beneficial technique (Agyei and Danquah, 2012). Matrix-assisted laser desorption/ionization time of flight (MALDI-TOF) mass spectrometric analysis is also utile method (Najafian and Babji, 2012). These methods can be used separately, but combination of two or more methods for production and isolation of peptides may be required (Agyei and Danquah, 2012). It has been reported that HPLC is commonly used with a UV detector or mass spectrometer (Najafian and Babji, 2012). Individual peptide fractions can be identified by using the combined techniques of mass spectrometry and protein sequencing, while liquid chromatography followed by tandem mass spectrometry detection (LC-MS/MS) or conventional membrane filtration with electrophoresis also can be applied (Ryan et al., 2011; Agyei and Danquah, 2012; Najafian and Babji, 2012).

Although, inactive within the sequence of the protein, after the releasing described above, bioactive peptides may induce many beneficial effects on human body (Möller et al., 2008; Korhonen and Pihlanto, 2006; Di Bernardini et al., 2011; Agyei and Danquah, 2012). Their properties were investigated under *in vitro* and *in vivo* conditions, and it has been reported that food derived bioactive peptides have antioxidative, antimicrobial, antihypertensive, antithrombotic, cytomodulatory, anticancer, immunomodulatory, opioid agonistic, mineral binding, hypocholesterolemic and anti-obesity effects. In addition, many of bioactive peptides possess multifunctional properties (Korhonen and Pihlanto, 2006; Möller et al., 2008; Shahidi and Zhong, 2008; Di Bernardini et al., 2011; Ryan et al., 2011; Agyei and Danquah, 2012; Udenigwe and Ashton, 2013). The activity of bioactive peptides depends on amino acid composition and sequence (Shahidi and Zhong, 2008). Moreover, compared to conventional small molecules, these peptides have high bioactivity, act on specific targets inside the body, have low levels of toxicity and they don't accumulate in small amounts in the tissues, which is why many researchers investigate their functional properties and potential applications (Marx, 2005; Agyei and Danquah, 2012).

## Antihypertensive properties

Hypertension is an increasing health problem which affects one third of the worldwide adult population, both men and women, especially in developed countries, and presents the most common type of cardiovascular disease (Ahmed and Muguruma, 2010; Hong et al., 2008; Shahidi and Zhong, 2008; Hernández-Ledesma et al., 2011; He et al., 2013). High blood pressure is predominant factor which contributes to cardiovascular diseases including myocardial infarction, heart failure, coronary heart disease, peripheral artery disease, stroke kidney failure, blindness, end-stage diabetes and even dementia (Hong et al., 2008; Ahmed and Muguruma, 2010; Hernández-Ledesma et al., 2011; Ryan et al., 2011; Sharp et al., 2011; Rutherford-Markwick, 2012; He et al., 2013). It is one of the main causes of the premature death, and WHO estimates that by 2020, heart disease and stroke will become the leading causes of death and disability worldwide (Erdmann et al., 2008; Onuh et al., 2013). There are a number of antihypertensive medications like captopril and analapril on the market, but apart from their advantages, their use may cause some side effects including coughing, taste disturbances, skin rashes, angio-oedema and many other disfunctions of human organs (Wu et al., 2008; Ahmed and Muguruma, 2010; Ryan et al., 2011). Also, these drugs are expensive, and only in the USA, cost of antihypertensive drug annually is approximately \$15 billion (Hong et al., 2008). As a results, over the last two decades, numerous researchers have investigated some effective natural alternatives which would be less expensive for production and cause no side-effects. One of such possibility is the use of food derived bioactive peptides which exhibit antihypertensive effect (Shahidi and Zhong, 2008; Wu et al., 2008; Ahmed and Muguruma, 2010; Hernández-Ledesma et al., 2011; Ryan et al., 2011).

Bioactive peptides act differently then hypotensive drugs. Synthetic substances directly block action of ACE (angiotensin-converting enzyme) responsible for converting angiotensin-I into angiotensin-II, major product of the renin-angiotensin system which presents a powerful vasoconstrictor. ACE hydrolyze bradykinin, a potent vasodilator, also induces the release of aldosterone in the adrenal cortex, which results in increasing of sodium concentration and blood pressure (Wu et al., 2008; Ahmed and Muguruma, 2010; Hernández-Ledesma et al., 2011; Cam and de Mejia, 2012; Escudero et al., 2012; He et al., 2013; Udenigwe and Ashton, 2013). Mechanism of action of ACE inhibitory peptides is based on competing with ACE and preventing the

production of angiotensin-II, causing relaxation of the arterial walls and reduction of fluid volume, in which way these bioactive peptides improve heart function and increase blood and oxygen flow to the heart, liver, and kidneys (Ahmed and Muguruma, 2010; Ryan et al., 2011; He et al., 2013). ACE inhibitory peptides may bind to the active site of the ACE enzyme, or to an inhibitor site located on the ACE enzyme, and in this way change the protein conformation and prevent the angiotensin-I from binding to the enzyme active site (Wijesekara and Kim, 2010; Ryan et al., 2011). Furthermore, some studies show that food-derived bioactive peptides can also inhibit the activity of renin, and in that way induce a reduction of blood pressure (Udenigwe and Aluko, 2012). There are three groups of ACE inhibitory peptides: true inhibitor type, substrate type and pro-drug type, and their classification is based on their inhibitory activity following preincubation with ACE (Iroyukifujita et al., 2000; Arihara and Ohata, 2006; Ryan et al., 2011).

Structure characteristics of peptides are associated with their antihypertensive properties. ACE-inhibitory peptides are mostly peptides with short amino acid sequences containing from 2 to 12 amino acids. Saiga et al., (2003) found that presence of hydroxyproline is crucial for binding of peptides and ACE in cases when peptides contain more than three amino acids in length. Many studies showed that C-terminal of ACE-inhibitory peptides usually contains hydrophobic amino acids residues and that these residues have a crucial role in competitive binding to the active site of ACE (Hernández-Ledesma et al., 2011; Ryan et al., 2011). It has been reported that ACE-inhibitory peptides with the highest antihypertensive activity contain aliphatic, basic and aromatic residues, at the penultimate positions, and aromatic, proline and aliphatic residues at the end of C-terminal. This is explained by interaction of these residues with the three hydrophobic sub-sites located on the active site of ACE (Matsufuji et al., 1994; Iroyukifujita et al., 2000; Ono et al., 2003; Hayes et al., 2007; Qian et al., 2007; Hernández-Ledesma et al., 2011; Ryan et al., 2011). For the same reason, hydrophilic peptides are incompatible with the active sites of ACE, and exhibit none or a weak ACE inhibitory activity (Li et al., 2004; Matsui and Matsumoto, 2006; Ryan et al., 2011). Studies found that N-terminal end of the peptides with ACE-inhibitory activity is hydrophobic (Hayes et al., 2007; Rho et al., 2009; Ryan et al., 2011). Moreover, it has been found that amino acid at the position three from the C-terminal requires the L-configuration (Hernández-Ledesma et al., 2011).

Numbers of peptides with ACE-inhibitory activity were isolated from porcine, beef and chicken muscles. Commonly used methods for the evaluation of ACE-inhibitory effects of bioactive peptides in *in vitro* conditions are those based on spectrophotometric and high-performance liquid chromatography (HPLC) assays (Vermeirssen et al., 2002; Li et al., 2005; Shalaby et al., 2006; Siemerink et al., 2010; Hernández-Ledesma et al., 2011). Studies conducted in *in vivo* systems, are generally based on oral or intravenous application of purified peptides to spontaneously hypertensive rats and measuring of blood pressure immediately after application or after a certain time (Ahmed and Muguruma, 2010; Hernández-Ledesma et al., 2011). Also, in some experiments for investigation of antihypertensive peptides properties normotensive Wistar-Kyoto rats were used (Hernández-Ledesma et al., 2011). Arihara et al. (2001) generated two ACE-inhibitory peptides with amino acid sequence MNPPK, and ITTNP known as from porcine myosin. These peptides were orally applied to spontaneously hypertensive rats (SHR) in order to investigate their effect on systolic blood pressure (SBP). 24 h after oral administration, the SBP of both test groups was still significantly lower than that of the control group, which proved that peptides exhibit antihypertensive effect *in vivo* (Nakashima et al., 2002; Ryan et al., 2011). MNPPK known as myopentapeptide A, is a precursor to tripeptide MNP which exhibited greater antihypertensive activity (Udenigwe and Ashton, 2013). Moreover, M6 peptide with amino acid sequence KRVITY, derived from porcine myosin B by pepsin treatment, showed antihypertensive effect after oral administration to SHR. Maximum decrease of 23 mmHg was noted 6 h after application, and this peptide retained his ACE-inhibitory activity even after thermal process (98 °C for 10 min), (Muguruma et al., 2009; Ryan et al., 2011). In other study, octapeptide with amino acid sequence VKKVLGNP also exhibited ACE-inhibitory effect and caused decrease of SBP in *in vivo* conditions after oral application to SHR (Katayama et al., 2007). Katayama et al. (2007) derived two bioactive peptides from porcine troponin. From troponin C they isolated nine amino acids peptide with sequence RMLGQTPTK, and from crude porcine troponin, peptide with the amino acid sequence KRQKYDI. These antihypertensive peptides are categorized as a non-competitive inhibitor and substrate type inhibitor, respectively (Katayama et al., 2003; Katayama et al., 2004; Katayama et al., 2008; Ryan et al., 2011).

Apart from peptides isolated from myosin and troponin, other meat proteins also present valuable sources for generating peptides with

antihypertensive activity. One anti-hypertensive peptide with amino acid sequence RPR was isolated from pork nebulin, while two antihypertensive peptides with amino acid sequences KAPVA and PTPVP were isolated from pork titin protein by Escudero et al. (2010). Although some of these peptides did not seem to have high ACE-inhibitory activity *in vitro*, they exhibited antihypertensive activity *in vivo*, which was explained by the bioconversion of ACE-inhibitory peptides or by antihypertensive mechanism which could be influenced by these peptides (Lopez-Fandino et al., 2006; Escudero et al., 2012).

In addition, Castellano et al. (2013) used a method of lactic acid bacterial (*L. sakei* CRL1862 and *L. curvatus* CRL705) fermentation in order to generate ACE-inhibitory peptides from porcine proteins. In a study conducted by Terashima et al. (2010), decapeptide with amino acid sequence VTVNPYKWLP was isolated from the myosin heavy chain of chicken leg meat, and its antihypertensive properties were determined (Udenigwe and Ashton, 2013). From three peptides isolated from the chicken breast muscle protein hydrolysates by Saiga et al. (2003) peptide P4, with amino acid sequence GFXGTXLXGF exhibited the strongest ACE inhibiting activity with IC value of 42.4  $\mu\text{M}$ , which was higher compared to IC value of 26  $\mu\text{M}$  what was later reported for the peptide by the same authors (Saiga et al., 2006), P4 was categorized as a non-competitive inhibitor of ACE. In the study P4 were administrated intravenous to SHR in doses of 30 mg per kilogram of body weight, and although it caused an immediate decrease in blood pressure it returned to base pressure 60 minutes post administration, which showed that this peptide does not act as a long-term vasodepressor *in vivo* (Saiga et al., 2006; Ryan et al., 2011; Udenigwe and Ashton, 2013). Jang and Lee (2005) generated a peptide with amino acid sequence VLAQYK, from beef sarcoplasmic proteins with ACE-inhibiting IC 23.1  $\mu\text{g/mL}$ . Furthermore, from dry-cured ham it was derived seven dipeptides (RP, KA, AA, GP, AR, GR and RR) which exhibited ACE inhibitory activity with IC<sub>50</sub> values of 15.2, 31.5, 51.4, 66.0, 95.5, 162.2 and 267  $\mu\text{M}$ , respectively (Udenigwe and Ashton, 2013).

Connective tissue also can be a source for obtaining bioactive peptides with hypotensive effects. For example, Kim et al. (2001) isolated two ACE-inhibitory peptides, EIICIII (GPV) and EIICIV (GPL), from the hydrolysate of bovine skin gelatin which was treated with five proteases (Alcalase, chymotrypsin, Neutrase, Pronase E, and trypsin) in specific order. EIICIII peptide had an IC<sub>50</sub> value of 4.7  $\mu\text{M}$ , while the peptide EIICIV had an IC<sub>50</sub> value

of 2.55  $\mu\text{M}$  (Kim et al., 2001; Ryan et al., 2011). ACE-inhibitory peptides were derived from hydrolysis of chicken collagen as well. At first, collagen was treated by an *Aspergillus oryzae* protease, and then hydrolyzed by treatment with four proteases more (protease FP, protease A, amino G and protease N), after which four oligopeptides were isolated with ACE-inhibitory IC<sub>50</sub> values of 29,4–60,8  $\mu\text{M}$ . They were administrated to SHR at dose of 3 g/kg body weight and SPB were measured. The greatest reduction occurred 6 h after administration (maximum value of -50 mm Hg), but peptide product showed long-term hypotensive effects *in vivo* (4-week) (Saiga et al., 2006; Ryan et al., 2011; Udenigwe and Ashton, 2013). Protein hydrolysates derived by Onuh et al. (2013) from chicken skin through alcalase or simulated gastrointestinal digestion showed to possess inhibitory activities against ACE and renin in *in vitro* tests.

Bioactive peptides with ACE-inhibitory activity were isolated and identified from many fish species such as shellfish, tuna, bonito, salmon and sardine (Yokoyama et al., 1992; Matsufuji et al., 1994; Ono et al., 2003; Qian et al., 2007; Hong et al., 2008; Ahmed and Muguruma, 2010; Ryan, 2011). Moreover, Wu et al. (2008) isolated and identified four peptides (CF, EY, MF and FE) with high ACE-inhibitory activity from shark meat and two of them (EY and FE) have never been reported before.

## Antithrombotic properties

Arterial thrombosis often presents a cause or complicate some vascular diseases and conditions like myocardial infarction and stroke. Some peptides obtained from meat sources showed antithrombotic properties and it is considered that their use in the future can be beneficial in the prevention or control of such conditions. (Udenigwe and Ashton, 2013). Shimizu et al. (2009) isolated a peptide with molecular weight of 2.5 kDa from defatted porcine *musculus longissimus dorsi* and investigated his effect on thrombosis. Pork meat was treated by papain protease and hydrolyzed peptide was implicated orally to mice in doses 210 mg/kg of body weight. After that a carotid artery thrombosis was induced with helium-neon laser and total thrombus size were calculated. Results of this study showed that peptide can significantly inhibit thrombus formation by decreasing platelet activity and have same effect as aspirin administration at 50 mg/kg body weight (Shimizu et al. 2009; Cam and de Mejia, 2012; Udenigwe and Ashton, 2013).

## Antioxidant properties

Reactive oxygen species (ROS) and free radicals attack and interact with membrane lipids, protein and DNA in the cell. They can be endogenous or exogenous origin, but in both case have influence on human health and play a great role in ethiology and progression of several diseases including cardiovascular diseases, atherosclerosis, arthritis, diabetes, inflammation, cancer, neuropathies, Alzheimer's and other degenerative diseases as well (Ames, 1983; Esterbauer, 1993; Cai *et al.*, 2002; Gimenez *et al.*, 2009; Ryan *et al.*, 2011; Jomova *et al.*, 2012; Udenigwe and Ashton, 2013). Oxidation by free radicals is also one of the primary mechanisms of quality deterioration in foods and especially in meat products, which also limits shelf-life and makes meat potentially dangerous for consumer's health (Simitzis *et al.*, 2009; Bošković *et al.*, 2013; Udenigwe and Ashton, 2013). In order to prevent or to retard lipid oxidation, a number of synthetic antioxidants, such as butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), tert-butyl hydroquinone (TBHQ) and propyl gallate (PG) are added to food (Saiga *et al.*, 2003a; Di Bernardini *et al.*, 2011; Ngo *et al.*, 2012). Their use may have negative influence on human health which is why food industry tends to find natural alternatives to synthetic antioxidants. (Saiga *et al.*, 2003a; Sakanaka and Tachibana, 2006; Kim *et al.*, 2001; Di Bernardini *et al.*, 2011). One of such possibilities is use of peptides from food sources, which have some advantages, compared to synthetic antioxidants. They are considered to be safe for consumers, economic for production, have high activity, they are easy to absorb, also have nutritional value and do not cause immunoreactions like enzymatic antioxidants. The antioxidant effect of peptides was firstly reported by Marcuse in 1960 (Gómez-Guillén *et al.*, 2011). Since then, in order to confirm their antioxidant properties numbers of studies were conducted on peptides from mostly plant and animal sources such as milk, milk-kefir and soymilk-kefir, casein, egg-yolk protein, soybean protein, wheat, potato, rice bran, sunflower protein, leaf protein, peanut kernels, corn gluten meal, frog skin, medicinal mushroom and fungi (Suetsuna *et al.*, 2000; Peña-Ramos *et al.*, 2004; Sun *et al.*, 2004; Wachtel-Galor *et al.*, 2004; Liu *et al.*, 2005; Zhu *et al.*, 2006; Sakanaka and Tachibana, 2006; Li *et al.*, 2008; Megias *et al.*, 2008; Pihlanto *et al.*, 2008; Qian *et al.*, 2008; Xie *et al.*, 2008; Revilla *et al.*, 2009; Hwang *et al.*, 2010; Gómez-Guillén *et al.*, 2011; Ryan *et al.*, 2011). In recent years, interest of scientists for peptides from

meat, especially fish sources, has increased (Ryan *et al.*, 2011).

In spite of all the research, the exact mechanism of antioxidant activity of peptides still has not been fully understood. Based on current knowledge, it is supposed that peptides are scavengers of free radicals and ROS, they inhibit lipid peroxidation and chelate transition metal ions (Suetsuna *et al.*, 2000; Wu *et al.*, 2003; Rajapakset *et al.*, 2005; Gómez-Guillén *et al.*, 2011; Young *et al.*, 2013). In addition, it has been proved that the antioxidant properties of peptides, and especially peptide composition and structure may be affected by a method used to isolate proteins, degree of hydrolysis, type of used protease, peptide concentration and hydrophobicity (Suetsuna *et al.*, 2000; Saiga *et al.*, 2003b; Peña-Ramos *et al.*, 2004; Erdmann *et al.*, 2008; Liu *et al.*, 2010). Type of amino acid, their number in the peptide, as well as the arrangements of amino acid sequence play an important role in antioxidant activity (Suetsuna *et al.*, 2000; Saito *et al.*, 2003a; Rajapakset *et al.*, 2005; Erdmann *et al.*, 2008). Tyr, Trp, Met, Lys, Cys, and His are those amino acids which contribute to antioxidant activity (Peña-Ramos *et al.*, 2004; Wang *et al.*, 2005; Sarmadi and Ismail, 2010; Di Bernardini *et al.*, 2011). Histidine-containing peptides possess imidazole group which is considered to be in relation with the hydrogen-donating, lipid peroxy radical trapping and the metal chelating, while SH group in cysteine has a main role in interaction with free radicals (Chan *et al.*, 1994; Erdmann *et al.*, 2008; Qian *et al.*, 2008; Sarmadi and Ismail, 2010). Moreover, it has been found that substitution of L-His by D-His in a peptide leads to reduction of the antioxidative activity, which proves that configuration of amino acids also has influence on antioxidant activity (Chen *et al.*, 1996; Sarmadi and Ismail, 2010). Some researchers have found that certain amino acids exhibit higher antioxidative activity when they are incorporated in dipeptides (Alabovskiy *et al.*, 1997; Takenaka *et al.*, 2003; Erdmann *et al.*, 2008; Sarmadi and Ismail, 2010).

The most studied hydrophilic antioxidants from meat are histidine-containing dipeptides, carnosine ( $\beta$ -alanyl-L-histidine) and anserine (N- $\beta$ -alanyl-1-methyl-L-histidine), (Decker *et al.*, 2000; Guiotto *et al.*, 2005; Arihara and Ohata, 2006; Di Bernardini *et al.*, 2011; Young *et al.*, 2013). They are found only in meat, poultry and in some fish (Young *et al.*, 2013). The concentration of carnosine in meat depends on type of meat, and ranges from 500 mg/kg in chicken to 2700 mg/kg in pork, while anserine is present in higher amounts in chicken muscle (Purchas and Busboom, 2005; Purchas *et*

al., 2004; Young et al., 2013). The antioxidant activity of these dipeptides is attributed mainly to their ability to chelate prooxidative metals such as copper, zinc and cobalt, but it has been found that carnosine is able to scavenge free radicals and form conjugates with potentially toxic aldehydic products from lipid oxidation as well (Brown, 1981; Decker et al., 2000; Young et al., 2013). It has been reported the ability of radioprotection of DNA by carnosine and anserine and protection of DNA by carnosine, against L-3, 4-dihydroxyphenylalanine Fe (III) induced damage. Some data showed that oral administration of L-carnosine has the same effect on increase of total antioxidant capacity of human serum as a consumption of beefsteaks (Di Bernardini et al., 2011).

Apart from carnosine and anserine, there are many antioxidative peptides from meat sources, which are generated from proteins by different methods. In one study, Saiga et al. (2003b) treated porcine myofibrillar proteins with two proteases, papain and actinase E, and found that hydrolyzates derived in this way exhibit high levels of antioxidant activity in a linolenic acid peroxidation system. Compared to five isolated peptides from papain hydrolysate (DSGVT-actin, IEAEGE-unknown, DAQEKLE-tropomyosin, EELDNLN-tropomyosin, VPSIDDQEELM-myosin heavy chain) DAQEKLE showed the highest level of activity, which was very similar to the activity of  $\alpha$ -tocopherol at pH 7. Also, it was reported that peptides which were obtained from myofibrillar proteins by actinase E treatment showed higher antioxidant activity, which proves that type of used proteolytic enzymes play an important role in determining the antioxidative properties of peptides (Arihara and Ohata, 2006; Di Bernardini et al., 2011; Ryan et al., 2011; Udenigwe and Ashton, 2013). In other study Arihara et al. (2005) found that peptides ALTA, SLTA, and VT, obtained from papain treated porcine skeletal muscle actomyosin exhibit antioxidative activity not only *in vitro*, but *in vivo* system, as well (Arihara and Ohata, 2006). Numerous studies were carried out on peptides derived from collagen (Gómez-Guillén et al., 2011). Li et al. (2007) treated porcine collagen with pepsin and then derived hydrolysate was treated with papain, protease from bovine pancreas (PP) and a cocktail of three enzymes (PP, bacterial proteases from *Streptomyces* and *Bacillus polymyxa*). The hydrolysate of collagen which was treated with cocktail of three enzymes showed the highest level of antioxidant activity, and four antioxidant peptides were isolated from this hydrolysate (QGAR, LQGM, LQGMH and HC) (Li et al., 2007; Di Bernardini et al., 2011; Ryan et al., 2011). A 36-amino acid residue

peptide GETGPAGPAGPIPVGARGPAGPQGPR GDKGETGEQ, which showed ability of free radical scavenging and metal chelating were isolated from bovine tendon collagen  $\alpha 1$  by Banerjee et al. (2012), (Udenigwe and Ashton, 2013). Result of other studies showed that peptides obtained from papain-hydrolyzed beef sarcoplasmic proteins, and antihypertensive peptides from dry-cured ham also exhibited antioxidant activities (Di Bernardini et al., 2012; Escudero et al., 2012; Udenigwe and Ashton, 2013).

## Anticancer properties

It has been proved that some peptides isolated from meat and marine organisms, especially fish, exhibit anti-cancer activity, inhibit cell proliferation and have cytotoxic effect against tumor cells (Shahidi and Zhong, 2008; Ryan et al., 2011; Najafian and Babji, 2012; Udenigwe and Aluko, 2012).

Peptides with antibacterial activity isolated from beef sarcoplasmic proteins were investigated by Jang et al. (2008) in order to prove their cytotoxic effect against human breast adenocarcinoma (MCF-7), gastric adenocarcinoma (AGS) and lung carcinoma (A549) cell lines. GFHI showed the strongest cytotoxic effect on MCF-7 cells and decreased the cell viability of AGS cells. GLSDGEWQ strongly inhibited the proliferation of AGS cells, while none of tested peptides had a cytotoxic effect on A549 cells (Jang et al., 2008; Ryan et al., 2011; Udenigwe and Ashton, 2013). Hsu et al. (2011) isolated two peptides from tuna dark muscle which was treated with two proteases, papain and protease XXII. Amino acid sequences of these peptides were LPHVLTPEAGAT from papain hydrolysate and PTAEGVYMVT, from protease XXIII and both of them exhibited dose-dependent antiproliferative activities against human breast adenocarcinoma (MCF-7) cells, (Hsu et al., 2011; Ryan et al., 2011; Udenigwe and Aluko, 2012). Picot et al. (2006) reported 18 protein hydrolysates isolated from blue whiting, cod, plaice, and salmon to have antiproliferative activity against 2 human breast cancer cell lines (MCF-7/6 and MDA-MB-231) (Picot et al., 2006; Shahidi and Zhong, 2008; Ryan et al., 2011). In addition, it has been shown that hydrophobic peptide isolated from anchovy sauce, with molecular weight of 440.9 Da, induced apoptosis in a human lymphoma cell line (U937), (Lee et al., 2003; Lee et al., 2004; Ryan et al., 2011).

These peptides, which showed to possess anticancer properties *in vivo*, could be further used to investigate their possibility to prevent the development of different types of cancer or even more, in their treatment.

## Antibacterial properties

Antimicrobial peptides are usually composed of less than 50 amino acids, and about a half of them are hydrophobic (Shahidi and Zhong, 2008; Najafian and Babji, 2012). Their antibacterial activity is different and varies depending on origin of the peptides, amino acid composition, peptide size, charge, hydrophobicity, and secondary structure of peptides (Shahidi and Zhong, 2008). In recent number of years the overuse of antibiotics in human and veterinary medicine in order to reduce pathogens has led to phenomenon of multi-drug-resistance bacteria (Sofos, 2008; Tohidpour et al., 2010; Bošković et al., 2013). One possible unconventional solution to this increasing problem could be the use of antimicrobial peptides in medical proposes (Najafian and Babji, 2012).

Antibacterial properties of peptides can be tested by several methods. The most commonly used method is agar diffusion. This method is based on measuring of the inhibition zone diameter formed on agar, but in order to determinate the exact antibacterial activity, the minimal inhibitory concentration (MIC) of peptides should be determined (Di Bernardini et al., 2011; Najafian and Babji, 2012).

Although a number of peptides with antimicrobial activity have been isolated from milk proteins, there is no much data on the antimicrobial peptides from meat sources in the available literature (Pihlanto, 2002; McCann et al., 2005; Hayes et al., 2006; McCann et al., 2006; Minervini et al., 2003; Di Bernardini et al., 2011; Ryan et al., 2011; Agyei and Danquah, 2012).

In one study, Jang et al., (2008) evaluated the antimicrobial effects of four peptides (GLSDGEWQ, GFHI, DFHING and FHG) isolated from beef sarcoplasmic proteins, which were previously determined to have ACE-I-inhibitory activity. Antimicrobial activity of these peptides was evaluated against three Gram-positive (*Bacillus cereus*; *Staphylococcus aureus* - KFRI00188; and *Listeria monocytogenes* -KFRI00719) and Gram-negative (*Salmonella typhimurium*-KFRI0025; *Escherichia coli*- ATCC43894 and *Pseudomonas aeruginosa*-KFRI00100) pathogenic bacteria. For this purpose the paper disc diffusion method was used, and peptides were applied at three different concentrations. All four tested peptides exhibited antimicrobial activities against one or more bacteria. Results showed that GLSDGEWQ had the highest level of antimicrobial activity, and was the only peptide that inhibited growth of both Gram-negative and Gram-positive bacteria at all three used concentrations. FHG inhibited *P. aeruginosa*, DFHING inhibited

*E. coli* at all tested concentrations and GFHI exhibited antibacterial activity against *E. coli* and *P. aeruginosa*, but neither one of them inhibited the growth of *L. monocytogenes* (Di Bernardini et al., 2011; Ryan et al., 2011; Udenigwe and Ashton, 2013). Numbers of peptides with antimicrobial activity have been isolated from fish sources. Liu et al. (2008) isolated a cysteine rich antimicrobial peptide (CgPep33) from oyster muscle by using a combination of alcalase and bromelin. This peptide showed antimicrobial activity against pathogenic bacteria, such as *E. coli*, *P. aeruginosa*, *B. subtilis* and *S. aureus*, and also some fungi (*Botrytis cinerea* and *Penicillium expansum*), (Ryan et al., 2011). In a study conducted by Gómez-Guillén et al. (2010) it has been found that peptides obtained from tuna and squid skin gelatins showed high level of antimicrobial activity against *Lactobacillus acidophilus* and *Bifidobacterium animalis* subsp. *lactis*, *Shewanella putrefaciens* and *Photobacterium phosphoreum* (Gómez-Guillén et al., 2011). Protein from skin homogenate of *Epinephelus fario* by trypsin digestion, which showed activity against Gram-positive bacteria (*Vibrio alginolyticus*, *Vibrio parahaemolyticus*, *Vibrio fluvialis*, *Pasteurella multocida*, *E. coli*, *Aeromonas hydrophila* and *P. aeruginosa*) (Najafian and Babji, 2012).

## Other properties

In recent years, obesity, hyperlipidemia and especially hypercholesterolemia became serious public health problems, which contribute mainly to cardiovascular diseases, but also to diabetes type 2, hypertension and stroke, certain forms of cancer and sleep-breathing disorder, as well. There is a great number of synthetic drugs with cholesterol-lowering effect, but nowadays researches are looking for natural alternatives which can be used in prevention and treatment of hypercholesterolemia (Shahidi and Zhong, 2008; Ngo et al., 2012). One of such possibilities is the use of food derived peptides. Although mainly soy and milk derived proteins showed lipid-lowering effect, researchers investigate and explore other possibilities among other peptides derived from meat. One study showed that protein hydrolysate isolated from pork with papain exhibit a hypocholesterolemic effect in cholesterol-fed rats (Morimatsu and Kimura, 1992; Morimatsu et al., 1996; Shahidi and Zhong, 2008).

There are some evidences that dipeptide carnosine exhibits significant pharmacological effects and could play a role in preventing or treating some pathological conditions, such as neurodegeneration,



diabetes and cataract (Guiotto et al., 2005; Lee et al., 2005). These effects of carnosine were mainly related to its antioxidant or antiglycating properties (Aldini et al., 2005; Fu et al., 2009). Baran (2000) found that carnosine zinc complex alleviates injuries of gastric mucosa, acts against stomach ulcers and inhibits growth of the main gastric pathogen *Helicobacter pylori*. Some studies found that this antioxidative peptide also plays role in injury healing, recovery from fatigue and prevention of diseases related to stress (Baran, 2000; Matsukura and Tanaka, 2000; Young et al., 2013). Arihara et al. (2005) isolated two peptides (ALTA and SLTA) from pork actomyosin by papain protease treatment. *In vitro* it has been found that these peptides showed antioxidative activity. In a study *in vivo* these peptides showed antifatigue effects after being orally applied to mice which were running on treadmill (Arihara and Ohata, 2006).

In addition, it has been reported by Nakatani et al. (2009) that dipeptide PX isolated from porcine meat contributes to reparation and maintenance of cartilage by preventing mature chondrocytes from

becoming mineralized and stimulating production of other protective peptides, while Iwai et al. (2005) found that peptides derived from collagen exhibit some immune-modulating activities by stimulating proliferation of fibroblasts, neutrophils, and monocytes (Iwai et al., 2005; Nakatani et al., 2009; Udenigwe and Ashton, 2013). Some peptides, such as commercial fish protein hydrolysate exhibits immunomodulatory activities by increasing the number of IgA+ cells and IL-4, IL-6 and IL-10 in the lamina propria of the small intestine in mice (Duarte et al., 2006; Möller et al., 2008).

## Conclusion

Although, there is still a small number of studies, especially *in vivo* studies, which should be conducted in order to confirm safety and beneficial effects of bioactive peptides, scientific, technological and consumer interest for these peptides and their potential use in controlling and promoting health increases, and results remains to be seen.

## References

- Agyei D., Danquah M. K., 2012. Rethinking food-derived bioactive peptides for antimicrobial and immunomodulatory activities. *Trends in Food Science & Technology*, 23, 2, 62–69.
- Ahmed A. M., Muguruma M., 2010. A review of meat protein hydrolysates and hypertension. *Meat Science*, 86, 1, 110–118.
- Alabovsky V. V., Boldyrev A. A., Vinokurov A. A., Shchavratsky V., 1997. Effect of histidine-containing dipeptides on isolated heart under ischemia/reperfusion. *Biochemistry (Moscow)*, 62, 77–87.
- Aldini G., Facino R. M., Beretta G., Carini M., 2005. Carnosine and related dipeptides as quenchers of reactive carbonyl species: From structural studies to therapeutic perspectives. *Biofactors* 24, 77–87.
- Ames B. N., 1983. Dietary carcinogens and anticarcinogens. Oxygen radicals and degenerative diseases. *Science*, 221, 1256–1264.
- Anand P., Kunnumakara B. A., Sundaram C., Harikumar H. K., Tharakan T. S., Lai S. O., Sung B., Aggarwal B. B., 2008. Cancer is a preventable disease that requires major lifestyle changes. *Pharmaceutical Research*, 25, 2097–2116.
- Arihara K., Nakashima Y., Mukai T. Ishikawa S., Itoh M., 2001. Peptide inhibitors for angiotensin I-converting enzyme from enzymatic hydrolysates of porcine skeletal muscle proteins. *Meat Science*, 57, 319–324.
- Arihara K., Tomita K., Ishikawa S., Itoh M., Akimoto M., Sameshima T., 2005. Anti-fatigue peptides derived from meat proteins. Japan patent, submitted to government.
- Arihara K., 2006. Strategies for designing novel functional meat products. *Meat Science* 74, 219–229.
- Arihara K., Ohata M., 2006. Functional Properties of Bioactive Peptides Derived from meat Proteins. In *Advanced Technologies for Meat Processing*; Toldra, F., Ed.; Springer: New York, NY, USA, 245–274.
- Babizhayev M. A., 2005. Analysis of lipid peroxidation and electron microscopic survey of maturation stages during human cataractogenesis: pharmacokinetic assay of CanC(TM) N-acetylcarnosine prodrug lubricant eye drops for cataract prevention. *Drugs R & D* 6, 345–369.
- Baltić Ž. M., Dragičević O., Karabasil N., 2002. Trends in meat consumption, Zbornik radova i kratkih sadržaja. 14. Savetovanje veterinara Srbije, Zlatibor, 10–14. septembar 2002.
- Baltić Ž. M., Bošković M., Mitrović R., 2013. *In Vitro Meat: Possibility of the Impossible*, International 57<sup>th</sup> Meat Industry Conference, Belgrade, 10<sup>th</sup>–12<sup>th</sup> June, 2013.
- Banerjee P., Suseela G., Shanthi C., 2012. Isolation and identification of cryptic bioactive regions in bovine Achilles tendon collagen. *The Protein Journal*, 31, 374–386.
- Baran E. J., 2000. Metal complexes of carnosine. *Biochemistry (Moscow)*, 65, 789–797.
- Biesalski, H. K., 2005. Meat as a component of a healthy diet – are there any risks or benefits if meat is avoided in the diet? *Meat Science*, 70, 509–524.
- Bošković M., Baltić Ž. M., Ivanović J., Đurić J., Lončina J., Dokmanović M., Marković R., 2013. Use of essential oils in order to prevent food borne illness caused by pathogens in meat, *Meat Technology*, 54, 1, 14–21.

- Brown C. E., 1981.** Interactions among carnosine, anserine, ophidine and copper in biochemical adaptation. *Journal of Theoretical Biology*, 88, 245–256.
- Cai W., Gao Q. D., Zhu L., Peppas M., He C., Vlassara H. 2002.** Oxidative stress-inducing carbonyl compounds from common foods: Novel mediators of cellular dysfunction. *Molecular Medicine*, 8, 337–346.
- Cam A., de Mejia E. G., 2012.** Role of dietary proteins and peptides in cardiovascular disease. *Molecular Nutrition & Food Research*, 56, 1, 53–66.
- Castellano P., Aristoy M. C., Sentandreu, M. Á., Vignolo, G., Toldrá, F., 2013.** Peptides with angiotensin I converting enzyme (ACE) inhibitory activity generated from porcine skeletal muscle proteins by the action of meat-borne *Lactobacillus*. *Journal of Proteomics*, 89, 183–190.
- CDC National Center for Chronic Disease Prevention., 2005.** Number of Americans with Diabetes Continues to Increase. In O. o. Communication (Ed.), (Vol. CDC).
- Chabeaud A., Vandanjon L., Bourseau P., Jaouen P., Chaplain-Derouiniot M., Guerard F., 2009.** Performances of ultrafiltration membranes for fractionating a fish protein hydrolysate: application to the refining of bioactive peptidic fractions. *Separation and Purification Technology*, 66, 3, 463–471.
- Chan K. M., Decker E. A., Feustman C., 1994.** Endogenous skeletal muscle antioxidants. *Critical Reviews in Food Science & Nutrition*, 34, 4, 403–426.
- Chan W., 2004.** Macronutrients in meat. In W. K. Jensen, C. Devine, & M. Dikeman (Eds.), *Encyclopedia of meat sciences*, 614–618. Oxford: Elsevier
- Chen H. M., Muramoto K., Yamauchi F., Nokihara K. 1996.** Antioxidant activity of design peptides based on the antioxidative peptide isolated from digests of a soybean protein. *Journal of Agricultural Food Chemistry*, 44, 2619–23.
- Decker A. E., Livisay S. A., Zhou S., 2000.** A Re-evaluation of the Antioxidant Activity of Purified Carnosine. *Biochemistry, (Moscow)*, 65, 7, 901–906.
- Decker E. A., Livisay S. A., Zhou S., 2000.** Mechanisms of endogenous skeletal muscle antioxidants: Chemical and physical aspects. In E. A. Decker, C. Faustman, & C. L. Lopez-Bote (Eds.), *Antioxidants in muscle foods*, 25–60. New York: WileyInterscience.
- Decker E. A., Park Y., 2010.** Healthier meat products as functional foods. *Meat Science*, 86, 1, 49–55.
- Descalzo A. M., Sancho A. M., 2008.** A review of natural antioxidants and their effects on oxidative status, odor and quality of fresh beef produced in Argentina. *Meat Science*, 79, 423–436.
- DHHS, 2010.** Chronic disease cost. [http://health.nv.gov/CD\\_ChronicDisease\\_Costs.htm](http://health.nv.gov/CD_ChronicDisease_Costs.htm).
- Di Bernardini R., Harnedy P., Bolton D., Kerry J., O'Neill E., Mullen A. M., Hayes M., 2011.** Antioxidant and antimicrobial peptidic hydrolysates from muscle protein sources and by-products. *Food Chemistry*, 124, 4, 1296–1307.
- Di Bernardini R., Mullen A. M., Bolton D., Jerry J., O'Neill E., Hayes M., 2012.** Assessment of the angiotensin-I-converting enzyme (ACE-I) inhibitory and antioxidant activities of hydrolysates of bovine brisket sarcoplasmic proteins produced by papain and characterization of associated bioactive peptidic fractions. *Meat Science*, 90, 226–235.
- Diplock A. T., Aggett P. J., Ashwell M., Bornet F., Fern E. B., Roberfroid M. B., 1999.** Scientific concept of functional foods in Europe Consensus document. *The British Journal of Nutrition*, 81, 1–27.
- Duarte J., Vinderola G., Ritz B., Perdigon G., Matar C., 2006.** Immunomodulating capacity of commercial fish protein hydrolysate for diet supplementation. *Immunobiology* 211, 341–350.
- Erdmann K., Cheung B. W., Schröder H., 2008.** The possible roles of food-derived bioactive peptides in reducing the risk of cardiovascular disease. *The Journal of Nutritional Biochemistry*, 19, 10, 643–654.
- Escudero E., Sentandreu M. A., Arihara K., Toldrá F., 2010.** Angiotensin I-converting enzyme inhibitory peptides generated from in vitro gastrointestinal digestion of pork meat. *Journal of Agricultural and Food Chemistry*, 58, 5, 2895–2901.
- Escudero E., Aristoy M. C., Nishimura H., Arihara K., Toldrá F., 2012.** Antihypertensive effect and antioxidant activity of peptide fractions isolated from Spanish dry-cured ham. *Meat Science*, 91, 306–311.
- Esterbauer H., 1993.** Cytotoxicity and genotoxicity of lipid-oxidation products. *The American Journal of Clinical Nutrition*, 57, 5, 779–785.
- Fu H., Katsumura Y., Lin M., Muroya Y., Hata K., Fujii K., Hatano Y., 2009.** Free radical scavenging and radioprotective effects of carnosine and anserine. *Radiation Physics and Chemistry*, 78, 12, 1192–1197.
- Giménez B., Alemán A., Montero P., Gómez-Guillén M. C., 2009.** Antioxidant and functional properties of gelatin hydrolysates obtained from skin of sole and squid. *Food Chemistry*, 114, 3, 976–983.
- Gómez-Guillén M. C., López-Caballero M. E., López de Lacey A., Alemán A., Giménez B., Montero P., 2010.** Antioxidant and antimicrobial peptide fractions from squid and tuna skin gelatin. In E. Le Bihan, N. Koueta (Eds.), *Sea by-products as a real material: New ways of application*, 89–115. Kerala, India: Transworld Research Network Signpost, Chapter 7.
- Gómez-Guillén C. M., Giménez B., López-Caballero M. E., Montero M. P., 2011.** Functional and bioactive properties of collagen and gelatin from alternative sources: A review. *Food Hydrocolloids* 25, 1813–1827.
- Guiotto A., Calderan A., Ruzza P., Borin G., 2005.** Carnosine and carnosine-related antioxidants: A review. *Current Medicinal Chemistry*, 12, 2293–2315.
- Hartmann R., Meisel H., 2007.** Food-derived peptides with biological activity: from research to food applications. *Current Opinion in Biotechnology*, 18, 2, 163–169.
- Hayes M., Ross R. P., Fitzgerald G. F., Hill C., Stanton C., 2006.** Casein-derived antimicrobial peptides generated by *Lactobacillus acidophilus* DPC6026. *Applied and Environmental Microbiology*, 72, 3, 2260–2264.
- Hayes M., Stanton C., Fitzgerald G. F., Ross R. P., 2007.** Putting microbes to work: dairy fermentation, cell factories and bioactive peptides. Part II: bioactive peptide functions. *Biotechnology journal*, 2, 4, 435–449.
- Hayes M., Stanton C., Slattery H., O'Sullivan O., Hill, C., Fitzgerald G. F., Ross R. P., 2007.** Casein fermentate of *Lactobacillus animalis* DPC6134 contains a range of novel propeptide angiotensin-converting enzyme inhibitors. *Applied and environmental microbiology*, 73, 14, 4658–4667.

- He R., Alashi A., Malomo S. A., Girgih A. T., Chao D., Ju X., Aluko R. E., 2013. Antihypertensive and free radical scavenging properties of enzymatic rapeseed protein hydrolysates. *Food Chemistry*, 141, 1, 153–159.
- Hernández-Ledesma B., del Mar Contreras M., Recio I., 2011. Antihypertensive peptides: production, bioavailability and incorporation into foods. *Advances in Colloid and Interface Science*, 165, 1, 23–35.
- Higgs J. D., 2000. The changing nature of red meat: 20 years of improving nutritional quality. *Trends in Food Science & Technology*, 11, 3, 85–95.
- Hong F., Ming L., Yi S., Zhanxia L., Yongquan W., Chi L., 2008. The antihypertensive effect of peptides: a novel alternative to drugs? *Peptides*, 29, 6, 1062–1071.
- Hsu K. C., Li-Chan E. C., Jao C. L., 2011. Antiproliferative activity of peptides prepared from enzymatic hydrolysates of tuna dark muscle on human breast cancer cell line MCF-7. *Food Chemistry*, 126, 2, 617–622.
- Hwang J. Y., Shyu Y. S., Wang Y. T., Hsu C. K., 2010. Antioxidative properties of protein hydrolysate from defatted peanut kernels treated with esperase. *LWT-Food Science and Technology*, 43, 2, 285–290.
- Iroyukifujita H., Eiichiyokoyama K., Yoshikawa M., 2000. Classification and Antihypertensive Activity of Angiotensin I-Converting Enzyme Inhibitory Peptides Derived from Food Proteins. *Journal of Food Science*, 65, 4, 564–569.
- Jang A., Lee, M., 2005. Purification and identification of angiotensin converting enzyme inhibitory peptides from beef hydrolysates. *Meat Science*, 69, 653–661.
- Jang A., Jo C., Kang K. S., Lee M., 2008. Antimicrobial and human cancer cell cytotoxic effect of synthetic angiotensin-converting enzyme (ACE) inhibitory peptides. *Food Chemistry*, 107, 327–336.
- Jiménez-Colmenero, F., Carballo, J., Cofrades, S., 2001. Healthier meat and meat products: their role as functional foods. *Meat Science*, 59, 1, 5–13.
- Jiménez-Colmenero F., Sanchez-Muniz F., Olmedilla-Alonso B., 2010. Design and development of meat-based functional foods with walnut: Technological, nutritional and health impact. *Food Chemistry*, 123, 959–967.
- Jiménez Colmenero F., Herrero A., Cofrades S., Ruiz-Capillas C., 2012. Meat and functional foods. In Y. H. Hui (Ed.), *Handbook of meat and meat processing* (225–248) (2<sup>nd</sup> ed.). Boca Raton: CRC Press. Taylor & Francis Group.
- Jomova K., Baros S., Valko M., 2012. Redox active metal-induced oxidative stress in biological systems. *Transition Metal Chemistry*, 37, 127–134.
- Katayama, K., Tomatsu, M., Fuchu, H., Sugiyama, M., Kawahara, S., Yamauchi, K., Kawamura, Y., Muguruma, M., 2003. Purification and characterization of an angiotensin I converting enzyme inhibitory peptide derived from porcine troponin C. *Animal Science Journal*, 74, 53–58.
- Katayama K., Makoto T., Satoshi K., Kiyoshi Y., Hidetaka F., Yoshiro K., Yukio K., Michio M., 2004. Inhibitory profile of nonapeptide derived from porcine troponin C against angiotensin I-converting enzyme. *Journal of agricultural and food chemistry* 52, 4, 771–775.
- Katayama K., Mori T., Kawahara S., Miake K., Kodama Y., Sugiyama M., Kawamura Y., Nakayama T., Maruyama M., Muguruma M., 2007. Angiotensin-I Converting Enzyme Inhibitory Peptide Derived from Porcine Skeletal Muscle Myosin and Its Antihypertensive Activity in Spontaneously Hypertensive Rats. *Journal of Food Science*, 72, 9, 702–706.
- Katayama, K., Anggraeni, H. E., Mori, T., Ahhmed, A. M., Kawahara, S., Sugiyama, M., Muguruma, M., 2008. Porcine skeletal muscle troponin is a good source of peptides with angiotensin-I converting enzyme inhibitory activity and antihypertensive effects in spontaneously hypertensive rats. *Journal of Agricultural and Food Chemistry*, 56, 2, 355–360.
- Kim S. K., Byun H. G., Park P. J., Shahidi F., 2001. Angiotensin I converting enzyme inhibitory peptides purified from bovine skin gelatin hydrolysate. *Journal of Agricultural and Food Chemistry*, 49, 6, 2992–2997.
- Korhonen H., Pihlanto A., 2006. Bioactive peptides: production and functionality. *International Dairy Journal*, 16, 9, 945–960.
- Lee Y. T., Hsu C. C., Lin M. H., Liu K. S., Yin M. C., 2005. Histidine and carnosine delay diabetic deterioration in mice and protect human low density lipoprotein against oxidation and glycation. *European Journal of Pharmacology*, 513, 145–150.
- Lee Y. G., Lee K. W., Kim J. Y., Kim K. H., Lee H. J., 2004. Induction of apoptosis in a human lymphoma cell line by hydrophobic peptide fraction separated from anchovy sauce. *Biofactors*, 21, 1, 63–67.
- Lee Y., Kim J. I. Y., Lee K., Kim K. H., Lee H., 2003. Peptides from Anchovy Sauce Induce Apoptosis in a Human Lymphoma Cell (U937) through the Increase of Caspase-3 and -8 Activities. *Annals of the New York Academy of Sciences*, 1010, 1, 399–404.
- Li B., Chen F., Wang X., Ji B., Wu Y., 2007. Isolation and identification of antioxidative peptides from porcine collagen hydrolysate by consecutive chromatography and electrospray ionization–mass spectrometry. *Food Chemistry*, 102, 4, 1135–1143.
- Li G. H., Le G. W., Shi Y. H., Shrestha S., 2004. Angiotensin I-converting enzyme inhibitory peptides derived from food proteins and their physiological and pharmacological effects. *Nutrition Research*, 24, 469–486.
- Li G. H., Liu H., Shi Y. H., Le G. W., 2005. Direct spectrophotometric measurement of angiotensin I-converting enzyme inhibitory activity for screening bioactive peptides. *Journal of Pharmaceutical and Biomedical Analysis*, 37, 2, 219–224.
- Li X. X., Han L. J., Chen L. J., 2008. In vitro antioxidant activity of protein hydrolysates prepared from corn gluten meal. *Journal of the Science of Food and Agriculture*, 88, 9, 1660–1666.
- Liu J. R., Chen M. J., Lin C. W., 2005. Antimutagenic and antioxidant properties of milk-kefir and soymilk-kefir. *Journal of Agricultural and Food Chemistry*, 53, 7, 2467–2474.
- Liu Q., Kong B., Xiong Y. L., Xia X., 2010. Antioxidant activity and functional properties of porcine plasma protein hydrolysate as influenced by the degree of hydrolysis. *Food Chemistry*, 118, 403–410.
- Liu Z.Y., Dong S. Y., Xu J., Zeng M. Y., Song H. X., Zhao Y.H., 2008. Production of cysteine-rich antimicrobial peptide by digestion of oyster (*Crassostrea gigas*) with alcalase and bromelain. *Food Control*, 19, 231–235.
- Lopez-Fandino R., Otte J., Van Camp J., 2006. Physiological, chemical and technological aspects of milk-protein-

- derived peptides with antihypertensive and ACE inhibitory activity. *International Dairy Journal*, 16, 1277–1293.
- Marković R., Baltić M. Ž., Šefer D., Radulović S., Drljačić A., Đorđević V., Ristić M., 2010.** Einfluss der Fütterung auf die Qualität von Broilern: Einfluss erhöhter Mengen an organischem Selen und Vitamin E in der Broile, *Fleischwirtschaft*, 10, 132–136.
- Marx V., 2005.** Watching peptides grow up. *Chemical and Engineering News*, 83, 11, 17–24.
- Matsufuji H., Matsui T., Seki E., Osajima K., Nakashima M., Osajima Y., 1994.** Angiotensin I-converting enzyme inhibitory peptides in an alkaline protease hydrolyzate derived from sardine muscle. *Bioscience, Biotechnology, and Biochemistry*, 58, 12, 2244–2245.
- Matsui T., Matsumoto K., 2006.** Antihypertensive peptides from natural resources. *Advances in Phytomedicine*, 2, 255–271.
- Matsukura T., Tanaka H., 2000.** Applicability of zinc complex of L-carnosine for medical use. *Biochemistry (Moscow)*, 65, 817–823.
- McCann K. B., Shiell B. J., Michalski W. P., Lee A., Wan J., Roginski H., Coventry M. J., 2005.** Isolation and characterisation of antibacterial peptides derived from the f (164–207) region of bovine  $\alpha$  i subS2 sub-casein. *International Dairy Journal*, 15, 2, 133–143.
- McCann K. B., Shiell B. J., Michalski W. P., Lee A., Wan Roginski H., Coventry M. J., 2006.** Isolation and characterization of a novel antibacterial peptide from bovine  $\alpha$ 1-casein. *International Dairy Journal*, 16, 316–323.
- Megías C., Pedroche J., Yust M. M., Girón-Calle J., Alaiz M., Millán F., 2008.** Production of copper-chelating peptides after hydrolysis of sunflower proteins with pepsin and pancreatin. *Food Science and Technology*, 41, 1973–1977.
- Minervini F., Algaron F., Rizzello C. G., Fox P. F., Monnet V., Gobetti M., 2003.** Angiotensin I-converting-enzyme-inhibitory and antibacterial peptides from *Lactobacillus helveticus* PR4 proteinase-hydrolysed casein of milk from six species. *Applied and Environmental Microbiology*, 5297, 5305.
- Möller N. P., Scholz-Ahrens K. E., Roos N., Schrezenmeir J., 2008.** Bioactive peptides and proteins from foods: indication for health effects. *European Journal of Nutrition*, 47, 4, 171–182.
- Morimatsu F., Kimura S., 1992.** *Journal of the Japanese Society for Food Science and Technology* 39, 770–777.
- Morimatsu F., Ito M., Budijanto S., Watanabe I., Furukawa Y., Kimura S., 1996.** Plasma cholesterol-suppressing effect of papain-hydrolyzed pork meat in rats fed hypercholesterolemic diet. *Journal of Nutritional Science and Vitaminology*, 42, 2, 145–153.
- Muguruma M., Ahmed A. M., Katayama K., Kawahara S., Maruyama M., Nakamura T., 2009.** Identification of pro-drug type ACE inhibitory peptide sourced from porcine myosin B: Evaluation of its antihypertensive effects in vivo. *Food Chemistry*, 114, 2, 516–522.
- Mulvihill B., 2004.** Micronutrients in meat. In W. K. Jensen, C. Devine, M. Dikeman (Eds.), *Encyclopedia of meat sciences*, 618–623. Oxford: Elsevier.
- Murray C. J., Lopez A. D., 1997.** Global mortality, disability, and the contribution of risk factors: Global Burden of Disease Study. *Lancet*, 349, 1436–1442.
- Najafian L., Babji A. S., 2012.** A review of fish-derived antioxidant and antimicrobial peptides: their production, assessment, and applications. *Peptides*, 33, 1, 178–185.
- Nakashima Y., Arihara K., Sasaki A., Mio H., Ishikawa S., Itoh M., 2002.** Antihypertensive activities of peptides derived from porcine skeletal muscle myosin in spontaneously hypertensive rats. *Journal of Food Science*, 67, 1, 434–437.
- Nakatani S., Mano H., Sampei C., Shimizu J., Wada M., 2009.** Chondroprotective effect of the bioactive peptide prolyl-hydroxyproline in mouse articular cartilage in vitro and in vivo. *Osteoarthritis and Cartilage*, 17, 1620–1627.
- Ngo D. H., Vo T. S., Ngo D. N., Wijesekara I., Kim S. K., 2012.** Biological activities and potential health benefits of bioactive peptides derived from marine organisms. *International Journal of Biological Macromolecules*, 51, 4, 378–383.
- Olmedilla-Alonso, B., Jiménez-Colmenero, F., Sánchez-Muniz, F. J., 2013.** Development and assessment of healthy properties of meat and meat products designed as functional foods. *Meat Science*, 95, 4, 919–930.
- Ono S., Hosokawa M., Miyashita K., Takahashi K., 2003.** Isolation of Peptides with Angiotensin I-converting Enzyme Inhibitory Effect Derived from Hydrolysate of Upstream Chum Salmon Muscle. *Journal of Food Science*, 68, 5, 1611–1614.
- Onuh J. O., Girgih A. T., Aluko R. E., Aliani, M., 2013.** Inhibitions of renin and angiotensin converting enzyme activities by enzymatic chicken skin protein hydrolysates. *Food Research International*, 53, 1, 260–267.
- Pedroche J., Yust M. M., Lqari H., Megias C., Giron-Calle J., Alaiz M., 2007.** Obtaining of Brassica carinata protein hydrolysates enriched in bioactive peptides using immobilized digestive proteases. *Food Research International*, 40, 7, 931–938.
- Peña-Ramos E. A., Xiong Y. L., Arteaga G. E., 2004.** Fractionation and characterisation for antioxidant activity of hydrolysed whey protein. *Journal of the Science of Food and Agriculture*, 84, 14, 1908–1918.
- Pereira P. M. D. C. C., Vicente A. F. D. R. B., 2013.** Meat nutritional composition and nutritive role in the human diet. *Meat Science*, 93, 3, 586–592.
- Picot L., Bordenave S., Didelot S., Fruitier-Arnaudin I., Sannier F., Thorkelsson G., Piot J. M., 2006.** Antiproliferative activity of fish protein hydrolysates on human breast cancer cell lines. *Process Biochemistry*, 41, 5, 1217–1222.
- Pihlanto A., Akkanen S., Korhonen H. J., 2008.** ACE-inhibitory and antioxidant properties of potato (*Solanum tuberosum*). *Food Chemistry*, 109, 104–112.
- Pihlanto-Leppala A., 2002.** Milk proteins j bioactive peptides. In R. Hubert (Ed.), *Encyclopedia of dairy sciences*, 1960. Oxford: Elsevier.
- Pownall T. L., Udenigwe C. C., Aluko R. E., 2010.** Amino acid composition and antioxidant properties of pea seed (*Pisum sativum* L.) enzymatic protein hydrolysate fractions. *Journal of Agricultural and Food Chemistry*, 58, 4712–4718.
- Purchas R. W., Rutherford S. M., Pearce P. D., Vather R., Wilkinson B. H. P., 2004.** Concentrations in beef and lamb of taurine, carnosine, coenzyme Q(10), and creatine. *Meat Science*, 66, 629–637.
- Purchas R. W., Busboom J., 2005.** The effect of production system and age on levels of iron, taurine, carnosine, coenzyme Q(10), and creatine in beef muscles and liver. *Meat Science*, 70, 589–596.

- Qian Z. J., Je J. Y., Kim S. K., 2007.** Antihypertensive effect of angiotensin I converting enzyme-inhibitory peptide from hydrolysates of bigeye tuna dark muscle, *Thunnus obesus*. *Journal of agricultural and food chemistry*, 55, 21, 8398–8403.
- Qian Z. J., Jung W. K., Kim S. K., 2008.** Free radical scavenging activity of a novel antioxidative peptide purified from hydrolysate of bullfrog skin, *Rana catesbeiana* Shaw. *Bioresour Technology*, 99, 1690–1698.
- Rajapakse N., Mendis E., Jung W. K., Je J. Y., Kim S. K., 2005.** Purification of a radical scavenging peptide from fermented mussel sauce and its antioxidant properties. *Food Research International*, 38, 175–82.
- Revilla E., Maria, C. S., Miramontes E., Bautista J., García-Martínez A., Cremades O., 2009.** Nutraceutical composition, antioxidant activity and hypocholesterolemic effect of a water-soluble enzymatic extract from rice bran. *Food Research International*, 42, 387–393.
- Rho S. J., Lee J. S., Chung Y. I., Kim Y. W., Lee H. G., 2009.** Purification and identification of an angiotensin I-converting enzyme inhibitory peptide from fermented soybean extract. *Process Biochemistry*, 44, 490–493.
- Rutherford-Markwick, K. J., 2012.** Food proteins as a source of bioactive peptides with diverse functions. *British Journal of Nutrition*, 108, 2, 149–157.
- Ryan J. T., Ross R. P., Bolton D., Fitzgerald G. F., Stanton C., 2011.** Bioactive peptides from muscle sources: meat and fish. *Nutrients*, 3, 9, 765–791.
- Saavedra L., Hebert E. M., Minahk C., Ferranti P., 2013.** An overview of omic analytical methods applied in bioactive peptide studies. *Food Research International*, 54, 1, 925–934.
- Saiga A., Okumura T., Makihara T., Katsuta S., Shimizu T., Yamada R., Nishimura, T., 2003a.** Angiotensin I-converting enzyme inhibitory peptides in a hydrolyzed chicken breast muscle extract. *Journal of Agricultural and Food Chemistry*, 51, 6, 1741–1745.
- Saiga A., Tanabe S., Nishimura T., 2003b.** Antioxidant activity of peptides obtained from porcine myofibrillar proteins by protease treatment. *Journal of Agricultural and Food Chemistry*, 51, 3661–3667.
- Saiga A., Okumura T., Makihara T., Katsuda S. I., Morimatsu F., Nishimura T., 2006.** Action mechanism of an angiotensin I-converting enzyme inhibitory peptide derived from chicken breast muscle. *Journal of Agriculture Food Chemistry*, 54, 942–945.
- Saito K., Jin D. H., Ogawa T., Muramoto K., Hatakeyama E., Yasuhara T., 2003.** Antioxidative properties of tripeptide libraries prepared by the combinatorial chemistry. *Journal of Agriculture Food Chemistry*, 51, 3668–3674.
- Sakanaka S., Tachibana Y., 2006.** Active oxygen scavenging activity of egg-yolk protein hydrolysates and their effects on lipid oxidation in beef and tuna homogenates. *Food Chemistry*, 95, 243–249.
- Sarmadi B. H., Ismail A., 2010.** Antioxidative peptides from food proteins: A review. *Pepdies*, 31, 10, 1949–1956.
- Shahidi F., Zhong Y., 2008.** Bioactive peptides. *Journal of AOAC International*, 91, 4, 914–931.
- Shalaby S. M., Zakora M., Otte J., 2006.** Performance of two commonly used angiotensin-converting enzyme inhibition assays using FA-PGG and HHL as substrates. *Journal of Dairy Research*, 73, 2, 178–186.
- Sharp S. I., Aarsland D., Day S., Sonnesyn H., Ballard C., Syst A., 2011.** Hypertension is a potential risk factor for vascular dementia: systematic review. *International Journal of Geriatric Psychiatry*, 26, 661–669.
- Shimizu M., Sawashita N., Morimatsu F., Ichikawa J., Taguchi Y., Ijiri Y., 2009.** Antithrombotic papain-hydrolyzed peptides isolated from pork meat. *Thrombosis Research*, 123, 753–757.
- Siemerink M., Schebb N. H., Liesener A., Perchuc A. M., Schöni, R., Wilmer, M., Vogel, M., 2010.** Development of a fast liquid chromatography/mass spectrometry screening method for angiotensin-converting enzyme (ACE) inhibitors in complex natural mixtures like snake venom. *Rapid Communications in Mass Spectrometry*, 24, 5, 687–697.
- Simitzis P. E., Symeon G. K., Charismiadou M. A., Bizelis J. A., Deligeorgis S. G., 2010.** The effects of dietary oregano oil supplementation on pig meat characteristics. *Meat Science*, 84, 670–676.
- Sofos J. N., 2008.** Challenges to meat safety in the 21<sup>st</sup> century. *Meat Science*, 78, 3–13.
- Suetsuna K., Ukeda H., Ochi H., 2000.** Isolation and characterization of free radical scavenging activities peptides derived from casein. *Journal of Nutritional Biochemistry*, 11, 128–131.
- Sun J., He H., Xie B. J., 2004.** Novel antioxidant peptides from fermented mushroom *Ganoderma lucidum*. *Journal of Agricultural and Food Chemistry*, 52, 21, 6646–6652.
- Takenaka A., Annaka H., Kimura Y., Aoki H., Igarashi K., 2003.** Reduction of paraquat-induced oxidative stress in rats by dietary soy peptide. *Bioscience, Biotechnology, and Biochemistry*, 67, 2, 278–283.
- Terashima M., Baba T., Ikemoto N., Katayama M., Morimoto T., Matsumura S., 2010.** Novel angiotensin-converting enzyme (ACE) inhibitory peptides derived from boneless chicken leg meat. *Journal of Agricultural and Food Chemistry*, 58, 7432–7436.
- Tohidpour A., Sattari M., Omidbaigi R., Yadegar A., Nazemi J., 2010.** Antibacterial effect of essential oils from two medicinal plants against Methicillin-resistant *Staphylococcus aureus* (MRSA). *Phytomedicine*, 17, 142–145.
- Toldra F., Reig M., 2011.** Innovations for healthier processed meats. *Trends in Food Science & Technology*, 22, 9, 517–522.
- Udenigwe C. C., Aluko, R. E., 2012.** Food Protein-Derived Bioactive Peptides: Production, Processing, and Potential Health Benefits. *Journal of Food Science*, 77, 1, R11–R24.
- Udenigwe C. C., Ashton H., 2013.** Meat proteome as source of functional biopeptides. *Food Research International* 54, 1021–1032.
- Vermeirssen V., Van Camp J., Verstraete W., 2002.** Optimisation and validation of an angiotensin-converting enzyme inhibition assay for the screening of bioactive peptides. *Journal of Biochemical and Biophysical Methods*, 51, 1, 75–87.
- Wachtel-Galor S., Szeto Y. T., Tomlinson B., Benzie I. F., 2004.** *Ganoderma lucidum* ('Lingzhi'); acute and short-term biomarker response to supplementation. *International Journal of Food Sciences and Nutrition*, 55, 1, 75–83.
- Wang W., Mejia D., Gonzalez E., 2005.** A New Frontier in Soy Bioactive Peptides that May Prevent Age-related Chronic Diseases. *Comprehensive Reviews in Food Science and Food Safety*, 4, 4, 63–78.

- Weiss J., Gibis M., Schuh V., Salminen H., 2010. Advances in ingredient and processing systems for meat and meat products. *Meat Science*, 86, 1, 196–213.
- Wijesekara I., Kim S. K., 2010. Angiotensin-I-converting enzyme (ACE) inhibitors from marine resources: prospects in the pharmaceutical industry. *Marine Drugs*, 8, 4, 1080–1093.
- Wu H. C., Chen, H. M., Shiau, C. Y., 2003. Free amino acids and peptides as related to antioxidant properties in protein hydrolysates of mackerel (*Scomber austriasicus*). *Food Research International*, 36, 9, 949–957.
- Wu H., He, H. L., Chen, X. L., Sun, C. Y., Zhang, Y. Z., Zhou, B. C., 2008. Purification and identification of novel angiotensin-I-converting enzyme inhibitory peptides from shark meat hydrolysate. *Process Biochemistry*, 43, 4, 457–461.
- Xie Z., Huang J., Xu X., Jin Z., 2008. Antioxidant activity of peptides isolated from alfalfa leaf protein hydrolysate. *Food Chemistry*, 111, 370–376.
- Yokoyama K., Chiba H., Yoshikawa M., 1992. Peptide inhibitors for angiotensin I-converting enzyme from thermolysin digest of dried bonito. *Bioscience, Biotechnology, and Biochemistry*, 56, 10, 1541–1545.
- Young F. J., Therkildsen M., Ekstrand B., Che B. N., Larsen M. K., Oksbjerg N., Stagsted J., 2013. Novel aspects of health promoting compounds in meat, *Meat Science* 95, 904–911.
- Zhu K., Zhou H., Qian H., 2006. Antioxidant and free radical-scavenging activities of wheat germ protein hydrolysates (WGPH) prepared with alcalase. *Process Biochemistry*, 41, 6, 1296–1302.

## Bioaktivni peptidi iz mesa i njihov uticaj na zdravlje ljudi

Baltić Ž. Milan, Bošković Marija, Ivanović Jelena, Janjić Jelena, Dokmanović Marija, Marković Radmila, Baltić Tatjana

*Rezime:* Bioaktivni peptidi predstavljaju funkcionalne komponente unutar proteina i mogu se izolovati iz hrane biljnog i animalnog porekla, uključujući i meso. Nakon oslobađanja iz proteina tokom digestije u gastrointestinalnom traktu, ili nekom od metoda koje se koriste u proizvodnji hrane, ovi peptidi ispoljavaju različite biološke efekte i poseduju različite aktivnosti poput antioksidativne, antimikrobne, antihipertenzivne, antitrombotične, citomodulatorne i imunomodulatorne aktivnosti, a imaju ulogu i u snižavanju nivoa holesterola i borbi protiv kancera i gojaznosti. Aktivnost bioaktivnih peptida zavisi od njihove strukture, ali i drugih karakteristika. Uzimajući u obzir njihove biološke aktivnosti i pozitivan uticaj na ljudsko zdravlje, sa jedne strane, kao i milione smrtnih slučajeva uzrokovanih kancerom, kardiovaskularnim bolestima, kao i drugim bolestima povezanim sa načinom života, sa druge strane, očigledno je da ovi peptidi mogu naći primenu u unapređivanju ljudskog zdravlja i smanjenju rizika od pojave različitih oboljenja. Takođe, bioaktivni peptidi pokazuju određene prednosti u odnosu na sintetičke lekove.

**Ključne reči:** funkcionalna hrana, ACE inhibitorni peptidi, proteini mišića, antioksidativne i antibakterijske osobine.

Paper received: 5.03.2014.

Paper accepted: 14.03.2014.