

Distribution of IS629 and *stx* genotypes among enterohemorrhagic *Escherichia coli* O157 isolates in Yamaguchi Prefecture, Japan, 2004–2013

Mitsuhiro KAMEYAMA^{1)*}, Kiyoshi TOMINAGA¹⁾, Junko YABATA¹⁾ and Yasuharu NOMURA¹⁾

¹⁾Department of Health Science, Yamaguchi Prefectural Institute of Public Health and Environment, 2–5–67 Aoi, Yamaguchi 753–0821, Japan

(Received 22 March 2015/Accepted 24 May 2015/Published online in J-STAGE 6 June 2015)

ABSTRACT. Patterns of insertion sequence (IS)629, *norV* genotype, and Shiga toxin (Stx) genotype distribution were investigated amongst 203 enterohemorrhagic *Escherichia coli* O157 isolates collected in Yamaguchi Prefecture, Japan, between 2004 and 2013. A total of 114 IS629 patterns were identified; these were divided into eight IS groups (A–H). Ninety isolates carried an intact *norV* gene, whereas 113 isolates carried a *norV* with a 204-bp deletion. Other than one isolate from IS group G, all isolates with an intact *norV* belonged to groups A–F, whereas isolates with a mutant *norV* belonged to IS groups G and H. Seven *stx* genotypes were identified, and of those, *stx1a/stx2a* was predominant (n=105), followed by *stx2c* (n=32) and *stx2a* (n=27). The *stx1a/stx2a* genotype was associated with the mutant *norV* isolates, whereas isolates with an intact *norV* had the *stx2c* genotype. Therefore, certain combinations of IS type and *stx* genotype appear to be more frequent among O157 clades which may be useful for detection of predominant subtypes in the interest of public health.

KEY WORDS: enterohemorrhagic *Escherichia coli*, insertion sequence 629, *norV*, O157, *stx* genotype

doi: 10.1292/jvms.15-0166; *J. Vet. Med. Sci.* 77(11): 1437–1441, 2015

Enterohemorrhagic *Escherichia coli* (EHEC), which produces Shiga toxin (Stx), is a human pathogen that causes hemorrhagic colitis, encephalopathy and hemolytic uremic syndrome (HUS) [1, 8]. In Japan, 3,768 cases of EHEC infection were reported in 2012. Among the EHEC isolates collected in 2012, the predominant serogroup was O157 (53%), followed by O26 (27%) and O103 (5%) [9].

Insertion sequences (IS), which are small mobile genetic elements, are widely distributed in bacterial genomes. IS629, a member of the IS3 family of insertion sequences, is prevalent in the O157 genome; for example, the genome of O157 strain Sakai contains 98 IS elements, and of these, 23 were identified as IS629 [2, 10]. O157 isolates show diverse patterns of IS629 insertion, and therefore, IS629 variability can be used for fingerprinting O157 isolates. Ooka *et al.* [12] developed a multiplex PCR tool for screening the distribution of IS629 in the O157 genome.

Several genomic subtyping tools have been developed for the EHEC O157 serogroup. Kulasekara *et al.* [6] showed that two forms of the anaerobic nitric oxide (NO) reductase gene, *norV*, are present in O157 isolates: an intact form and a form with a 204-bp deletion; the intact *norV* is a putative determinant of virulence in some O157 isolates, because NO inhibits Stx2 expression under anaerobic conditions [15]. In addition, Manning *et al.* [7] showed that single nucleotide polymorphisms (SNPs) could be used to classify

O157 strains into nine clades and that HUS patients were significantly more likely to be infected with clade 8 strains. Further, a lineage-specific polymorphism assay can be used to classify O157 isolates (lineages I, II and I/II), and lineage I isolates are more commonly associated with human disease than lineage II isolates [5, 13, 16, 18].

Some IS elements are thought to play an important role in the diversification and evolution of bacteria, including those of EHEC O157 isolates [11, 13]. Yokoyama *et al.* [17] and Hirai *et al.* [3] showed a biased distribution of IS629 among different lineages or clades of O157 isolates. However, the isolates examined in these studies were collected from a limited number of geographical areas, and the distribution of IS629 among O157 isolates from Yamaguchi Prefecture, Japan, has not been investigated. In the present study, we investigated the distribution of IS629 as well as *norV* types in EHEC O157 isolates collected in Yamaguchi Prefecture between 2004 and 2013. We also examined the association between *stx* genotype and O157 phylogeny.

MATERIALS AND METHODS

EHEC O157 isolates: A total of 203 EHEC O157 isolates were used in the present study. The isolates were sent to our laboratory from hospitals and health care centers in Yamaguchi Prefecture, Japan, between 2004 and 2013, and originated from epidemiologically unrelated patients (189 isolates) and asymptomatic carriers (14 isolates). The clinical symptoms of the patients included watery diarrhea (n=165), abdominal pain (n=144), bloody diarrhea (n=115), fever (n=59) and vomiting (n=31). Five patients developed HUS. All isolates were negative for sorbitol fermentation, and 192 isolates were serotyped as O157:H7. The remaining isolates were non-motile and were therefore classified as O157:NM.

*CORRESPONDENCE TO: KAMEYAMA, M., Department of Health Science, Yamaguchi Prefectural Institute of Public Health and Environment, 2–5–67 Aoi, Yamaguchi 753–0821, Japan.
e-mail: kameyama.mitsuhiro@pref.yamaguchi.lg.jp

©2015 The Japanese Society of Veterinary Science

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives (by-nc-nd) License <<http://creativecommons.org/licenses/by-nc-nd/3.0/>>.

DNA preparation: To extract DNA for PCR analysis, individual isolates were cultured on Mueller-Hinton agar (Oxoid, Basingstoke, U.K.), and the DNA was extracted using a QIAamp DNA Blood mini kit (Qiagen, Hilden, Germany), according to the manufacturer's instructions. Prepared DNA was stored at -20°C until use.

IS629 typing: Multiplex PCR using an O157 IS-printing system (Toyobo, Osaka, Japan), which can be used to detect all 32 IS629 loci, was used to type the isolates based on the number and location of the insertion sequences. The assay also screens for the presence of the four virulence genes, *eaec*, *hlyA*, *stx1* and *stx2*. The resulting banding patterns were analyzed using BioNumerics software v. 7.1 (Applied Maths, Sint-Martens-Latem, Belgium), which excluded bands corresponding to the 4 virulence genes. A simple matching coefficient and an unweighted pair group method with arithmetic mean (UPGMA) algorithm were used to generate a dendrogram. For the purposes of this study, isolates with $\geq 80\%$ similarity were considered to belong to the same group.

Detection of the intact and deletion variants of *norV*: The 2 versions of *norV*, the intact version and the version containing a 204-bp deletion, were detected by PCR, as described previously [4].

Detection of isolates belonging to clade 8: We screened for the presence of clade 8 isolates using mismatch amplification mutation assay (MAMA) PCR, as described previously [4].

***stx* genotyping:** The *stx* subtypes, *stx1a*, *stx1c*, *stx1d* and *stx2a-stx2g*, were detected by PCR, as described previously [14].

RESULTS

IS629 distribution: A total of 114 IS629 patterns were distinguished amongst the 203 O157 isolates. As shown in Fig. 1, the isolates could be divided into 8 groups (groups A–H). Group G was predominant ($n=103$), followed by group C ($n=35$). Isolates belonging to each of these 2 groups were identified in all years of the study (Table 1).

Detection of *norV* types: Intact *norV* was detected in 90 isolates (intact *norV*-type), and *norV* with the 204-bp deletion was detected in 113 isolates (deletion *norV*-type). All isolates belonging to groups A–F, and one isolate from group G, carried the intact *norV* (Fig. 1 and Table 1).

Clade 8 isolates: In total, 15 of the 203 O157 isolates were determined to belong to clade 8. All of these isolates contained an intact *norV* gene and were classified into IS group E.

***stx* genotypes:** The *stx1a*, *stx2a* and *stx2c* subtypes were detected in 123, 154 and 68 isolates, respectively. The *stx1a/stx2a* genotype was predominant ($n=105$), followed by *stx2c* ($n=32$), *stx2a* ($n=27$), *stx2a/stx2c* ($n=21$), *stx1a/stx2c* ($n=14$) and *stx1a* ($n=3$). Only one isolate carried all three *stx* subtypes, *stx1a*, *stx2a* and *stx2c* (Table 2).

Among the seven *stx* genotypes detected, the *stx2a* genotype was common in both intact *norV*-type and deletion *norV*-type isolates. The *stx2c*, *stx1a/stx2c*, *stx2a/stx2c* and *stx1a/stx2a/stx2c* genotypes were common in the intact

norV-type isolates, whereas the *stx1a* and *stx1a/stx2a* genotypes were associated with the deletion *norV*-type isolates.

Of the 15 clade 8 isolates, 7 showed the *stx2a* genotype, and 8 showed the *stx2a/stx2c* genotype.

DISCUSSION

Various IS629 patterns were observed among the O157 isolates collected in Yamaguchi Prefecture, Japan, between 2004 and 2013. The majority of the isolates collected over this 10-year period belonged to IS groups C and G, although isolates from 8 different groups were identified. As IS629 is thought to contribute to the diversity of O157 isolates, it is not surprising that isolates with distinct IS patterns are distributed in Yamaguchi Prefecture.

Recent studies demonstrated that the distribution of IS629 was biased among O157 clades or lineages [3, 17]. Iyoda *et al.* [4] also revealed that O157 clades were associated with certain *norV* types, with isolates belonging to clades 4–8 carrying an intact *norV*, whereas isolates belonging to clades 1–3 contained the deletion type. Our results showed that the 15 clade 8 isolates shared identical or highly similar IS patterns and belonged to a single IS group (group E). Moreover, with the exception of one isolate, the intact *norV*-type and deletion *norV*-type isolates were clearly divided into distinct IS groups (IS groups A–F and G–H, respectively). Therefore, our findings, along with those of previous studies [3, 4, 17], indicate that it is rare for isolates belonging to different clades to share similar IS profiles.

Kulasekara *et al.* [6] reported that most deletion *norV*-type isolates contained *stx1a*, whereas only 10% of intact *norV*-type isolates harbored *stx1a*. In the present study, 95.6% of deletion *norV*-type isolates and 16.7% of intact *norV*-type isolates carried *stx1a*, in accord with the previous study. Moreover, all isolates carrying *stx2c* also carried an intact *norV*. A previous study demonstrated that 98.8% of all *stx2c*-positive isolates carried an intact *norV* and belonged to clades 4–8 [4]. Thus, *stx* genotypes appear to be associated with specific O157 clades.

An additional association between *stx* genotypes and IS distribution was observed in this study. Among the seven *stx* genotypes identified, the predominant genotype, *stx1a/stx2a*, was only identified in isolates in IS groups G and H, whereas the 4 genotypes including the *stx2c* subtype were identified in IS groups A–F. Thus, particular *stx* genotypes may be associated with O157 isolates with particular IS629 distributions.

Five of the 203 isolates tested in the present study were obtained from HUS patients. These 5 isolates did not belong to clade 8 and were genotyped as *stx1a/stx2a* ($n=3$, deletion *norV*-type) and *stx2a/stx2c* ($n=2$, intact *norV*-type). Similar results were obtained by Iyoda *et al.* [4], who found that isolates with these 2 genotypes were significantly more likely to be associated with HUS patients than with asymptomatic carriers. Thus, the *stx1a/stx2a* and *stx2a/stx2c* genotypes may be associated with an increased risk of developing HUS. However, Manning *et al.* [7] also investigated the association between O157 clades, *stx* subtypes and HUS and




Fig. 1. Cluster analysis of 203 enterohemorrhagic *Escherichia coli* O157 isolates based on IS-printing profiles. Isolates with >80% similarity were considered to belong to the same group. The hatched bar indicates the intact *norV*, and the open bar indicates the 204-bp-deletion *norV*. A “☆” indicates a HUS patient, and “★” indicates an asymptomatic carrier.

Table 1. Distribution of O157 isolates according to IS groups by year

IS group	norV ^{a)}	Clade ^{b)}	n	No. of isolates in each year									
				2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
A	Intact	-	18	5			4	2	2	4		1	
B	Intact	-	1				1						
C	Intact	-	35	4	1	1	5	7	5	3	3	3	3
D	Intact	-	1	1									
E	Intact	-	3				1			2			
		clade 8	15	3	1	1	1		1	3	4		1
F	Intact	-	16	2	1	1	4	3	1	3	1		
G	Intact	-	1					1					
	Deletion	-	102	14	15	8	10	9	13	8	8	9	
H	Deletion	-	11			1		3	2		2	2	1
Total			145	29	18	12	26	25	24	23	18	15	13

a) Intact or 204-bp-deletion *norV* gene. b) “-” indicates non-clade 8 isolates.

Table 2. *stx* genotypes of intact or deletion *norV*-type isolates

<i>norV</i> type ^{a)}	No. of isolates in each <i>stx</i> genotype							
	<i>stx1a</i>	<i>stx2a</i>	<i>stx2c</i>	<i>stx1a/stx2a</i>	<i>stx1a/stx2c</i>	<i>stx1a/stx2a/stx2c</i>	<i>stx2a/stx2c</i>	
Intact	0	21	32	0	14	1		21
Deletion	3	6	0	105	0	0		0
Total	3	27	32	105	14	1		21

a) Intact or 204-bp-deletion *norV* gene.

found that *stx* type alone cannot account for the variation in hospitalization and HUS rates by clade.

In the present study, the diversity of IS629 patterns and *stx* genotypes in O157 isolates collected in Yamaguchi Prefecture, Japan, were determined. The clade 8 isolates, which are considered to show high virulence, shared a particular IS629 distribution and *stx* genotype. Because analysis of patterns of IS629 distribution and *stx* genotype in O157 isolates can help to determine whether or not an isolate has high virulence in humans, it is important to identify the characteristics of O157 isolates to alert clinical laboratories to the risk of developing severe diseases, such as HUS.

ACKNOWLEDGMENT. This work was supported in part by a grant-in-aid for Research on Emerging and Re-emerging Infectious Diseases (H24-Shinkou-Ippan-005) from the Ministry of Health, Labour and Welfare, Japan.

REFERENCES

- Feng, P. 1995. *Escherichia coli* serotype O157:H7: novel vehicles of infection and emergence of phenotypic variants. *Emerg. Infect. Dis.* **1**: 47–52. [Medline] [CrossRef]
- Hayashi, T., Makino, K., Ohnishi, M., Kurokawa, K., Ishii, K., Yokoyama, K., Han, C. G., Ohtsubo, E., Nakayama, K., Murata, T., Tanaka, M., Tobe, T., Iida, T., Takami, H., Honda, T., Sasakawa, C., Ogasawara, N., Yasunaga, T., Kuhara, S., Shiba, T., Hattori, M. and Shinagawa, H. 2001. Complete genome sequence of enterohemorrhagic *Escherichia coli* O157:H7 and genomic comparison with a laboratory strain K-12. *DNA Res.* **8**: 11–22. [Medline] [CrossRef]
- Hirai, S., Yokoyama, E. and Yamamoto, T. 2013. Linkage disequilibrium of the IS629 insertion among different clades of enterohemorrhagic *Escherichia coli* O157:H7/H- strains. *Infect. Genet. Evol.* **18**: 94–99. [Medline] [CrossRef]
- Iyoda, S., Manning, S. D., Seto, K., Kimura, K., Isobe, J., Etoh, Y., Ichihara, S., Migita, Y., Ogata, K., Honda, M., Kubota, T., Kawano, K., Matsumoto, K., Kudaka, J., Asai, N., Yabata, J., Tominaga, K., Terajima, J., Morita-Ishihara, T., Izumiya, H., Ogura, Y., Saitoh, T., Iguchi, A., Kobayashi, H., Hara-Kudo, Y. and Ohnishi, M., EHEC Working Group in Japan. 2014. Phlogenetic clades 6 and 8 of enterohemorrhagic *Escherichia coli* O157:H7 with particular *stx* subtypes are more frequently found in isolates from hemolytic uremic syndrome patients than from asymptomatic carriers. *Open Forum Infect. Dis.* Summer 2014. (first published online July 18, 2014. doi: [CrossRef]). [Medline]
- Kim, J., Nietfeldt, J., Ju, J., Wise, J., Fegan, N., Desmarchelier, P. and Benson, A. K. 2001. Ancestral divergence, genome diversification, and phylogeographic variation in subpopulation of sorbitol-negative, beta-grucuronidase-negative enterohemorrhagic *Escherichia coli* O157. *J. Bacteriol.* **183**: 6885–6897. [Medline] [CrossRef]
- Kulasekara, B. R., Jacobs, M., Zhou, Y., Wu, Z., Sims, E., Saenphimmachak, C., Rohmer, L., Ritchie, J. M., Radey, M., McKeivitt, M., Freeman, T. L., Hayden, H., Haugen, E., Gillett, W., Fong, C., Chang, J., Beskhlebna, V., Waldor, M. K., Samadpour, M., Whittam, T. S., Kaul, R., Brittnacher, M. and Miller, S. I. 2009. Analysis of the genomes of the *Escherichia coli* O157:H7 2006 spinach-associated outbreak isolate indicates candidate genes that may enhance virulence. *Infect. Immun.* **77**: 3713–3721. [Medline] [CrossRef]
- Manning, S. D., Motiwala, A. S., Springman, A. C., Qi, W., Lacher, D. W., Ouellette, L. M., Mladonicky, J. M., Somsel, P., Rudrik, J. T., Dietrich, S. E., Zhang, W., Swaminathan, B., Alland, D. and Whittam, T. S. 2008. Variation in virulence among

- clades of *Escherichia coli* O157:H7 associated with disease outbreaks. *Proc. Natl. Acad. Sci. U.S.A.* **105**: 4868–4873. [[Medline](#)] [[CrossRef](#)]
8. Mellmann, A., Bielaszewska, M., Köck, R., Friedrich, A. W., Fruth, A., Middendorf, B., Harmsen, D., Schmidt, M. A. and Karch, H. 2008. Analysis of collection of hemolytic uremic syndrome-associated enterohemorrhagic *Escherichia coli*. *Emerg. Infect. Dis.* **14**: 1287–1290. [[Medline](#)] [[CrossRef](#)]
 9. National Institute of Infectious Diseases. 2013. Enterohemorrhagic *Escherichia coli* infections in Japan as of April 2013. *Infect. Agents Surveillance Rep.* **34**: 123–124.
 10. Ohnishi, M., Terajima, J., Kurokawa, K., Nakayama, K., Murata, T., Tamura, K., Ogura, Y., Watanabe, H. and Hayashi, T. 2002. Genomic diversity of enterohemorrhagic *Escherichia coli* O157 revealed by whole genome PCR scanning. *Proc. Natl. Acad. Sci. U.S.A.* **99**: 17043–17048. [[Medline](#)] [[CrossRef](#)]
 11. Ooka, T., Ogura, Y., Asadulghani, M., Ohnishi, M., Nakayama, K., Terajima, J., Watanabe, H. and Hayashi, T. 2009. Inference of the impact of insertion sequence (IS) elements on bacterial genome diversification through analysis of small-size structural polymorphisms in *Escherichia coli* O157 genomes. *Genome Res.* **19**: 1809–1816. [[Medline](#)] [[CrossRef](#)]
 12. Ooka, T., Terajima, J., Kusumoto, M., Iguchi, A., Kurokawa, K., Ogura, Y., Asadulghani, S., Nakayama, K., Murase, K., Ohnishi, M., Iyoda, S., Watanabe, H. and Hayashi, T. 2009. Development of a multiplex PCR-based rapid typing method for enterohemorrhagic *Escherichia coli* O157 strains. *J. Clin. Microbiol.* **47**: 2888–2894. [[Medline](#)] [[CrossRef](#)]
 13. Rump, L. V., Fischer, M. and Gonzalez-Escalona, N. 2011. Prevalence, distribution and evolutionary significance of the IS629 insertion element in the stepwise emergence of *Escherichia coli* O157:H7. *BMC Microbiol.* **11**: 133. [[Medline](#)] [[CrossRef](#)]
 14. Scheutz, F., Teel, L. D., Beutin, L., Piérard, D., Buvens, G., Karch, H., Mellmann, A., Caprioli, A., Tozzoli, H., Morabito, S., Strockbine, N. A., Melton-Celsa, A., Sanchez, M., Persson, S. and O'Brien, A. D. 2012. Multicenter evaluation of a sequence-based protocol for subtyping Shiga toxins and standardizing Stx nomenclature. *J. Clin. Microbiol.* **50**: 2951–2963. [[Medline](#)] [[CrossRef](#)]
 15. Shimizu, T., Tsutsuki, H., Matsumoto, A., Nakaya, H. and Noda, M. 2012. The nitric oxide reductase of enterohemorrhagic *Escherichia coli* plays an important role for the survival within macrophages. *Mol. Microbiol.* **85**: 492–512. [[Medline](#)] [[CrossRef](#)]
 16. Yang, Z., Kovar, J., Kim, J., Nietfeldt, J., Smith, D. R., Moxley, R. A., Olson, M. E., Fey, P. D. and Benson, A. K. 2004. Identification of common subpopulations of non-sorbitol-fermenting, β -glucuronidase-negative *Escherichia coli* O157:H7 from bovine production environments and human clinical samples. *Appl. Environ. Microbiol.* **70**: 6846–6854. [[Medline](#)] [[CrossRef](#)]
 17. Yokoyama, E., Hashimoto, R., Etoh, Y., Ichihara, S., Horikawa, K. and Uchimura, M. 2011. Biased distribution of IS629 among strains in different lineages of enterohemorrhagic *Escherichia coli* serovar O157. *Infect. Genet. Evol.* **11**: 78–82. [[Medline](#)] [[CrossRef](#)]
 18. Zhang, Y., Laing, C., Steele, M., Ziebell, K., Johnson, R., Benson, A. K., Taboada, E. and Gannon, V. P. J. 2007. Genomic evolution in major *Escherichia coli* O157:H7 lineages. *BMC Genomics* **8**: 121. [[Medline](#)] [[CrossRef](#)]