

Diet, *Escherichia coli* O157:H7, and Cattle: A Review After 10 Years

Todd R. Callaway^{1*}, M. A. Carr², T. S. Edrington¹,
Robin C. Anderson¹, and David J. Nisbet¹

¹Food and Feed Safety Research Unit, Southern Plains Agricultural Research Center, Agricultural Research Service, USDA, College Station, TX 77845, USA

²National Cattlemen's Beef Association, Centennial, CO, USA

Abstract

Escherichia coli are commensal bacteria that can account for up to 1% of the bacterial population of the gut. Ruminant animals are reservoirs of the pathogenic bacteria *E. coli* strain O157:H7, and approximately 30% of feedlot cattle shed *E. coli* O157:H7. Feedlot and high-producing dairy cattle are fed high grain rations in order to increase feed efficiency. When cattle are fed high grain rations, some starch escapes ruminal microbial degradation and passes to the hindgut where it undergoes fermentation. Ten years ago researchers demonstrated that populations of total *E. coli* were higher in grain-fed than in forage-fed cattle, and when cattle were abruptly switched from a high grain diet to an all hay diet, total *E. coli* populations declined 1000-fold within 5 days and reduced the ability of the surviving *E. coli* to survive an acid shock mimicking passage through the human gastric stomach. This research provoked many questions about the effects of diet or *E. coli* O157:H7 populations that have not been conclusively answered to date. Subsequent research has shown that diet does affect *E. coli* O157:H7 populations, but the effects have varied in magnitude and impact. Further studies have demonstrated that the effects of forage feeding on *E. coli* O157:H7 populations may be due to concentrations of tannins and phenolic acids in forages. Other ration components such as rapidly ruminally fermented grains (e.g., barley) increase the shedding of *E. coli* O157:H7, and in some situations, feeding distillers grains can increase fecal shedding of *E. coli* O157:H7 due to VFA concentrations. Data from researchers across North America indicate that diet does impact STEC/EHEC populations in cattle prior to slaughter; however the economic, logistic and practical impacts of dietary changes must be examined and accounted for.

Disclaimer: "Proprietary or brand names are necessary to report factually on available data; however, the USDA neither guarantees nor warrants the standard of the product, and the use of the name by the USDA implies no approval of the product, and exclusion of others that may be suitable."

*Corresponding author: Email: Todd.Callaway@ars.usda.gov Phone: +1 (979) 260-9374; Fax: +1 (979) 260-9332.

Introduction

Escherichia coli is a facultative anaerobic bacterium commonly found in the mammalian intestinal tract (Drasar and Barrow, 1985). *Escherichia coli* lives a fecal-oral lifestyle and can comprise up to 1% of the gastrointestinal population of mammals and is used as an indicator of environmental fecal contamination of water supplies (Winfield and Groisman, 2003). Most *E. coli* strains are commensal; however, some *E. coli* strains can be pathogenic to humans, and are harbored within food animals (Drasar, 1974, Drasar and Barrow, 1985). Some *E. coli* strains can cause hemorrhagic colitis in humans; but the best known enterohemorrhagic *E. coli* (EHEC) strain remains O157:H7 (Scotland et al., 1990). Each year, more than 60 people die and 73,000 people are made ill by *Escherichia coli* O157:H7 in the U.S. (Mead et al., 1999), and EHEC infections are estimated to cost the economy more than \$1 billion per year (USDA-ERS, 2001).

Ground beef is the most frequently implicated source of *E. coli* O157:H7 outbreaks, and bovine-derived products are linked to approximately 75% of *E. coli* O157:H7 outbreaks (USDA-APHIS, 1997, Vugia et al., 2007). Cattle are a major reservoir of *E. coli* O157:H7 and repeated hemorrhagic colitis outbreaks linked to consumption of ground beef, animal contact, manure amendment, or cattle manure-contaminated runoff has firmly established the connection between cattle and *E. coli* O157:H7 epidemiologically, and in public perception (Jay et al., 2007, Keen et al., 2007, Steinmuller et al., 2006). Repeated large-scale recalls of *E. coli* O157:H7 contaminated ground beef, and the well-publicized deaths of children who consumed foods contaminated by exposure to beef products or ruminants have further shaken the confidence of consumers in the wholesomeness and safety of beef (Gage, 2001). In-plant post-harvest sanitation efforts effectively reduce contamination of carcasses with *E. coli* O157:H7 (Arthur et al., 2007a, Bosilevac et al., 2006, Woerner et al., 2006). However, no matter the effectiveness of these strategies, they are not sufficient to ensure human food safety and health.

Studies have shown that up to 30% of all cattle are asymptomatic carriers of *E. coli* O157:H7 (Callaway et al., 2006, Reinstein et al., 2007, Stanford et al., 2005). Manure from cattle production facilities can contain viable *E. coli* O157:H7 and be washed into the water supply and consumed directly in drinking water, or be used as irrigation water on crops, or transmitted by other animal vectors (Hill et al., 2006, LeJeune et al., 2001, Sargeant et al., 2003, Thurston-Enriquez et al., 2005). Therefore, methods that reduce *E. coli* O157:H7 populations in food animals prior to entry to the food chain have great potential to reduce human illnesses (Callaway et al., 2004, Loneragan and Brashears, 2005, Sargeant et al., 2007).

One method proposed to reduce EHEC in cattle is to abruptly change the diet from a high grain to a forage based ration. This suggestion was based on research results first published in 1998 that demonstrated that an abrupt shift from grain to hay-based rations reduced generic *E. coli* populations significantly (Diez-Gonzalez et al., 1998). This study elicited a great deal of subsequent research that has yielded variable results (Hancock et al., 2000, Hovde et al., 1999, Keen et al., 1999). Therefore, as the 10 year anniversary of this hypothesis is reached, this review examines the current state of knowledge about the effects of dietary and other cattle management manipulations on *E. coli* and O157:H7 populations.

Enterohaemorrhagic *E. coli* and their reservoir, ruminant animals

The microbial population of the ruminant is diverse and microbes live throughout the gastrointestinal tract of mammals, and this includes the ubiquitous and adaptable *E. coli* (Drasar and Barrow, 1985, Yokoyama and Johnson, 1988). *Escherichia coli* are rarely cultured in high numbers from the rumen of cattle, typically less than 10^6 cells/ml out of a population of $>10^{10}$ cells/ml (Laven et al., 2003, Min et al., 2007, Wolin, 1969). *Escherichia coli* were not considered "important" ruminal bacteria (Rasmussen et al., 1999, Wolin, 1969) because of the competitive nature of the rumen and the fact that high concentrations of volatile fatty acids (VFA) are bactericidal (Wallace et al., 1989, Wolin, 1969). In the lower intestinal tract, conditions are generally more favorable for *E. coli*, where they can be found at concentrations ranging from 10^2 to 10^7 cells/g feces at slaughter (Davidson and Taylor, 1978, Jordan and McEwen, 1998). In spite of the severity of illness in humans caused by EHEC, they are not predominant members of the *E. coli* population (Laven et al., 2003, Tkalcic et al., 2003, Zhao et al., 1998). In addition to gastrointestinal populations, the oral cavities of cattle also can contain EHEC (Keen and Elder, 2002, Smith et al., 2005), likely due to the process of rumination. However, it must be noted that the genotype of *E. coli* O157:H7 isolates from the oral cavity can differ significantly from fecal isolates (Keen and Elder, 2002). Researchers have shown that *E. coli* were associated with the digesta rather than with the intestinal wall (Laven et al., 2003). Other studies have demonstrated that the terminal end of the colon was the major site of *E. coli* O157:H7 colonization (Grauke et al., 2002) and the lymphoid tissue located at the recto-anal junction (RAJ) has been demonstrated to be the primary site of colonization of cattle (Naylor et al., 2003). Other researchers have subsequently verified that the RAJ is intimately involved with *E. coli* O157:H7 colonization (Davis, 2006, Greenquist et al., 2005, Lim et al., 2007, Rice et al., 2003) leading to the suggestion that RAJ colonization may be involved in the phenomenon of "super-shedders" (Cobbold, 2007, LeJeune and Kauffman, 2006).

The EHEC strain responsible for most human illnesses and that colonize cattle vary geographically and temporally (Cookson et al., 2007, Fagan et al., 1999, Wang et al., 2000). Studies have found cattle can be concomitantly colonized by up to 26 different serotypes of EHEC (Schurman et al., 2000). In the U.S. most illnesses and subsequent surveys and interest have focused on strain O157:H7, although

other EHEC are found in the U.S. in humans and cattle (Hussein et al., 2003b, Midgley et al., 1999). Therefore it is emphasized that although the U.S. focus remains on strain O157:H7, that we not forget the other EHEC (Acheson, 2000), because the natural ecology of EHEC suggests that if strain O157:H7 decreases then another EHEC strain (or another pathogen) would simply fill the vacuum.

Shedding of EHEC in cattle appears to be widespread, but sporadic (Meyer-Brosseta et al., 2001). EHEC shedding is highly dependent on season of the year (Gyles, 2007, Sargeant et al., 2007) and can range from as many as 80% of all feedlot cattle in the summer, to as few as 5-10% shedding during the winter (Barkocy-Gallagher et al., 2003, Elder et al., 2000, Naumova et al., 2007). This correlates with an increase in human outbreaks during each summer/early fall, thus highlighting a linkage between the animal (reservoir) populations and consumers via food-borne outbreaks (USDA-APHIS, 1997, Vugia et al., 2007). Seasonality of shedding has long been theorized to be related to temperature or weather (Barkocy-Gallagher et al., 2003, Naumova et al., 2007); however a recent theory has emerged that day length and melatonin or seasonal hormones may play a role in this phenomenon, however much further research is needed to investigate this intriguing hypothesis (Edrington et al., 2007, Edrington et al., 2006a, Schultz et al., 2005).

Hides a reservoir of E. coli O157:H7

Bovine manure can harbor *E. coli* O157:H7 at the typical environmental temperatures for ≥ 49 d (Wang et al., 1996). Dirt and feces that collects on the hides of cattle can therefore be contaminated with *E. coli* O157:H7 for long periods of time (Arthur et al., 2007b, Barkocy-Gallagher et al., 2004, Keen and Elder, 2002, Reid et al., 2002). Research has indicated that the number of hides positive for *E. coli* O157:H7 is a more accurate predictor for carcass contamination than is fecal prevalence (Barkocy-Gallagher et al., 2003). The incidence of hides contaminated by *E. coli* O157:H7 in studies ranged from 14% to 85% (Elam et al., 2003, Stephens et al., 2007, Woerner et al., 2006, Younts-Dahl et al., 2004). Recently it was demonstrated that when a pen had a $>20\%$ fecal incidence rate, then the percentage of hides positive for *E. coli* O157:H7 rose to 26%; however when the fecal incidence rate in the pens was $<20\%$ then the hide contamination level was only 5% (Woerner et al., 2006). Thus a common-sense approach to decreasing the prevalence of *E. coli* O157:H7 on hides has led to the development of methods to remove the hair from hides prior to dehiding/evisceration (Castillo et al., 1998, Nou et al., 2003) or to specialized treatments that reduce the pathogen load on the hide (Bosilevac et al., 2005, Bosilevac et al., 2004). Gregory et al. (2000) found that when cattle arrived at the slaughter plant the hides of cattle fed hay for 48 h prior to transport were as clean as the hides of fasted cattle, and were significantly cleaner than pasture-fed cattle (Gregory et al., 2000).

Do *E. coli* O157:H7 levels in the live animal really matter?

Data indicate that in-plant intervention strategies reduce the spread of *E. coli* O157:H7 on and between carcasses (Arthur et al., 2002, Barkocy-Gallagher et al., 2003, Bosilevac et al.,

2006, Woerner *et al.*, 2006). If pre-harvest interventions can be introduced to reduce levels of *E. coli* O157:H7 that enter the abattoir within the live animal, we should be able to further enhance the effectiveness of in-plant intervention strategies (Callaway *et al.*, 2004, Loneragan and Brashears, 2005). Thus reducing the burden of pathogens entering the abattoir should enhance human health (Hynes and Wachsmuth, 2000). However direct infection via food is not the only route of human exposure to foodborne pathogenic bacteria.

The increasing disconnect between the consumer and their agricultural food supply is reflected in a growing number of direct-contact illnesses in humans contracted in farmyards, open farms, petting zoos, and zoological parks (Chapman *et al.*, 2000, Edwards *et al.*, 2008, Keen *et al.*, 2007, Keen *et al.*, 2003, Pritchard *et al.*, 2000). Recent years have seen an increase in human *E. coli* O157:H7 illnesses linked to water contaminated by run-off from feedlots and dairies (Anonymous, 2000, Edwards *et al.*, 2008, Goss and Richards, 2007, Jay *et al.*, 2007). The spread of *E. coli* O157:H7 in runoff from farms has only been recently assessed to understand the movement of pathogens from farms during rainfall events (Berry *et al.*, 2007, Ferguson *et al.*, 2007). Further concerns have been raised about the spread of *E. coli* O157:H7 to humans through crops (e.g., spinach or lettuce) irrigated with water from cattle farms (Gerba and Smith, 2005, Manshadi *et al.*, 2001, Natvig *et al.*, 2002). The recent (2006), widespread *E. coli* O157:H7 outbreaks linked to spinach and lettuce (Jay *et al.*, 2007) contaminated by swine that had foraged on a dairy highlights the ability of EHEC to cause human illnesses through a variety of vectors, further emphasizing the need to reduce foodborne pathogenic bacteria in the live animal before they contact human consumers (Hynes and Wachsmuth, 2000, Loneragan and Brashears, 2005, Sargeant *et al.*, 2007).

Dietary effects on *E. coli* O157:H7 shedding in cattle

Fasting

Before and during transport, cattle can at times be fasted for up to 48 h, which can affect their susceptibility to colonization by *E. coli* O157:H7. Ruminal and intestinal VFA concentrations limit *E. coli* populations because VFA are toxic (Hollowell and Wolin, 1965, Russell and Diez-Gonzalez, 1998, Wolin, 1969). This has allowed VFA and other organic acids to be used to reduce pathogen populations in the gut (Ohya *et al.*, 2000, Prohaszka and Baron, 1983, Van Immerseel *et al.*, 2006). Because feed withdrawal and/or starvation results in decreased VFA concentrations in the gut, it has been suggested that this shift plays a role in the effects of transport on the shedding of EHEC.

Fasting increased *E. coli*, *Enterobacter* and total anaerobic bacterial populations throughout the intestinal tract (Buchko *et al.*, 2000b, Gregory *et al.*, 2000), and increased *Salmonella* and *E. coli* populations in the rumen (Brownlie and Grau, 1967, Grau *et al.*, 1969), furthermore, fasting can cause “apparently *E. coli* (O157:H7) negative animals to become positive” (Kudva *et al.*, 1995). Further studies indicated that fasting made calves more susceptible to colonization by inoculated *E. coli* O157:H7, and demonstrated that fasted inoculated calves shed more *E. coli* O157:H7 than did

calves fed normally (Cray *et al.*, 1998). Cattle fasted for 48 h prior to slaughter also were shown to contain significantly greater *E. coli* populations throughout the gut than cattle fed hay or pasture (Gregory *et al.*, 2000). In contrast, it was demonstrated that a fasting period reduced ruminal VFA concentrations, but this did not influence *E. coli* O157:H7 shedding (Harmon *et al.*, 1999). In general, studies examining the intestinal environment have repeatedly indicated that low pH and high concentrations of short chain VFA result in lower EHEC populations (Bach *et al.*, 2002a, Bach *et al.*, 2005b, Cobbold and Desmarchelier, 2004, Shin *et al.*, 2002). Thus research indicates fasting increases shedding or makes cattle more susceptible to colonization due to decreased short chain VFA and increased pH in the gastrointestinal tract.

Can feed additives and antimicrobials affect *E. coli* O157:H7?

Ionophores, such as monensin and lasalocid, are included in most feedlot and dairy rations to inhibit gram-positive bacteria, thereby improving feed:gain ratios and production efficiency (Callaway *et al.*, 2003). Because these feed additives can alter the microbial population, possibly giving gram-negative bacteria (such as *E. coli*) a competitive advantage, they were an obvious risk factor to investigate in regard to their role in colonization and shedding of *E. coli* O157:H7. As expected, based upon the gram-negative membrane physiology of *E. coli* O157:H7, the most widely used ionophore (monensin) did not affect the growth or growth rate of this pathogen in vitro when added at concentrations similar to that found in the rumen of cattle fed monensin (Bach *et al.*, 2002b), subsequently *E. coli* O157:H7 has been shown to be resistant to concentrations of monensin as high as 3-fold higher than that normally found in the rumen (Van Baale *et al.*, 2004). Ionophoric feed additives (monensin, lasalocid, laidlomycin and bambarmycin) demonstrated no effect on *E. coli* O157:H7 in vitro (Edrington *et al.*, 2003).

Early epidemiological studies demonstrated a marginally significant increase of EHEC shedding by heifers fed ionophores (Herriott *et al.*, 1998), but others could not draw any conclusions (Dargatz *et al.*, 1997). Cattle fed a forage ration that included monensin shed *E. coli* O157:H7 for a shorter period of time than forage-fed cattle not supplemented with monensin, but monensin had no effect on shedding when cattle were fed a corn-based ration (Van Baale *et al.*, 2004). In a recent in vitro study, it was found that monensin and the co-approved antibiotic tylosin (tylan) treatment reduced *E. coli* O157:H7 populations up to 2 log₁₀ CFU/ml in ruminal fermentations from cows fed forage, but did not affect *E. coli* O157:H7 populations in ruminal fluid from cows fed corn (McAllister *et al.*, 2006). The inclusion of monensin and tylosin did not alter fecal shedding of experimentally-inoculated *E. coli* O157:H7 when either were fed alone or in combination in cattle fed a barley (grain)-based diet (McAllister *et al.*, 2006). These results are intriguing in that they suggest there is a potential interaction between diet type and antimicrobial treatment; however, no definitive proof of this linkage has been demonstrated.

Other feed additives have been examined such as ractopamine and neomycin. Researchers found that the β -agonist ractopamine, which is used to increase animal

growth performance, increased fecal shedding and cecal populations of *E. coli* O157:H7 in sheep (Edrington et al., 2006b). Neomycin is an antibiotic that is used to a limited degree in human medicine and is approved for use in cattle to treat enteric infections. Neomycin was demonstrated to reduce *E. coli* O157:H7 populations in the gut (Elder et al., 2002, Ransom et al., 2003) and on the hides of cattle (Ransom et al., 2003). However, this treatment has not been recommended to reduce *E. coli* O157:H7 in cattle due to antimicrobial resistance concerns.

Probiotics and E. coli O157:H7

In order to enhance ruminant production efficiency, various probiotics (including yeast cultures, competitive exclusion [CE] products, and direct-fed microbials [DFM]) have been widely used in the cattle industry for many years (Dawson et al., 1990, Yoon and Stern, 1996). These probiotic products have been primarily utilized to increase growth rate, milk production, or production efficiency (Callaway et al., 2005b, Fuller, 1989). A commercial *Saccharomyces cerevisiae* DFM culture was found to reduce *E. coli* O157:H7 populations in batch culture but not in a continuous flow culture system that simulated a bovine gut (Bach et al., 2003). A probiotic culture comprised of *Streptococcus bovis* and *Lactobacillus gallinarum* from the rumen of cattle reduced *E. coli* O157 shedding when given to experimentally-infected calves, and this decrease was attributed to an increase in VFA concentration in the gut (Ohya et al., 2001).

Other researchers have specifically developed probiotic products to reduce *E. coli* O157:H7 shedding in cattle. A probiotic that contained *S. faecium*, or a mixture of *S. faecium*, *L. acidophilus*, *L. casei*, *L. fermentum* and *L. plantarum* significantly reduced fecal shedding of *E. coli* O157:H7 in sheep from 2-4 log₁₀ CFU/g feces (Lema et al., 2001); however, a probiotic composed solely of *Lactobacillus acidophilus* was ineffective (Lema et al., 2001). A *L. acidophilus* culture that was derived from a cattle rumen reduced *E. coli* O157:H7 shedding by more than 50% when fed to feedlot cattle (Brashears and Galyean, 2002, Brashears et al., 2003a, Brashears et al., 2003b). This culture reduced fecal shedding of *E. coli* O157:H7 in cattle from 46% to 13% (Ransom et al., 2003). In a further refinement, *L. acidophilus* cultures were combined with *Propionibacterium freudenreichii* (a propionate-producing commensal bacteria) reduced the prevalence of *E. coli* O157:H7 in the feces from approximately 27% to 16% and reduced the prevalence on hides from 14% to 4% (Elam et al., 2003, Younts-Dahl et al., 2004). Further work has again shown reductions in *E. coli* O157:H7 and *Salmonella* in feces and on hides (Stephens et al., 2007). This probiotic improves the growth efficiency of cattle, which can economically pay for the cost of its inclusion in cattle rations as a food safety enhancement.

Competitive exclusion (CE) is another strategy to eliminate *E. coli* O157:H7 (as well as *Salmonella*) from cattle intestinal tracts (Brashears and Galyean, 2002, Brashears et al., 2003a, Brashears et al., 2003b, Zhao et al., 2003). A defined population of multiple non-EHEC *E. coli* strains that were isolated from cattle that did not contain *E. coli* O157:H7, and found this generic *E. coli* CE culture could

displace an established *E. coli* O157:H7 population from calves (Zhao et al., 1998).

Feedstuffs and *E. coli* O157:H7 populations?

Finishing beef and lactating dairy cattle in the United States are fed high grain rations in order to maximize animal performance and production efficiency (Huntington, 1997). Ruminant bacteria can break down dietary starch, although some starch does reach the colon in an undegraded form where it undergoes microbial fermentation (Huntington, 1997). Fecal samples from cattle fed dry rolled corn, high-moisture corn and wet corn gluten feed did not contain different populations of generic *E. coli*, or extreme acid-resistant *E. coli* during a limit-feeding period (Scott et al., 2000). However, feces from cattle fed wet corn gluten *ad libitum* contained significantly higher concentrations of extreme acid resistant *E. coli* (resistant to an acid shock simulating passage through the human stomach) than did feces of cattle fed dry-rolled or high moisture corn (Scott et al., 2000).

Barley is often fed in cattle rations, and is fermented more quickly than corn by the ruminal microbial population. This means that more starch is fermented in the lower gut of corn-fed cattle than in barley-fed cattle; resulting in barley-fed cattle having a higher fecal pH and lower VFA concentration compared with corn-fed animals (Bach et al., 2005a, Berg et al., 2004, Buchko et al., 2000a). Barley feeding was linked (albeit at a low correlation) to increased *E. coli* O157:H7 shedding (Dargatz et al., 1997); and in experimental infection studies barley feeding was again associated with increased shedding of *E. coli* O157:H7 by feedlot cattle (Buchko et al., 2000a). Feeding of barley-based diets resulted in higher fecal pH compared to corn diets and resulted in higher *E. coli* O157:H7 prevalence and quantity of *E. coli* O157:H7 shedding compared to cattle fed corn-based rations (Berg et al., 2004). Survival of *E. coli* O157:H7 in manure from corn- and barley fed cattle is approximately equal, therefore simple survival in the feces is not responsible for the increased prevalence of *E. coli* O157:H7 in barley-fed cattle (Bach et al., 2005b).

Recent research has demonstrated that steam-flaked grains increased *E. coli* O157 shedding in the feces compared to diets composed of dry-rolled grains (Fox et al., 2007). This difference was theorized to be due to dry rolling allowing the passage of more starch to the hindgut where it was fermented to produce VFA thereby killing *E. coli* O157 (Fox et al., 2007). This theory is supported by the fact that post-ruminal starch infusion, increased generic *E. coli* populations in the lower gut numerically (Van Kessel et al., 2002).

Comparing grain-fed to forage-fed cattle still indicates that more *E. coli* (including O157:H7) are present in the feces of cattle fed grain diets. In experimental inoculation studies the calves that consistently shed the highest concentrations of *E. coli* O157:H7 were fed a high concentrate (grain) diet (Tkalcic et al., 2000). Ruminant fluid collected from steers fed a high-forage diet allowed *E. coli* O157:H7 to proliferate to higher populations in vitro than did ruminant fluid from high-grain fed steers (Tkalcic et al., 2000), this was possibly due to differences in VFA concentrations between the ruminal fluids. Other researchers found that feeding forage actually

increased the shedding of *E. coli* O157:H7 in cattle (Van Baale et al., 2004). When cattle were fed forage *E. coli* O157:H7 was shed for 60 d compared to 16 for cattle on a grain-based diet (Van Baale et al., 2004). However, it must be emphasized, that although populations *E. coli* O157:H7 are generally lower in cattle fed forage diets, EHEC are still isolated from cattle solely fed forage, so forage feeding should not be viewed as a magic bullet (Hussein et al., 2003b, Thran et al., 2001).

One of the most interesting developments involving dietary effects on pathogens in cattle has evolved from the recent surge in the use of corn to produce ethanol. Distillers grains were shown to increase the shedding of *E. coli* O157:H7 in cow-calf operations in Scotland (Synge et al., 2003). Other researchers found that feeding a related product (brewers grain) to cattle was also associated with increased *E. coli* O157 shedding, and increased the odds of shedding by more than 6-fold (Dewell et al., 2005). More recent studies have shown that there is a positive association between distillers grain feeding and an increased prevalence of *E. coli* O157 (Jacob et al., 2008a, Jacob et al., 2008b). The individual animal prevalence of feedlot cattle shedding *E. coli* O157 on d 122 (but not d 136) was higher in cattle fed 25% wet distiller's grain (WDG) compared to control diets lacking WDG (Jacob et al., 2008b), but the pen-level shedding was unaffected by WDG feeding. In a follow-up study, cattle were fed a steam-flaked corn diet supplemented with 0% or 25% dried distiller's grain (DDG). Pen floor fecal sample prevalence of *E. coli* O157 was significantly higher across a 12 week finishing period in cattle fed 25% DDG and either 15% or 5% corn silage compared with cattle fed 0% DDG and 15% corn silage (Jacob et al., 2008a). Further studies found that DDG significantly increased fecal shedding and intestinal populations of inoculated *E. coli* O157:H7 compared to calves fed a steam-flaked corn based ration (Jacob et al., 2008c). It is important to note that the extent of the increase in *E. coli* O157:H7 is variable, and a great deal of variation occurs between sources and even batches of distillers grains. The underlying biology behind this effect has not been elucidated to this point, but it has been suggested that difference could be due to intermediate endproducts of the yeast fermentation (e.g., vitamins, organic acids), however these suggestions remain hypotheses. *In vitro* studies in our laboratory have detected no effects of DG on *E. coli* O157:H7 populations in mixed ruminal and fecal fluid fermentations (Callaway et al., unpublished). While the magnitude of these DG effects is relatively small and variable, it underlines the point that diet can potentially significantly impact *E. coli* O157:H7 populations in the gut of cattle.

Orange peel and pulp included at 2% of the total volume have been demonstrated to have anti-*E. coli* O157:H7 activity in *in vitro* fermentations (Callaway et al., 2008, Callaway et al., 2005a, Fisher and Phillips, 2006). This effect appears to be a result of the antimicrobial action of essential oils (e.g., limonene) found in the peel (Callaway et al., 2008). Other research has found that feeding an *Ascophyllum nodosum* supplement reduced *E. coli* O157:H7 populations in feces from 35% to 10% and reduced hide sample positives from 85% to less than 50% (Braden et al., 2004). These results underscore that certain feedstuffs and diet can exert

potent effects on the microbial population and can be used to control pathogens in certain circumstances and dietary regimens.

Forage to grain dietary shifts: their effects on fecal *E. coli* and *E. coli* O157:H7 populations

Escherichia coli can and does thrive in the lower gut of animals fed high grain diets, as well as those fed forage diets (Hussein et al., 2003a, Hussein et al., 2003b, Jacobson et al., 2002). However, shifting the forage to grain ratio in cattle rations can affect *E. coli* O157:H7 shedding. Early studies investigating *E. coli* and dietary effects indicated that a sudden decrease in hay intake by cattle increased fecal *E. coli* populations (Brownlie and Grau, 1967) and overfeeding of cattle with grain caused an increase in total fecal coliform counts (Allison et al., 1975). Other studies using experimentally infected sheep found a sudden switch from an alfalfa pellet diet to a low quality forage diet increased *E. coli* O157:H7 shedding (Kudva et al., 1995). Sheep shifted from a 50:50 corn/alfalfa ration to poor quality grass hay shed greater populations of *E. coli* O157:H7 than did sheep fed a corn/alfalfa ration (Kudva et al., 1997).

In the study that started a great deal of controversy on this topic, cattle fed a 90% corn/soybean meal ration (feedlot-type ration) contained generic *E. coli* populations that were 1000-fold higher than cattle fed a 100% good-quality hay (Timothy) diet (Diez-Gonzalez et al., 1998). The *E. coli* recovered from the feces of grain-fed cattle in this study were 1000-fold more resistant to an "extreme" acid shock that simulated passage through the human stomach than were *E. coli* from cattle fed only hay (Diez-Gonzalez et al., 1998). When cattle were abruptly switched from a 90% grain finishing ration to a 100% hay diet, fecal *E. coli* populations declined 1000-fold, and the population of *E. coli* resistant to an acid environment similar to that of the human stomach declined more than 100,000-fold within 5 d (Diez-Gonzalez et al., 1998). It is important to note that in this study no *E. coli* O157:H7 were detected. Based on these results the authors suggested that feedlot cattle be switched from high grain diets to hay for 5 days prior to slaughter to reduce *E. coli* contamination entering the abattoir (Diez-Gonzalez et al., 1998).

Although it appears that brief periods of hay feeding can affect *E. coli* populations research indicates that a brief period does not have a significant impact on carcass characteristics but does change final BW (Stanton and Schutz, 2000). When cattle were fed hay during the final portion of the finishing period, they had lower DMI and lost an average of 2.2 lb/head/d and did not significantly impact carcass weight, dressing percentage, carcass grades, or quality parameters, but did significantly reduce total coliform counts as well as generic *E. coli* counts (Stanton and Schutz, 2000), but the impact was not as great as that seen by Diez-Gonzalez et al. (1998). Keen et al., (1999) found that switching cattle from grain to hay caused a decrease in body weight (approximately 1.25 lb/hd/d compared to controls). Over 200 cattle maintained on a grain ration were screened for natural *E. coli* O157:H7 infection and 53% were found to be positive (Keen et al., 1999). When these cattle were divided into two groups and one was fed grain

and the other abruptly switched to hay, 52% of the grain-fed controls remained *E. coli* O157:H7 positive, but only 18% of the hay-fed cattle continued to shed *E. coli* O157:H7 (Keen et al., 1999).

The proposal of such a dietary switch to reduce *E. coli* O157:H7 shedding provoked a great deal of scientific controversy (Diez-Gonzalez et al., 1998, Hancock et al., 2000) and led to several studies that have subsequently evaluated the effect of radical dietary shifts on *E. coli* populations in cattle, however these studies have also produced conflicting results (Table 1). When cattle were fed a high-concentrate diet and switched to a diet containing 50% corn silage and 50% alfalfa hay, generic *E. coli* counts decreased (Jordan and McEwen, 1998). Cattle fed an 80% barley ration (which as shown previously tends to increase EHEC shedding) were fasted for 48 h and then subsequently switched to 100% alfalfa silage did not exhibit any change in *E. coli* O157:H7 shedding (Buchko et al., 2000b). However, when these same forage-fed animals were again fasted for 48 h and re-fed 100% alfalfa silage, the prevalence of *E. coli* O157:H7 shedding increased significantly (Buchko et al., 2000b). In a study using experimentally infected cattle, Researchers found that cattle fed hay shed *E. coli* O157:H7 significantly longer than did grain-fed cattle (42 d vs. 4 d), but *E. coli* O157:H7 populations shed were similar between dietary regimes and the diet shift did not affect the acid resistance of *E. coli* O157:H7 (Hovde et al., 1999). When cattle were abruptly switched from a finishing diet that contained wet corn gluten feed to alfalfa hay for 5 d, colonic pH increased almost 1 pH unit, total *E. coli* populations decreased approximately 10-fold (Scott et al., 2000). These authors concluded "increased colonic pH was not associated with reduced populations of acid resistant *E. coli*" but "feeding hay for a short duration can reduce acid-resistant *E. coli* populations" (Scott et al., 2000).

In research that approached this question from a different perspective, it was found that when cattle switched from forage-type diets (bromegrass hay or corn silage) to a high grain finishing ration, fecal and ruminal generic *E. coli* concentrations increased (Berry et al., 2006). However, in this study *E. coli* O157:H7 levels were not significantly impacted (Berry et al., 2006). In another study, switching cattle from pasture to hay for 48 h prior to slaughter significantly reduced the *E. coli* population throughout the gut, and found that hay feeding increased intestinal *Enterococci* populations that were capable of inhibiting *E. coli* populations (Gregory et al., 2000). Based on their data, the authors concluded, "the most effective way of manipulating gastro-intestinal counts of *E. coli* was to feed hay" (Gregory et al., 2000). Collectively, these results emphasize that dietary manipulations could be a powerful method to reduce *E. coli*/EHEC populations in cattle prior to harvest.

A theory addressing the effects of forage and dietary shifts on *E. coli* O157:H7 populations

It appears that a dietary shift does cause changes in the microbial populations, including *E. coli* O157:H7, yet studies investigating this phenomenon have often produced contradictory results (Table 1). Shifting the diet abruptly obviously causes a change in the availability and

concentrations of substrates available for fermentation in the lower intestinal tract, as well as the members of the microbial intestinal population. Grain-based diets tend to yield higher levels of *E. coli* in the feces due to the higher availability of starch. A change from grain diets to hay shifts the site and extent of digestion toward the rumen, and reduces starch availability in the colon. This change in nutrient availability furthermore causes a rapid shift in the microbial population of the gut, and some organisms selected for by hay feeding (such as *Enterococcus*) can exhibit competitive exclusion type behavior (Gregory et al., 2000), displacing established *E. coli* O157:H7 populations or preventing colonization (Callaway and Martin, 2006, Schneitz, 2005). Increasing the fiber component of the diet also increases the undigestible component of the diet which can physically "scrape" the gut mucosa which has been shown to be colonized by *E. coli* O157:H7 (Lim et al., 2007, Low et al., 2005, Naylor et al., 2003), and could physically remove these organisms.

Comparing results from dietary switch studies, it appears that some component of forage quality is involved in some of the differences (e.g. switching to alfalfa hay vs. switching to sagebrush). To date, however it has not been clearly demonstrated which aspect of forage quality is involved because none of these studies have used similar diets, protocols or *E. coli*/EHEC isolation techniques, thereby preventing direct comparison. We theorize that some factors intrinsic to forages may explain some of this effect, as well as some of the inconsistencies found between forages used.

Tannins are anti-nutritional polyphenols that are found in some forages, which have been shown to inhibit the growth of ruminal bacterial (Nelson et al., 1997, Nelson et al., 1998). Tannins inhibit and kill *E. coli* O157:H7 in *in vitro* studies and were found to reduce the shedding of generic *E. coli* over a 15 d period in steers fed tannins (Min et al., 2007). Additionally, as forages mature concentrations of the carbohydrate lignin, which contains carboxylic phenols such as *p*-coumaric acid and vanillin (Heppner, 1968, Martin, 1970). It has been long known that these components of lignin are bactericidal (Martin, 1990), but only recently has research demonstrated that these compounds are capable of killing *E. coli* O157:H7 *in vitro* and in manure (Wells et al., 2005). These data suggest a theory that forage quality and components may play a role in reducing *E. coli* O157:H7 in food animals, and that variation in the concentration of tannins and/or lignin in forages tested in various studies may be responsible for the variance in the dietary shift studies. In general, the greatest reductions in *E. coli* O157:H7 populations have been observed when cattle were switched to high quality forages; yet high concentrations of tannins and lignin are negatively correlated with forage quality. Therefore, details surrounding this hypothesis obviously need to be examined further. However, given the effects phenolic compounds have in altering the efficiency of the ruminal fermentation (Martin, 1990, Nelson et al., 1997), it is possible that, while tannins and carboxylic phenols do directly kill *E. coli* O157:H7 at physiologically unrealistic concentrations, when phenolics are added to in the intestinal consortium the phenolic compounds could still provide a competitive advantage to organisms that compete against

Table 1. Published reports of effects of diet on fecal <i>E. coli</i> populations in cattle					
Authors	Concentrate diet	<i>E. coli</i>	Forage diet or alternative feed	<i>E. coli</i> CFU	Log ₁₀ impact
Allison, 1975	Normal diet Over fed grain	8 x 10 ⁶ CFU coliforms/g 1 x 10 ¹⁰ CFU coliforms/g			
Kudva, et al., 1995	100% Alfalfa pellets	Shed O157:H7 for 4 d	Sagebrush/bunchgrass	Shed O157:H7 for 15 d	
Diez-Gonzalez et al., 1998	90% Concentrate	8 x 10 ⁷ CFU/g feces	100% Timothy Hay	3 x 10 ⁴ CFU/g feces	-3.5
Jordan and McEwen, 1998	44% Dry corn 7% Dry Gluten 7% Distiller's Dried Grains	7 x 10 ⁶ CFU/g feces	50% Corn silage 50% Alfalfa	4 x 10 ⁶ CFU/g feces	-0.3
Keen et al., 1999	85% Concentrate	52% shedding O157:H7	100% Forage	18% shedding O157:H7	
Hovde et al., 1999	62% Barley/19% corn 90% Corn	7 x 10 ⁶ CFU/g feces peak (4 d of shedding)	Alfalfa or Grass Hay	7 x 10 ⁶ CFU/g feces peak (39 or 42 d shedding)	
Stanton and Schutz, 2000	85% Whole Corn	3.2 x 10 ⁷ CFU/g	30% Millet hay 62% whole corn	1 x 10 ⁶ CFU/g feces	-1.2
Scott et al., 2000	84% DRC or 41% DRC 45% WCG	3 x 10 ⁸ CFU/g 5 x 10 ⁸ CFU/g feces	100% Alfalfa hay	1 x 10 ⁷ CFU/g 9 x 10 ⁶ CFU/g feces	-1.2 -1.6
Bucko et al., 2000a	80% Concentrate	5% shedding O157:H7	100% Alfalfa silage (after 48 h fast) Re-fed 100% alfalfa silage (after 48 h fast)	5% shedding O157:H7 42% shedding O157:H7 after 5 d	
Van Baale et al, 2004	80% corn 15% alfalfa hay	16 d of shedding O157:H7	70% prairie hay 15% alfalfa hay	60 d of shedding O157:H7	
Berry et al., 2006	70% corn and 24% corn silage	Approximately 10 ⁶ CFU/g generic <i>E. coli</i>	100% brome grass hay or 87% corn silage	Approximately 10 CFU/g generic <i>E. coli</i>	
Fox et al., 2007	Steam flaked corn/wheat and sorghum vs. dry-rolled corn/wheat and sorghum	65% shedding O157 Vs. 30% shedding			
Jacob et al., 2008a	85% steam-flaked corn 15% corn silage	3.6% shedding O157	25% of SFC replaced with Dried Distillers Grain	9.8% shedding O157	
Jacob et al., 2008b	84% steam flaked corn	14% shedding O157	25% of grain replaced with Wet Distillers Grain	3% shedding O157	

E. coli O157:H7, helping to reduce its populations in the intestinal ecosystem.

Conclusions

The United States has one of the safest food supplies, yet food-borne pathogenic bacteria are still significant threats to human health, including Enterohaemorrhagic *E. coli*. Post-harvest sanitation strategies have reduced *E. coli* O157:H7 in meat products, but pre-harvest intervention strategies offer methods to reduce pathogen populations in food animals before they enter the food chain. Reductions in *E. coli* O157:H7 shedding on farms can reduce human exposures through water supplies, fruits and vegetables, as well as via direct animal contact. Some feedstuffs do appear to alter shedding levels of *E. coli* O157:H7, but these effects have not always been consistent. Fasting and feeding poor quality forages have been shown to increase shedding of *E. coli* O157:H7 in cattle; however abruptly switching cattle from a high grain ration to a high-quality hay-based diet has been shown to reduce generic *E. coli* and *E. coli* O157:H7 populations. However, switching all feedlot cattle in the U.S. from grain-based diets to hay prior to slaughter is not practical. Further research is needed to elucidate the mechanism (e.g., competitive exclusion, physical removal, forage quality, tannins, lignin, other phenolics) by which forage-feeding impacts the microbial ecology of the bovine intestinal tract, including the ecology of *E. coli* and *E. coli* O157:H7 populations, so that economically viable and practical dietary modifications can be implemented.

References

- Acheson, D.W.K. 2000. How does *Escherichia coli* O157:H7 testing in meat compare with what we are seeing clinically? *J. Food Prot.* 63:819-821.
- Allison, M.J., I.M. Robinson, R.W. Dougherty and J.A. Bucklin. 1975. Grain overload in cattle and sheep: changes in microbial populations in the cecum and rumen. *Amer. J. Vet. Res.* 36:181-185.
- Anonymous. 2000. Waterborne outbreak of gastroenteritis associated with a contaminated municipal water supply, Walkerton, Ontario, May-June 2000. *Can. Commun. Dis. Rep.* 26:170-173.
- Arthur, T.M., G.A. Barkocy-Gallagher, M. Rivera-Betancourt and M. Koohmaraie. 2002. Prevalence and characterization of non-O157 shiga toxin-producing *Escherichia coli* on carcasses in commercial beef cattle processing plants. *Appl. Environ. Microbiol.* 68:4847-4852.
- Arthur, T.M., J.M. Bosilevac, D.M. Brichta-Harhay, M.N. Guerini, N. Kalchayanand, S.D. Shackelford, T.L. Wheeler and M. Koohmaraie. 2007a. Transportation and lairage environment effects on prevalence, numbers, and diversity of *Escherichia coli* O157:H7 on hides and carcasses of beef cattle at processing. *J. Food Prot.* 70:280-286.
- Arthur, T.M., J.M. Bosilevac, X. Nou, S.D. Shackelford, T.L. Wheeler and M. Koohmaraie. 2007b. Comparison of the molecular genotypes of *Escherichia coli* O157:H7 from the hides of beef cattle in different regions of North America. *J. Food Prot.* 70:1622-1626.
- Bach, S.J., T.A. McAllister, J. Baah, L.J. Yanke, D.M. Veira, V.P.J. Gannon and R.A. Holley. 2002a. Persistence of *Escherichia coli* O157:H7 in barley silage: Effect of a bacterial inoculant. *J. Appl. Microbiol.* 93:288-294.
- Bach, S.J., T.A. McAllister, D.M. Veira, V.P.J. Gannon and R.A. Holley. 2002b. Effect of monensin on survival and growth of *Escherichia coli* O157:H7 in vitro. *Can. Vet. J.* 43:718-719.
- Bach, S.J., T.A. McAllister, D.M. Veira, V.P.J. Gannon and R.A. Holley. 2003. Effects of a *Saccharomyces cerevisiae* feed supplement on *Escherichia coli* O157:H7 in ruminal fluid in vitro. *Anim. Feed Sci. Technol.* 104:179-189.
- Bach, S.J., L.J. Selinger, K. Stanford and T. McAllister. 2005a. Effect of supplementing corn- or barley-based feedlot diets with canola oil on faecal shedding of *Escherichia coli* O157:H7 by steers. *J. Appl. Microbiol.* 98:464-475.
- Bach, S.J., K. Stanford and T.A. McAllister. 2005b. Survival of *Escherichia coli* O157:H7 in feces from corn- and barley-fed steers. *FEMS Microbiol. Lett.* 252:25-33.
- Barkocy-Gallagher, G.A., T.M. Arthur, M. Rivera-Betancourt, X. Nou, S.D. Shackelford, T.L. Wheeler and M. Koohmaraie. 2003. Seasonal prevalence of shiga toxin-producing *Escherichia coli*, including O157:H7 and non-O157 serotypes, and *Salmonella* in commercial beef processing plants. *J. Food Prot.* 66:1978-1986.
- Barkocy-Gallagher, G.A., T.M. Arthur, M. Rivera-Betancourt, X. Nou, S.D. Shackelford, T.L. Wheeler and M. Koohmaraie. 2004. Characterization of O157:H7 and other *Escherichia coli* isolates recovered from cattle hides, feces, and carcasses. *J. Food Prot.* 67:993-998.
- Berg, J.L., T.A. McAllister, S.J. Bach, R.P. Stillborn, D.D. Hancock and J.T. LeJeune. 2004. *Escherichia coli* O157:H7 excretion by commercial feedlot cattle fed either barley- or corn-based finishing diets. *J. Food Prot.* 67:666-671.
- Berry, E.D., J.E. Wells, S.L. Archibeque, C.L. Ferrell, H.C. Freetly and D.N. Miller. 2006. Influence of genotype and diet on steer performance, manure odor, and carriage of pathogenic and other fecal bacteria. 2. Pathogenic and other fecal bacteria. *J. Anim. Sci.* 84:2523-2532.
- Berry, E.D., B.L. Woodbury, J.A. Nienaber, R.A. Eigenberg, J.A. Thurston and J.E. Wells. 2007. Incidence and persistence of zoonotic bacterial and protozoan pathogens in a beef cattle feedlot runoff control-vegetative treatment system. *J. Environ. Qual.* 36:1873-1882.
- Bosilevac, J.M., T.L. Wheeler, M. Rivera-Betancourt, X. Nou, T.M. Arthur, S.D. Shackelford, M.P. Kent, D. Jaroni, M.S. Osborn, M.L. Rossman, J.O. Reagan and M. Koohmaraie. 2004. Protocol for evaluating the efficacy of cetylpyridinium chloride as a beef hide intervention. *J. Food Prot.* 67:303-309.
- Bosilevac, J.M., S.D. Shackelford, D.M. Brichta and M. Koohmaraie. 2005. Efficacy of ozonated and electrolyzed oxidative waters to decontaminate hides of cattle before slaughter. *J. Food Prot.* 68:1393-1398.
- Bosilevac, J.M., X. Nou, G.A. Barkocy-Gallagher, T.M. Arthur and M. Koohmaraie. 2006. Treatments using hot water instead of lactic acid reduce levels of aerobic bacteria and Enterobacteriaceae and reduce the prevalence of *Escherichia coli* O157:H7 on previsceration beef carcasses. *J. Food Prot.* 69:1808-1813.
- Braden, K.W., J.R. Blanton Jr, V.G. Allen, K.R. Pond and M.F. Miller. 2004. *Ascophyllum nodosum* supplementation: A preharvest intervention for reducing *Escherichia coli* O157:H7 and *Salmonella* spp. in feedlot steers. *J. Food Prot.* 67:1824-1828.
- Brashears, M.M. and M.L. Galyean. 2002. Testing of

- probiotic bacteria for the elimination of *Escherichia coli* O157:H7 in cattle. Amer. Meat Inst. Found. Last Accessed: Access Date: 24 April 2007, 2002. Available at: <http://www.amif.org/PRProbiotics042302.htm>
- Brashears, M.M., M.L. Galyean, G.H. Loneragan, J.E. Mann and K. Killinger-Mann. 2003a. Prevalence of *Escherichia coli* O157:H7 and performance by beef feedlot cattle given *Lactobacillus* direct-fed microbials. J. Food Prot. 66:748-754.
- Brashears, M.M., D. Jaroni and J. Trimble. 2003b. Isolation, selection, and characterization of lactic acid bacteria for a competitive exclusion product to reduce shedding of *Escherichia coli* O157:H7 in cattle. J. Food Prot. 66:355-363.
- Brownlie, L.E. and F.H. Grau. 1967. Effect of food intake on growth and survival of *Salmonellas* and *Escherichia coli* in the bovine rumen. J. Gen. Microbiol. 46:125-134.
- Buchko, S.J., R.A. Holley, W.O. Olson, V.P.J. Gannon and D.M. Veira. 2000a. The effect of different grain diets on fecal shedding of *Escherichia coli* O157:H7 by steers. J. Food Prot. 63:1467-1474.
- Buchko, S.J., R.A. Holley, W.O. Olson, V.P.J. Gannon and D.M. Veira. 2000b. The effect of fasting and diet on fecal shedding of *Escherichia coli* O157:H7 by cattle. Can. J. Anim. Sci. 80:741-744.
- Callaway, T.R., T.S. Edrington, J.L. Rychlik, K.J. Genovese, T.L. Poole, Y.S. Jung, K.M. Bischoff, R.C. Anderson and D.J. Nisbet. 2003. Ionophores: Their use as ruminant growth promotants and impact on food safety. Curr. Iss. Intest. Microbiol. 4:43-51.
- Callaway, T.R., R.C. Anderson, T.S. Edrington, K.J. Genovese, R.B. Harvey, T.L. Poole and D.J. Nisbet. 2004. Recent pre-harvest supplementation strategies to reduce carriage and shedding of zoonotic enteric bacterial pathogens in food animals. Anim. Health Res. Rev. 5:35-47.
- Callaway, T.R., J.B. Carroll, J.D. Arthington, R.C. Anderson, T.S. Edrington, K.J. Genovese and D.J. Nisbet. 2005a. Orange pulp reduces growth of *E. coli* O157:H7 and *Salmonella Typhimurium* in pure culture and in vitro mixed ruminal microorganism fermentation. Procs. Amer. Soc. Anim. Sci./Amer. Dairy Sci. Assoc./Can. Anim. Sci. Assoc Annual Meeting. Cincinnati, OH. 236.
- Callaway, T.R., K.D. Dunkley, R.C. Anderson, T.S. Edrington, K.J. Genovese, T.L. Poole, R.B. Harvey and D.J. Nisbet. 2005b. Probiotics, vaccines and other intervention strategies. In: Raw Material Safety: Meat. Sofos, J.N. (eds), Woodhead Pub., Cambridge, UK pp. 192-213.
- Callaway, T.R., T.S. Edrington, A.D. Brabban, J.E. Keen, R.C. Anderson, M.L. Rossman, M.J. Engler, K.J. Genovese, B.L. Gwartney, J.O. Reagan, T.L. Poole, R.B. Harvey, E.M. Kutter and D.J. Nisbet. 2006. Fecal prevalence of *Escherichia coli* O157, *Salmonella*, *Listeria*, and bacteriophage infecting *E. coli* O157:H7 in feedlot cattle in the southern plains region of the United States. Foodborne Pathog. Dis. 3:234-244.
- Callaway, T.R. and S.A. Martin. 2006. Use of competitive exclusion cultures and oligosaccharides. In: Feedstuffs Direct-fed Microbial, Enzyme and Forage Additive Compendium, 8th Ed. (eds), Miller Publishing, Minnetonka, MN pp. 34-39.
- Callaway, T.R., J.A. Carroll, J.D. Arthington, C. Pratt, T.S. Edrington, R.C. Anderson, M.L. Galyean, S.C. Ricke, P. Crandall and D.J. Nisbet. 2008. Citrus products decrease growth of *E. coli* O157:H7 and *Salmonella* Typhimurium in pure culture and in fermentation with mixed ruminal microorganism in vitro. Foodborne Path. Dis. (Accepted 18 April 2008):
- Castillo, A., J.S. Dickson, R.P. Clayton, L.M. Lucia and G.R. Acuff. 1998. Chemical dehairing of bovine skin to reduce pathogenic bacteria and bacteria of fecal origin. J. Food Prot. 61:623-625.
- Chapman, P.A., J. Cornell and C. Green. 2000. Infection with verocytotoxin-producing *Escherichia coli* O157 during a visit to an inner city open farm. Epidemiol. Infect. 125:531-536.
- Cobbold, R.N. and P.M. Desmarchelier. 2004. In vitro studies on the colonization of bovine colonic mucosa by Shiga-toxigenic *Escherichia coli* (STEC). Epidemiol. Infect. 132:87-94.
- Cobbold, R.N. 2007. Rectoanal Junction Colonization of feedlot cattle by *Escherichia coli* O157:H7 and its association with supershedders and excretion dynamics. Appl. Environ. Microbiol. 73, no.5:1563-1568.
- Cookson, A.L., J. Bennett, F. Thomson-Carter and G.T. Attwood. 2007. Intimin subtyping of *Escherichia coli*: Concomitant carriage of multiple intimin subtypes from forage-fed cattle and sheep. FEMS Microbiol. Lett. 272:163-171.
- Cray, W.C., T.A. Casey, B.T. Bosworth and M.A. Rasmussen. 1998. Effect of dietary stress on fecal shedding of *Escherichia coli* O157:H7 in calves. Appl. Environ. Microbiol. 64:1975-1979.
- Dargatz, D.A., S.J. Wells, L.A. Thomas, D.D. Hancock and L.P. Garber. 1997. Factors associated with the presence of *Escherichia coli* O157 in feces of feedlot cattle. J. Food Prot. 60:466-470.
- Davidson, C.M. and M. Taylor. 1978. Variability of *E. coli* levels in bovine feces and its implications on guidelines for ground beef. Can. Inst. Food Sci. Technol. J. 11:53.
- Davis, M.A. 2006. Comparison of cultures from rectoanal-junction mucosal swabs and feces for detection of *Escherichia coli* O157 in dairy heifers. Appl. Environ. Microbiol. 72:3766-3770.
- Dawson, K.A., K.E. Newman and J.A. Boling. 1990. Effects of microbial supplements containing yeast and lactobacilli on roughage-fed ruminal microbial activities. J Anim. Sci. 3392-3398.
- Dewell, G.A., J.R. Ransom, R.D. Dewell, K. McCurdy, I.A. Gardner, A.E. Hill, J.N. Sofos, K.E. Belk, G.C. Smith and M.D. Salman. 2005. Prevalence of and risk factors for *Escherichia coli* O157 in market-ready beef cattle from 12 U.S. feedlots. Foodborne Path. Dis. 2:70-76.
- Diez-Gonzalez, F., T.R. Callaway, M.G. Kizoulis and J.B. Russell. 1998. Grain feeding and the dissemination of acid-resistant *Escherichia coli* from cattle. Science. 281:1666-1668.
- Drasar, B.S. 1974. Some factors associated with geographical variations in the intestinal microflora. In: The normal microbial flora of man. (eds), Academic Press, London pp. 187-196.
- Drasar, B.S. and P.A. Barrow. 1985. Intestinal Microbiology. In: (eds), Amer. Soc. Microbiol. Press, Washington, DC pp. 19-40.
- Edrington, T.S., T.R. Callaway, P.D. Varey, Y.S. Jung, K.M. Bischoff, R.O. Elder, R.C. Anderson, E. Kutter, A.D.

- Brabban and D.J. Nisbet. 2003. Effects of the antibiotic ionophores monensin, lasalocid, laidlomycin propionate and bambarmycin on *Salmonella* and *E. coli* O157:H7 in vitro. *J. Appl. Microbiol.* 94:207-213.
- Erdington, T.S., T.R. Callaway, S.E. Ives, M.J. Engler, M.L. Loofer, R.C. Anderson and D.J. Nisbet. 2006a. Seasonal shedding of *Escherichia coli* O157:H7 in ruminants: a new hypothesis. *Foodborne Pathog. Dis.* 3:413-421.
- Erdington, T.S., T.R. Callaway, S.E. Ives, M.J. Engler, T.H. Welsh, D.M. Hallford, K.J. Genovese, R.C. Anderson and D.J. Nisbet. 2006b. Effect of Ractopamine HCl Supplementation on Fecal Shedding of *Escherichia coli* O157:H7 and *Salmonella* in Feedlot Cattle. *Curr. Microbiol.* 53:340-345.
- Erdington, T.S., T.R. Callaway, D.M. Hallford, R.C. Anderson and D.J. Nisbet. 2007. Influence of exogenous triiodothyronine (T₃) on fecal shedding of *Escherichia coli* O157 in cattle. *Microbial Ecology.* 53:664-669.
- Edwards, A.C., D. Kay, A.T. McDonald, C. Francis, J. Watkins, J.R. Wilkinson and M.D. Wyer. 2008. Farmyards, an overlooked source for highly contaminated runoff. *J. Environ. Man.* 86:In Press.
- Elam, N.A., J.F. Gleghorn, J.D. Rivera, M.L. Galyean, P.J. Defoor, M.M. Brashears and S.M. Younts-Dahl. 2003. Effects of live cultures of *Lactobacillus acidophilus* (strains NP45 and NP51) and *Propionibacterium freudenreichii* on performance, carcass, and intestinal characteristics, and *Escherichia coli* strain O157 shedding of finishing beef steers. *J. Anim. Sci.* 81:2686-2698.
- Elder, R.O., J.E. Keen, G.R. Siragusa, G.A. Barkocy-Gallagher, M. Koohmaraie and W.W. Lagreid. 2000. Correlation of enterohemorrhagic *Escherichia coli* O157 prevalence in feces, hides, and carcasses of beef cattle during processing. *Proc. Nat. Acad. Sci. (USA).* 97:2999-3003.
- Elder, R.O., J.E. Keen, T.E. Wittum, T.R. Callaway, T.S. Erdington, R.C. Anderson and D.J. Nisbet. 2002. Intervention to reduce fecal shedding of enterohemorrhagic *Escherichia coli* O157:H7 in naturally infected cattle using neomycin sulfate. *J. Anim. Sci.* 80 (Suppl. 1):15 (Abstr.).
- Fagan, P.K., M.A. Hornitzky, K.A. Bettelheim and S.P. Djordjevic. 1999. Detection of shiga-like toxin (stx1 and stx2), intimin (eaeA), and enterohemorrhagic *Escherichia coli* (EHEC) hemolysin (EHEC hlyA) genes in animal feces by multiplex PCR. *Appl. Environ. Microbiol.* 65:868-872.
- Ferguson, C.M., C.M. Davies, C. Kaucner, M. Krogh, J. Rodehutsors, D.A. Deere and N.J. Ashbolt. 2007. Field scale quantification of microbial transport from bovine faeces under simulated rainfall events. *J. Water Health.* 5:83-95.
- Fisher, K. and C.A. Phillips. 2006. The effect of lemon, orange and bergamot essential oils and their components on the survival of *Campylobacter jejuni*, *Escherichia coli* O157, *Listeria monocytogenes*, *Bacillus cereus* and *Staphylococcus aureus* in vitro and in food systems. *J. Appl. Microbiol.* 101:1232-1240.
- Fox, J.T., B.E. Dejenbusch, J.S. Drouillard and T.G. Nagaraja. 2007. Dry-rolled or steam-flaked grain-based diets and fecal shedding of *Escherichia coli* O157 in feedlot cattle. *J. Anim. Sci.* 85:1207-1212.
- Fuller, R. 1989. Probiotics in man and animals. *J. Appl. Bacteriol.* 66:365-378.
- Gage, R. 2001. Outbreaks of *Escherichia coli* O157:H7 infections among children associated with farm visits-- Pennsylvania and Washington, 2000. *Morbid. Mortal. Weekly Rep.* 50:293-297.
- Gerba, C.P. and J.E. Smith. 2005. Sources of pathogenic microorganisms and their fate during land application of wastes. *J. Environ. Qual.* 34:42-48.
- Goss, M. and C. Richards. 2007. Development of a risk-based index for source water protection planning, which supports the reduction of pathogens from agricultural activity entering water resources. *J. Environ. Man.* 86:In Press.
- Grau, F.H., L.E. Brownlie and M.G. Smith. 1969. Effects of food intake on numbers of *Salmonellae* and *Escherichia coli* in rumen and faeces of sheep. *J. Appl. Bact.* 32:112-117.
- Grauke, L.J., I.T. Kudva, J.W. Yoon, C.W. Hunt, C.J. Williams and C.J. Hovde. 2002. Gastrointestinal tract location of *Escherichia coli* O157:H7 in ruminants. *Appl. Environ. Microbiol.* 68:2269-2277.
- Greenquist, M.A., J.S. Drouillard, J.M. Sargeant, B.E. Dejenbusch, X. Shi, K.F. Lechtenberg and T.G. Nagaraja. 2005. Comparison of rectoanal mucosal swab cultures and fecal cultures for determining prevalence of *Escherichia coli* O157:H7 in feedlot cattle. *Appl. Environ. Microbiol.* 71:6431-6433.
- Gregory, N.G., L.H. Jacobson, T.A. Nagle, R.W. Muirhead and G.J. Leroux. 2000. Effect of preslaughter feeding system on weight loss, gut bacteria, and the physico-chemical properties of digesta in cattle. *New Zealand J. Agric. Res.* 43:351-361.
- Gyles, C.L. 2007. Shiga toxin-producing *Escherichia coli*: An overview. *J. Anim. Sci.* 85:E45-62.
- Hancock, D.D., T.E. Besser, C. Gill and C.H. Bohach. 2000. Cattle, hay, and *E. coli*. *Science.* 284:49-50.
- Harmon, B.G., C.A. Brown, S. Tkalcic, P.O.E. Mueller, A. Parks, A.V. Jain, T. Zhao and M.P. Doyle. 1999. Fecal shedding and rumen growth of *Escherichia coli* O157:H7 in fasted calves. *J. Food Prot.* 62:574-579.
- Hepner, M.B. 1968. The tricky chemistry of a blade of grass. In: *Yearbook of agriculture.* (eds), pp. 72-76.
- Herriott, D.E., D.D. Hancock, E.D. Ebel, L.V. Carpenter, D.H. Rice and T.E. Besser. 1998. Association of herd management factors with colonization of dairy cattle by shiga toxin-positive *Escherichia coli* O157. *J. Food Prot.* 61:802-807.
- Hill, D.D., W.E. Owens and P.B. Tchounwou. 2006. Prevalence of *Escherichia coli* O157:H7 bacterial infections associated with the use of animal wastes in Louisiana for the period 1996-2004. *Int. J. Environ. Res. Pub. Health.* 3:107-113.
- Hollowell, C.A. and M.J. Wolin. 1965. Basis for the exclusion of *Escherichia coli* from the rumen ecosystem. *Appl. Microbiol.* 13:918-924.
- Hovde, C.J., P.R. Austin, K.A. Cloud, C.J. Williams and C.W. Hunt. 1999. Effect of cattle diet on *Escherichia coli* O157:H7 acid resistance. *Appl. Environ. Microbiol.* 65:3233-3235.
- Huntington, G.B. 1997. Starch utilization by ruminants: from basics to the bunk. *J. Anim. Sci.* 75:852-867.
- Hussein, H.S., B.H. Thran and H.A. Glimp. 2003a. Verotoxin-producing *Escherichia coli* in sheep grazing an irrigated pasture or arid rangeland forages. *Exp. Biol.*

- Med. 228:358-364.
- Hussein, H.S., B.H. Thrane, M.R. Hall, W.G. Kvasnicka and R.C. Torell. 2003b. Verotoxin-producing *Escherichia coli* in culled beef cows grazing rangeland forages. *Exp. Biol. Med.* 228:352-357.
- Hynes, N.A. and I.K. Wachsmuth. 2000. *Escherichia coli* O157:H7 risk assessment in ground beef: A public health tool. *Procs. Proc. 4th Int. Symp. on Shiga Toxin-Producing Escherichia coli Infections*. Kyoto, Japan. 46.
- Jacob, M.E., J.T. Fox, J.S. Drouillard, D.G. Renter and T.G. Nagaraja. 2008a. Effects of dried distillers' grain on fecal prevalence and growth of *Escherichia coli* O157 in batch culture fermentations from cattle. *Appl. Environ. Microbiol.* 74:38-43.
- Jacob, M.E., J.T. Fox, S.K. Narayanan, J.S. Drouillard, D.G. Renter and T.G. Nagaraja. 2008b. Effects of feeding wet corn distiller's grains with solubles with or without monensin and tylosin on the prevalence and antimicrobial susceptibilities of fecal food-borne pathogenic and commensal bacteria in feedlot cattle. *J. Anim. Sci.* 86:1182-1190.
- Jacob, M.E., G.L. Parsons, M.K. Shelor, J.T. Fox, J.S. Drouillard, D.U. Thomson, D.G. Renter and T.G. Nagaraja. 2008c. Feeding supplemental dried distiller's grains increases faecal shedding of *Escherichia coli* O157 in experimentally inoculated calves. *Zoon. Pub. Health.* 55:125-132.
- Jacobson, L.H., T.A. Nagle, N.G. Gregory, R.G. Bell, G. Le Roux and J.M. Haines. 2002. Effect of feeding pasture-finished cattle different conserved forages on *Escherichia coli* in the rumen and faeces. *Meat Sci.* 62:93-106.
- Jay, M.T., M. Cooley, D. Carychao, G.W. Wiscumb, R.A. Sweitzer, L. Crawford-Miksza, J.A. Farrar, D.K. Lau, J. O'Connell, A. Millington, R.V. Asmundson, E.R. Atwill and R.E. Mandrell. 2007. *Escherichia coli* O157:H7 in feral swine near spinach fields and cattle, central California coast. *Emerg. Infect. Dis.* Available from <http://www.cdc.gov/EID/content/13/12/1908.htm>.
- Jordan, D. and S.A. McEwen. 1998. Effect of duration of fasting and a short-term high-roughage ration on the concentration of *Escherichia coli* biotype 1 in cattle feces. *J. Food Prot.* 61:531-534.
- Keen, J.E., G.A. Uhlich and R.O. Elder. 1999. Effects of hay- and grain-based diets on fecal shedding in naturally-acquired enterohemorrhagic *E. coli* (EHEC) O157 in beef feedlot cattle. *Procs. 80th Conf. Research Workers in Animal Diseases*. Chicago, IL. (Abstr.).
- Keen, J.E. and R.O. Elder. 2002. Isolation of shiga-toxicogenic *Escherichia coli* O157 from hide surfaces and the oral cavity of finished beef feedlot cattle. *J. Amer. Vet. Med. Assoc.* 220:756-763.
- Keen, J.E., T.E. Wittum, J.R. Dunn, J.L. Bono and M.E. Fontenot. 2003. Occurrence of STEC O157, O111, and O26 in livestock at agricultural fairs in the United States. *Procs. Proc. 5th Int. Symp. on Shiga Toxin-Producing Escherichia coli Infections*. Edinburgh, UK. 22 (Abstr.).
- Keen, J.E., L.M. Durso and T.P. Meehan. 2007. Isolation of *Salmonella enterica* and shiga-toxicogenic *Escherichia coli* O157 from feces of animals in public contact areas of United States zoological parks. *Appl. Environ. Microbiol.* 73:362-365.
- Kudva, I.T., P.G. Hatfield and C.J. Hovde. 1995. Effect of diet on the shedding of *Escherichia coli* O157:H7 in a sheep model. *Appl. Environ. Microbiol.* 61:1363-1370.
- Kudva, I.T., P.G. Hatfield and C.J. Hovde. 1997. Characterization of *Escherichia coli* O157:H7 and other shiga toxin-producing *E. coli* serotypes isolated from sheep. *J. Clin. Microbiol.* 35:892-899.
- Laven, R.A., A. Ashmore and C.S. Stewart. 2003. *Escherichia coli* in the rumen and colon of slaughter cattle, with particular reference to *E. coli* O157. *Vet. J.* 165:78-83.
- LeJeune, J. and M. Kauffman. 2006. Bovine *E. coli* O157 supershedders: mathematical myth or meaningful monsters? *Procs. Proceedings of the 2006 VTEC Conference*. Melbourne, Australia. 26.
- LeJeune, J.T., T.E. Besser and D.D. Hancock. 2001. Cattle water troughs as reservoirs of *Escherichia coli* O157. *Appl. Environ. Microbiol.* 67:3053-3057.
- Lema, M., L. Williams and D.R. Rao. 2001. Reduction of fecal shedding of enterohemorrhagic *Escherichia coli* O157:H7 in lambs by feeding microbial feed supplement. *Small Rum. Res.* 39:31-39.
- Lim, J.Y., J. Li, H. Sheng, T.E. Besser, K. Potter and C.J. Hovde. 2007. *Escherichia coli* O157:H7 colonization at the rectoanal junction of long-duration culture-positive cattle. *Appl. Environ. Microbiol.* 73:1380-1382.
- Loneragan, G.H. and M.M. Brashears. 2005. Pre-harvest interventions to reduce carriage of *E. coli* O157 by harvest-ready feedlot cattle. *Meat Science.* 71:72.
- Low, J.C., I.J. McKendrick, C. McKechnie, D.R. Fenlon, S.W. Naylor, C. Currie, D.G.E. Smith, L. Allison and D.L. Gally. 2005. Rectal carriage of enterohemorrhagic *Escherichia coli* O157 in slaughtered cattle. *Appl. Environ. Microbiol.* 71:93-97.
- Manshadi, F.D., P. Gortares, C.P. Gerba, M. Karpiscak and R.J. Frentas. 2001. Role of irrigation water in contamination of domestic fresh vegetables. *Procs. Gen. Mtg. Amer. Soc. Microbiol.* 561.
- Martin, A.K. 1970. Effect of Stage of Maturity of Perennial Ryegrass on its Content of some Organic Acids and Phenolic Compounds. *J. Sci. Food Agric.* 21:496-501.
- Martin, S.A. 1990. Effect of phenolic compounds on fiber-degrading enzymes from rumen bacteria. In: Akin, D.E. (eds), *Elevier science publishing co, inc., Athens* pp. 289-300.
- McAllister, T.A., S.J. Bach, K. Stanford and T.R. Callaway. 2006. Shedding of *Escherichia coli* O157:H7 by cattle fed diets containing monensin or tylosin. *J. Food Prot.* 69:2075-2083.
- Mead, P.S., L. Slutsker, V. Dietz, L.F. McCraig, J.S. Bresee, C. Shapiro, P.M. Griffin and R.V. Tauxe. 1999. Food-related illness and death in the United States. *Emerg. Infect. Dis.* 5:607-625.
- Meyer-Brosseta, S., S.N. Bastian, P.D. Arne, O. Cerf and M. Sanaa. 2001. Review of epidemiological surveys on the prevalence of contamination of healthy cattle with *Escherichia coli* serogroup O157:H7. *Int. J. Hyg. Environ. Health.* 203:347-361.
- Midgley, J., N. Fegan and P. Desmarchelier. 1999. Dynamics of shiga toxin-producing *Escherichia coli* (STEC) in feedlot cattle. *Lett. Appl. Microbiol.* 29:85-89.
- Min, B.R., W.E. Pinchak, R.C. Anderson and T.R. Callaway. 2007. Effect of tannins on the in vitro growth of *Escherichia coli* O157:H7 and in vivo growth of generic *Escherichia coli* excreted from steers. *J. Food Prot.* 70:543-550.

- Natvig, E.E., S.C. Ingham, B.H. Ingham, L.R. Cooperband and T.R. Roper. 2002. *Salmonella enterica* serovar typhimurium and *Escherichia coli* contamination of root and leaf vegetables grown in soils with incorporated bovine manure. *Appl. Environ. Microbiol.* 68:2737-2744.
- Naumova, E.N., J.S. Jagai, B. Matyas, A. DeMaria Jr, I.B. MacNeill and J.K. Griffiths. 2007. Seasonality in six enterically transmitted diseases and ambient temperature. *Epidemiol. infect.* 135:281-292.
- Naylor, S.W., J.C. Low, T.E. Besser, A. Mahajan, G.J. Gunn, M.C. Pearce, I.J. McKendrick, D.G.E. Smith and D.L. Gally. 2003. Lymphoid follicle-dense mucosa at the terminal rectum is the principal site of colonization of enterohaemorrhagic *Escherichia coli* O157:H7 in the bovine host. *Infect. Immun.* 71:1505-1512.
- Nelson, K.E., A.N. Pell, P.H. Doane, B.I. Giner-Chavez and P. Schofield. 1997. Chemical and biological assays to evaluate bacterial inhibition by Tannins. *J. Chem. Ecol.* 23:1175-1194.
- Nelson, K.E., M.L. Thonney, T.K. Woolston, S.H. Zinder and A.N. Pell. 1998. Phenotypic and phylogenetic characterization of ruminant tannin-tolerant bacteria. *Appl. Environ. Microbiol.* 64:3824-3830.
- Nou, X., M. Rivera-Betancourt, J.M. Bosilevac, T.L. Wheeler, S.D. Shackelford, B.L. Gwartney, J.O. Reagan and M. Koohmaraie. 2003. Effect of chemical dehairing on the prevalence of *Escherichia coli* O157:H7 and the levels of aerobic bacteria and *Enterobacteriaceae* on carcasses in a commercial beef processing plant. *J. Food Prot.* 66:2005-2009.
- Ohya, T., T. Marubashi and H. Ito. 2000. Significance of fecal volatile fatty acids in shedding of *Escherichia coli* O157 from calves: experimental infection and preliminary use of a probiotic product. *J. Vet. Med. Sci.* 62:1151-1155.
- Ohya, T., M. Akiba and H. Ito. 2001. Use of a trial probiotic product in calves experimentally infected with *Escherichia coli* O157. *Japan Agric. Res. Quart.* 35:189-194.
- Pritchard, G.C., G.A. Willshaw, J.R. Bailey, T. Carson and T. Cheasty. 2000. Verocytotoxin-producing *Escherichia coli* O157 on a farm open to the public: outbreak investigation and longitudinal bacteriological study. *Vet. Rec.* 147:259-264.
- Prohaszka, L. and F. Baron. 1983. Antibacterial effect of volatile fatty acids on *Enterobacteriaceae* in the large intestine. *Acta. Vet. Hung.* 30:9-16.
- Ransom, J.R., K.E. Belk, J.N. Sofos, J.A. Scanga, M.L. Rossman, G.C. Smith and J.D. Tatum. 2003. Investigation of on-farm management practices as pre-harvest beef microbiological interventions.
- Rasmussen, M.A., T.L. Wickman, W.C. Cray and T.A. Casey. 1999. *Escherichia coli* O157:H7 and the rumen environment. In: *E. coli* O157 in Farm Animals. (eds), CAB International, pp. 39.
- Reid, C.A., A. Small, S.M. Avery and S. Buncic. 2002. Presence of food-borne pathogens on cattle hides. *Food Control.* 13:411-415.
- Reinstein, S., J.T. Fox, X. Shi and T.G. Nagaraja. 2007. Prevalence of *Escherichia coli* O157:H7 in gallbladders of beef cattle. *Applied and Environmental Microbiology.* 73:1002-1004.
- Rice, D.H., H.Q. Sheng, S.A. Wynia and C.J. Hovde. 2003. Rectoanal Mucosal Swab Culture Is More Sensitive Than Fecal Culture and Distinguishes *Escherichia coli* O157:H7- Colonized Cattle and Those Transiently Shedding the Same Organism. *J. Clin. Microbiol.* 41:4924-4929.
- Russell, J.B. and F. Diez-Gonzalez. 1998. The effects of fermentation acids on bacterial growth. *Adv. Microb. Physiol.* 39:205-234.
- Sargeant, J.M., M.W. Sanderson, R.A. Smith and D.D. Griffin. 2003. *Escherichia coli* O157 in feedlot cattle feces and water in four major feeder-cattle states in the USA. *Prev. Vet. Med.* 61:127-135.
- Sargeant, J.M., M.R. Amezcua, A. Rajic and L. Waddell. 2007. Pre-harvest interventions to reduce the shedding of *E. coli* O157 in the faeces of weaned domestic ruminants: a systematic review. *Zoonos. Pub. Health.* 54:260-277.
- Schneitz, C. 2005. Competitive exclusion in poultry--30 years of research. *Food Control.* 16:657-667.
- Schultz, C.L., T.S. Edrington, S.B. Schroeder, D.M. Hallford, K.J. Genovese, T.R. Callaway, R.C. Anderson and D.J. Nisbet. 2005. Effect of the thyroid on faecal shedding of *E. coli* O157:H7 and *Escherichia coli* in naturally infected yearling beef cattle. *Journal of Applied Microbiology.* 99:1176-1180.
- Schurman, R.D., H. Hariharan, S.B. Heaney and K. Rahn. 2000. Prevalence and characteristics of shiga toxin-producing *Escherichia coli* in beef cattle slaughtered on Prince Edward Island. *J. Food Prot.* 63:1583-1586.
- Scotland, S.M., G.A. Willshaw, H.R. Smith and B. Rowe. 1990. Properties of strains of *Escherichia coli* O26:H11 in relation to their enteropathogenic or enterohemorrhagic classification. *J. Infect. Dis.* 162:1069-1074.
- Scott, T., C. Wilson, D. Bailey, T. Klopfenstein, T. Milton, R. Moxley, D. Smith, J. Gray and L. Hungerford. 2000. Influence of diet on total and acid resistant *E. coli* and colonic pH. *Nebraska Beef Rep.* 39-41.
- Shin, R., M. Suzuki and Y. Morishita. 2002. Influence of intestinal anaerobes and organic acids on the growth of enterohaemorrhagic *Escherichia coli* O157:H7. *J. Med. Microbiol.* 51:201-206.
- Smith, D.R., R.A. Moxley, S.L. Clowser, J.D. Folmer, S. Hinkley, G.E. Erickson and T.J. Klopfenstein. 2005. Use of rope devices to describe and explain the feedlot ecology of *Escherichia coli* O157:H7 by time and place. *Foodborne Path. Dis.* 2:50-60.
- Stanford, K., D. Croy, S.J. Bach, G.L. Wallins, H. Zahiroddini and T.A. McAllister. 2005. Ecology of *Escherichia coli* O157:H7 in commercial dairies in southern Alberta. *J. Dairy Sci.* 88:4441.
- Stanton, T.L. and D. Schutz. 2000. Effect of switching from high grain to hay five days prior to slaughter on finishing cattle performance.
- Steinmuller, N., L. Demma, J.B. Bender, M. Eidson and F.J. Angulo. 2006. Outbreaks of enteric disease associated with animal contact: Not just a foodborne problem anymore. *Clin. Infect. Dis.* 43:1596-1602.
- Stephens, T.P., G.H. Loneragan, E. Karunasena and M.M. Brashears. 2007. Reduction of *Escherichia coli* O157 and *Salmonella* in feces and on hides of feedlot cattle using various doses of a direct-fed microbial. *J. Food Prot.* 70:2386-2391.
- Synge, B.A., M.E. Chase-Topping, G.F. Hopkins, I.J. McKendrick, F. Thomson-Carter, D. Gray, S.M. Rusbridge, F.I. Munro, G. Foster and G.J. Gunn. 2003. Factors influencing the shedding of verocytotoxin-producing *Escherichia coli* O157 by beef suckler cows. *Epidemiol.*

- Infect. 130:301-312.
- Thran, B.H., H.S. Hussein, M.R. Hall and S.F. Khaiboullina. 2001. Shiga toxin-producing *Escherichia coli* in beef heifers grazing an irrigated pasture. *J. Food Prot.* 64:1613-1616.
- Thurston-Enriquez, J.A., J.E. Gilley and B. Eghball. 2005. Microbial quality of runoff following land application of cattle manure and swine slurry. *J. Water Health.* 3:157-171.
- Tkalcic, S., C.A. Brown, B.G. Harmon, A.V. Jain, E.P.O. Mueller, A. Parks, K.L. Jacobsen, S.A. Martin, T. Zhao and M.P. Doyle. 2000. Effects of diet on rumen proliferation and fecal shedding of *Escherichia coli* O157:H7 in calves. *J. Food Prot.* 63:1630-1636.
- Tkalcic, S., T. Zhao, B.G. Harmon, M.P. Doyle, C.A. Brown and P. Zhao. 2003. Fecal shedding of enterohemorrhagic *Escherichia coli* in weaned calves following treatment with probiotic *Escherichia coli*. *J. Food Prot.* 66:1184-1189.
- USDA-APHIS. 1997. An update: *Escherichia coli* O157:H7 in humans and cattle.
- USDA-ERS. 2001. ERS estimates foodborne disease costs at \$6.9 billion per year. Economic Research Service-United States Department of Agriculture. Last Accessed: 16 October 2007, 2004. Available at: <http://www.ers.usda.gov/publications/aer741/aer741.pdf>
- Van Baale, M.J., J.M. Sargeant, D.P. Gnad, B.M. DeBey, K.F. Lechtenberg and T.G. Nagaraja. 2004. Effect of forage or grain diets with or without monensin on ruminal persistence and fecal *Escherichia coli* O157:H7 in cattle. *Appl. Environ. Microbiol.* 70:5336-5342.
- Van Immerseel, F., J.B. Russell, M.D. Flythe, I. Gantois, L. Timbermont, F. Pasmans, F. Haesebrouck and R. Ducatelle. 2006. The use of organic acids to combat *Salmonella* in poultry: A mechanistic explanation of the efficacy. *Avian Pathol.* 35:182-188.
- Van Kessel, J.S., P.C. Nedoluha, A. Williams-Campbell, R.L. Baldwin and K.R. McLeod. 2002. Effects of ruminal and post-ruminal infusion of starch hydrolysate or glucose on the microbial ecology of the gastrointestinal tract in growing steers. *J. Anim. Sci.* 80:3027-3034.
- Vugia, D., A. Cronquist, J. Hadler, M. Tobin-D'Angelo, D. Blythe, K. Smith, S. Lathrop, D. Morse, P. Cieslak, T. Jones, K.G. Holt, J.J. Guzewich, O.L. Henao, E. Scallan, F.J. Angulo, P.M. Griffin and R.V. Tauxe. 2007. Preliminary FoodNet data on the incidence of infection with aathogens transmitted commonly through food --- 10 states, 2006. *Morbidity and Mortality Weekly Report.* 56:336-339.
- Wallace, R.J., M.L. Falconer and P.K. Bhargava. 1989. Toxicity of volatile fatty acids at rumen pH prevents enrichment of *Escherichia coli* by sorbitol in rumen contents. *Current Microbiol.* 19:277-281.
- Wang, G., T. Zhao and M.P. Doyle. 1996. Fate of enterohemorrhagic *Escherichia coli* O157:H7 in bovine feces. *Appl. Environ. Microbiol.* 62:2567-2570.
- Wang, L., D. Rothmund, H. Curd and P.R. Reeves. 2000. Sequence diversity of the *Escherichia coli* H7 fliC genes: implication for a DNA-based typing scheme for *E. coli* O157:H7. *J. Clin. Microbiol.* 38:1786-1790.
- Wells, J.E., E.D. Berry and V.H. Varel. 2005. Effects of common forage phenolic acids on *Escherichia coli* O157:H7 viability in bovine feces. *Appl. Environ. Microbiol.* 71:7974-7979.
- Winfield, M.D. and E.A. Groisman. 2003. Role of nonhost environments in the lifestyles of *Salmonella* and *Escherichia coli*. *Appl. Environ. Microbiol.* 69:3687-3694.
- Woerner, D.R., J.R. Ransom, J.N. Sofos, G.A. Dewell, G.C. Smith, M.D. Salman and K.E. Belk. 2006. Determining the prevalence of *Escherichia coli* O157 in cattle and beef from the feedlot to the cooler. *J. Food Prot.* 69:2824-2827.
- Wolin, M.J. 1969. Volatile fatty acids and the inhibition of *Escherichia coli* growth by rumen fluid. *Appl. Microbiol.* 17:83-87.
- Yokoyama, M.G. and K.A. Johnson. 1988. Microbiology of the rumen and intestine. In: *The Ruminant Animal: Digestive Physiology and nutrition.* Church, D.C. (eds), Waveland Press, Englewood Cliffs, NJ pp. 125-144.
- Yoon, I.K. and M.D. Stern. 1996. Effects of *Saccharomyces cerevisiae* and *Aspergillus oryzae* cultures on ruminal fermentation in dairy cows. *J. Dairy Sci.* 79:411-417.
- Younts-Dahl, S.M., M.L. Galyean, G.H. Loneragan, N.A. Elam and M.M. Brashears. 2004. Dietary supplementation with Lactobacillus- and Propionibacterium-based direct-fed microbials and prevalence of *Escherichia coli* O157 in beef feedlot cattle and on hides at harvest. *J. Food Prot.* 67:889-893.
- Zhao, T., M.P. Doyle, B.G. Harmon, C.A. Brown, P.O.E. Mueller and A.H. Parks. 1998. Reduction of carriage of enterohemorrhagic *Escherichia coli* O157:H7 in cattle by inoculation with probiotic bacteria. *J. Clin. Microbiol.* 36:641-647.
- Zhao, T., S. Tkalcic, M.P. Doyle, B.G. Harmon, C.A. Brown and P. Zhao. 2003. Pathogenicity of enterohemorrhagic *Escherichia coli* in neonatal calves and evaluation of fecal shedding by treatment with probiotic *Escherichia coli*. *J. Food Prot.* 66:924-930.

Caliciviruses

Molecular and Cellular Virology

Edited by: G.S. Hansman, J. Jiang, K.Y. Green
c. 250 pp., April 2010

ISBN: 978-1-904455-63-9 \$310 / £159

The most important research findings.
Timely and comprehensive reviews.
Discussion of past and current research.

Epstein-Barr Virus

Latency and Transformation

Edited by: Erle S. Robertson
c. 220 pp., April 2010

ISBN: 978-1-904455-62-2 \$310 / £159

Expert virologists comprehensively review this important subject from a genetic, biochemical, immunological, and cell biological perspective. Essential reading.

Anaerobic Parasitic Protozoa

Genomics and Molecular Biology

Edited by: C.G. Clark, P.J. Johnson, R.D. Adam
c. 210 pp., March 2010

ISBN: 978-1-904455-61-5 \$310 / £159

Internationally acclaimed researchers critically review the most important aspects of research on anaerobic parasitic protozoa.

Lentiviruses and Macrophages

Molecular and Cellular Interactions

Edited by: Moira Desport
c. 410 pp., March 2010

ISBN: 978-1-904455-60-8 \$310 / £159

Top lentivirus and macrophage specialists comprehensively review cutting-edge topics in the molecular and cellular biology of the lentivirus-macrophage interaction.

Microbial Population Genetics

Edited by: Jianping Xu
c. 230 pp., March 2010

ISBN: 978-1-904455-59-2 \$310 / £159

Details the major current advances in microbial population genetics and genomics.

Borrelia

Molecular Biology, Host Interaction and Pathogenesis

Edited by: D. Scott Samuels and Justin D. Radolf
c. 630 pp., March 2010

ISBN: 978-1-904455-58-5 \$310 / £159

Written by renowned scientists in the field who have made seminal contributions to the field, this book is a comprehensive guide to the pathogenic *Borrelia*.

Influenza

Molecular Virology

Edited by: Qinghua Wang and Yizhi Jane Tao
c. 200 pp., February 2010

ISBN: 978-1-904455-57-8 \$310 / £159

NS1, hemagglutinin, nucleoprotein, glycoproteins, M2 channel, virulence, polymerase, microarrays, vaccine design.

RNA Interference and Viruses

Current Innovations and Future Trends

Edited by: Miguel Angel Martínez
c. 280 pp., February 2010

ISBN: 978-1-904455-56-1 \$310 / £159

Expert RNAi specialists from around the world have teamed up to produce a timely and thought-provoking review of the area.

Retroviruses

Molecular Biology, Genomics and Pathogenesis

Edited by: Reinhard Kurth and Norbert Bannert
c. 520 pp., January 2010

ISBN: 978-1-904455-55-4 \$310 / £159

Genomics, molecular biology and pathogenesis, comprehensively covering all the recent advances.

Metagenomics

Theory, Methods and Applications

Edited by: Diana Marco
x + 212 pp., January 2010

ISBN: 978-1-904455-54-7 \$310 / £159

Essential reading for all researchers performing metagenomics studies. Highly recommended.

Aspergillus

Molecular Biology and Genomics

Edited by: M. Machida and K. Gomi
x + 238 pp., January 2010

ISBN: 978-1-904455-53-0 \$310 / £159

Systematics, bioinformatics, systems biology, regulation, genetics, genomics, metabolism, ecology, development.

Environmental Molecular Microbiology

Edited by: Wen-Tso Liu and Janet K. Jansson
viii + 232 pp., January 2010

ISBN: 978-1-904455-52-3 \$310 / £159

Current technology and applications. Microbial diversity, phylogeny, communities, 16S rRNA, metagenomics, metaproteomics, microarrays, fingerprinting, soil, water, plants, humans, biofilms.

Neisseria

Molecular Mechanisms of Pathogenesis

Edited by: Caroline Genco and Lee Wetzler
x + 270 pp., January 2010

ISBN: 978-1-904455-51-6 \$310 / £150

Genomics, biofilms, adhesion, invasion, immunity, complement, apoptosis, vaccine, epidemiology, antibiotic resistance.

Frontiers in Dengue Virus Research

Edited by: K.A. Hanley and S.C. Weaver
viii + 304 pp., January 2010

ISBN: 978-1-904455-50-9 \$310 / £150

Evolution, epidemiology, translation, replication, pathogenesis, host, animal models, mosquito interactions, transmission, vaccine, drugs, immunotherapy.

ABC Transporters in Microorganisms

Edited by: Alicia Ponte-Sucre
xii + 260 pp., August 2009

ISBN: 978-1-904455-49-3 \$310 / £150

Pili and Flagella

Current Research and Future Trends

Edited by: Ken Jarrell

x + 238 pp., August 2009

ISBN: 978-1-904455-48-6 \$310 / £150

Lab-on-a-Chip Technology

Edited by: K. E. Herold and A. Rasooly

Vol 1: Fabrication and Microfluidics

ISBN: 978-1-904455-46-2 \$310 / £150

Vol 2: Biomolecular Separation and Analysis

ISBN: 978-1-904455-47-9 \$310 / £150

Bacterial Polysaccharides

Current Research and Future Trends

Edited by: Matthias Ullrich

xii + 358 pp., June 2009

ISBN: 978-1-904455-45-5 \$310 / £150

Microbial Toxins:

Current Research and Future Trends

Edited by: Thomas Proft

viii + 192 pp., May 2009

ISBN: 978-1-904455-44-8 \$310 / £150

Acanthamoeba:

Biology and Pathogenesis

Author: Naveed Khan

viii + 290 pp., February 2009

ISBN: 978-1-904455-43-1 \$310 / £150

Bacterial Secreted Proteins: Secretory Mechanisms and Role in Pathogenesis

Edited by: Karl Wooldridge

xii + 512 pp., April 2009

ISBN: 978-1-904455-42-4 \$310 / £150