

# Estimating the Relative Role of Various Subcategories of Food, Water, and Animal Contact Transmission of 28 Enteric Diseases in Canada

Ainslie J. Butler, Katarina D.M. Pintar, and M. Kate Thomas

## Abstract

**Objective:** Enteric illness represents a significant burden of illness in Canada and internationally. Building on previous research, an expert elicitation was undertaken to explore the routes of transmission for 28 pathogens involved in enteric illness in Canada. This article considers the subcategories of foodborne, waterborne, and animal contact transmission.

**Methods:** As part of an expert elicitation, 31 experts were asked to provide estimates of source attribution for subcategories of foodborne ( $n=15$ ), waterborne ( $n=10$ ), and animal contact ( $n=3$ ) transmission. The results from an online survey were combined using triangular probability distributions, and median and 90% credible intervals were produced. The total proportion and estimated number of cases of enteric illness attributable to each type of food commodity, water source, and animal exposure route were calculated using results from the larger elicitation survey and from a recent Canadian foodborne burden of illness study (Thomas *et al.*, 2013).

**Results:** Thirty experts provided foodborne subcategory estimates for 15/28 pathogens, waterborne subcategory estimates for 14/28 pathogens and animal contact subcategory estimates for 5/28. The elicitation identified raw produce, recreational water, and farm animal contact as important risk factors for enteric illness. These results also highlighted the complexity of transmission, with greater uncertainty for certain pathogens and routes of transmission.

**Conclusions:** This study is the first of its kind to explore subcategories of foodborne, waterborne, and animal contact transmission across such a range of enteric pathogens. Despite inherent uncertainty, these estimates present an important quantitative synthesis of the roles of foodborne commodities, water sources, and pathways of animal contact in the transmission of enteric illness in Canada.

## Introduction

ENTERIC INFECTIONS REPRESENT a significant burden of illness in Canada and globally. In Canada, there are an estimated 20.5 million cases of enteric illness annually (95% confidence interval [CI]: 19.3–21.7 million) of which 4.0 million (95% CI: 3.1–5.8 million) are estimated to be domestically acquired foodborne cases (Thomas *et al.*, 2013). The pathways to human enteric illness are varied and complex and often poorly characterized.

Expert elicitation has previously been used as a source attribution tool, to explore the role of transmission in the most common enteric infections in Canada and around the world (Cressey and Lake, 2005; Hoffmann *et al.*, 2006; Havelaar *et al.*, 2008; Ravel *et al.*, 2010; Vally *et al.*, 2014; Butler *et al.*, 2015). Some have focused particularly on subcategories of foodborne transmission (Hoffmann *et al.*, 2006;

Havelaar *et al.*, 2008; Davidson *et al.*, 2011), but very limited data exist on waterborne and animal contact routes.

As part of the Public Health Agency of Canada's (PHAC) burden of enteric illness and source attribution efforts, an expert elicitation was undertaken to attribute transmission of 28 enteric pathogens (Butler *et al.*, 2015). An objective of this study involved the elicitation of specific subcategory estimates for foodborne, waterborne, and animal contact transmission of these same pathogens, which is the focus of this article. This research will help inform future research and surveillance efforts in Canada.

## Materials and Methods

A six-stage expert elicitation for the attribution of 28 enteric illnesses to five major routes of transmission and their respective subcategories was undertaken in January–April

---

Centre for Foodborne, Environmental, and Zoonotic Infectious Diseases, Public Health Agency of Canada, Ottawa, Canada.

© Ainslie J. Butler, et al. 2016; Published by Mary Ann Liebert, Inc. This Open Access article is distributed under the terms of the Creative Commons Attribution Noncommercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits any non-commercial use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

2014, methods described elsewhere (Butler *et al.*, 2015). Briefly, a panel of 31 Canadian experts were each assigned 10 pathogens, based on self-assessed expertise collected in a presurvey.

Experts were asked to attribute 100 domestically acquired cases transmitted via food, water, or animal contact to their subcategories (Supplementary Material S1; Supplementary Data are available online at [www.liebertpub.com/fpd](http://www.liebertpub.com/fpd)). Experts could decline to produce subcategory estimates. Food subcategories in this study were based on Centers for Disease Control and Prevention guidelines for outbreak reporting (Painter *et al.*, 2009), used in previous elicitation (Hoffmann *et al.*, 2006; Havelaar *et al.*, 2008; Davidson *et al.*, 2011). Waterborne subcategories were differentiated by source water (recreational, bottled, private well, municipal groundwater, municipal surface water, and municipal ground water under the influence of surface water [GUDI]) and size of system (municipal system > or <10,000 population serviced) (Murphy *et al.*, 2015). Animal contact subcategories included exposure to wildlife, or domestic or farm animals. Experts were prompted to consider whether their foodborne subcategory estimates would change if considering attribution at the point of entry of contaminated foods into the domestic or commercial kitchen, and if so to quantify attribution of foodborne subcategories at the kitchen door.

Estimates were excluded where experts assessed confidence as 1/5 (low). Median and 90% credible intervals (CrI) for each subcategory were calculated from cumulative triangular probability distributions built using the Microsoft Excel add-in @Risk (Version 6.1.2; Palisade Corporation, Newfield, NY). Statistical methods are described elsewhere (Butler *et al.*, 2015).

Cumulative estimates of transmission were combined with estimated distributions of annual cases for 28 pathogens obtained from recent Canadian burden of enteric illness estimates (Thomas *et al.*, 2013). For enterotoxigenic *Escherichia coli* (EPEC), verotoxin-producing *E. coli* (VTEC) non-O157 and *E. coli*, other (non-VTEC, non-EPEC strains of *E. coli*), estimated domestic cases were determined by multiplying the domestic cases of VTEC O157 from Thomas *et al.* (2013) by the identified ratios relative to VTEC O157 from Thomas *et al.* (2013), derived from Scallan *et al.* (2011) and Chui *et al.* (2011). The estimated distributions of domestic cases for each pathogen ( $N_{\text{pathogen}}$ ) were multiplied by the cumulative distributions estimated via each of food, water, and animal contact transmission (e.g.,  $P_{\text{pathogen}}[\text{food}]$ ), derived from the first stage of the elicitation survey (Butler *et al.*, 2015) using @Risk. These estimates were then multiplied by the distributions of the proportion of cases transmitted by each subcategory (e.g.,  $P_{\text{pathogen}}[\text{beef}]$ ) to estimate how many cases were attributable to specific commodities (e.g.,  $N_{\text{beef}}$ ), and their 90% CrI.

Ethics approval was obtained from Health Canada and PHAC's Research Ethics Board on January 13, 2014 (REB 2013-0033).

## Results

Attribution estimates for specific subcategories of foodborne, waterborne, and animal contact transmission routes were provided by 30 of 31 participating experts. Responses were excluded where confidence was ranked as 1/5 (low),

representing 7/148 (5%) foodborne, 3/71 (4%) waterborne, and 6/70 (9%) animal contact transmission estimates. Data were also excluded for foodborne ( $n=3$ ) and waterborne ( $n=1$ ) transmission where major transmission route analysis indicated inappropriate clusters of responses (implying nonenteric transmission) (Butler *et al.*, 2015).

Five or more experts provided foodborne subcategory estimates for 15/28 pathogens; waterborne subcategory estimates for 5/28 pathogens; and animal contact estimates for 5/28 pathogens. No subcategory estimates were provided for pathogens where less than 10 cases per 100 were attributed to that particular route (e.g., 8.3% of adenovirus transmission attributed to food and no foodborne subcategory estimates were provided).

Kitchen door attribution estimates for foodborne transmission were provided for 13/28 pathogens, with an average of two expert estimates per pathogen where provided. The shifts in attribution calculated from these estimates were negligible, but implied that cross-contamination, particularly from (raw) poultry in the kitchen, may play an important role in disease transmission, especially for *Campylobacter* spp. and nontyphoidal *Salmonella* spp.

Medians and 90% CrI from the cumulative distributions for foodborne, waterborne, and animal contact transmission subcategories are presented in Tables 1, 2, and 3, respectively, for pathogens with  $\geq 5$  experts responding. The elicited probability distributions should be interpreted with caution considering the uncertainty demonstrated in the CrI.

Using total pathogen estimates from the Canadian foodborne burden of illness estimates (Thomas *et al.*, 2013), 90% of cases of enteric illness with the 28 pathogens of interest were attributed to viruses, 8% to bacteria and 2% to parasites.

### Food

For the foodborne estimates, raw produce was most frequently identified as the dominant source (3/14 pathogens) (Table 1); 21.8% (90% CrI: 8.1–37.5) of all foodborne cases were attributed to raw produce, 18.3% (90% CrI: 7.2–31.2) to seafood, and <10% each to the remaining subcategories (Fig. 1). For bacterial infections, the most attributed food vehicle was “other” (15.4%; 90% CrI: 3.2–28.7), followed by poultry (12.7%; 90% CrI: 2.9–24.8); raw produce was the most common food vehicle for parasitic (25.2%; 90% CrI: 4.6–54.3) and viral (29.1%; 90% CrI: 0.0–48.9) infections (Fig. 1).

### Water

For waterborne transmission, recreational water sources were the most commonly identified subcategory (3/5) (Table 2). Recreational water was attributed to the most cases overall within the waterborne transmission category (23.8%; 90% CrI: 2.6–52.8), and specifically for parasites (24.2%; 90% CrI: 8.3–46.1) (Fig. 2). Private well water was most frequently implicated for bacterial (29.4%; 90% CrI: 2.4–64.9) and viral infections (23.5%; 90% CrI: 1.0–53.0) (Fig. 2).

### Animal

Farm animal contact was the most common vehicle within animal contact transmission (5/5) (Table 3). The majority of

TABLE 1. MOST LIKELY PROPORTION AND 90% CREDIBLE INTERVAL OF FOODBORNE TRANSMISSION ATTRIBUTABLE TO SUBCATEGORIES FOR 14 OF 28 PATHOGENS AT THE POINT OF CONSUMPTION FOR WHICH MORE THAN FIVE EXPERTS PRODUCED ESTIMATES

	<i>Campylobacter spp.</i>	<i>Clostridium botulinum</i>	<i>Clostridium perfringens</i>	<i>Cyclospora cayetanensis</i>	<i>Giardia spp.</i>
Count	8	5	5	9	5
Beef	5.9 (1.4–18.7)	2.4 (0.2–7.5)	16.8 (1.4–47.0)	5.9 (0.5–23.3)	10.8 (0.9–32.9)
Beverages	2.9 (0.2–17.6)	2.3 (0.2–8.6)	0.1 (0.0–0.5)	0.4 (0.0–2.3)	3.2 (0.3–12.2)
Breads & bakery	2.9 (0.2–17.5)	0.7 (0.0–2.8)	0.3 (0.0–1.1)	0.4 (0.0–2.3)	0.1 (0.0–0.5)
Dairy	9.5 (0.1–27.5)	0.2 (0.0–1.1)	1.4 (0.0–5.3)	2.2 (0.0–12.4)	1.6 (0.0–5.4)
Eggs	7.8 (0.6–27.7)	0.0 (0.0–0.0)	2.8 (0.2–7.7)	0.4 (0.0–2.3)	1.1 (0.0–5.0)
Game	3.5 (0.3–17.5)	17.4 (1.4–50.3)	17.0 (1.4–49.2)	0.4 (0.0–2.3)	4.4 (0.3–25.7)
Lamb & goat	3.2 (0.3–17.7)	2.1 (0.2–6.9)	4.7 (0.4–14.1)	0.4 (0.0–2.3)	4.6 (0.3–25.7)
Lunchmeat	4.6 (0.4–17.8)	2.9 (0.3–9.1)	4.4 (0.4–12.3)	0.9 (0.1–3.3)	1.1 (0.0–5.0)
Nuts & seeds	3.1 (0.2–17.6)	1.2 (0.0–5.4)	0.1 (0.0–0.5)	0.9 (0.1–3.3)	0.1 (0.0–0.5)
Pork	4.1 (0.3–18.1)	1.8 (0.1–6.1)	6.0 (0.5–16.4)	2.0 (0.1–11.0)	3.9 (0.3–18.5)
Poultry	33.0 (9.6–48.9)	1.8 (0.1–6.0)	14.8 (1.4–38.5)	0.4 (0.0–2.4)	1.1 (0.0–5.0)
Produce, cooked	4.4 (0.4–17.9)	25.6 (7.6–43.4)	1.9 (0.1–8.1)	1.2 (0.1–4.8)	0.4 (0.0–2.4)
Produce, raw	9.1 (2.4–22.0)	2.9 (0.2–9.2)	0.6 (0.1–1.8)	61.7 (20.8–85.0)	56.7 (28.9–75.5)
Seafood	3.1 (0.2–17.8)	19.6 (6.0–43.1)	0.7 (0.0–2.8)	3.0 (0.1–18.0)	5.4 (0.4–28.0)
Other	3.0 (0.0–17.5)	19.1 (1.8–51.7)	28.4 (2.4–72.5)	19.9 (1.6–53.8)	5.6 (0.5–15.3)

	<i>Hepatitis A</i>	<i>Listeria monocytogenes</i>	<i>Salmonella spp., nontyphoidal</i>	<i>Trichinella spp.</i>	<i>Verotoxin-producing Escherichia coli (VTEC) non-O157</i>
Count	5	9	11	9	6
Beef	3.4 (0.2–19.0)	6.5 (0.5–22.0)	6.2 (0.5–23.4)	1.7 (0.1–5.8)	34.9 (14.8–51.1)
Beverages	4.8 (0.8–19.2)	3.1 (0.2–18.2)	4.1 (0.3–22.9)	0.0 (0.0–0.0)	4.1 (0.3–17.1)
Breads & bakery	5.8 (0.5–20.7)	3.0 (0.1–17.6)	4.0 (0.3–22.9)	0.0 (0.0–0.0)	2.8 (0.1–17.0)
Dairy	0.8 (0.0–3.1)	17.0 (0.0–29.6)	5.2 (0.0–23.4)	0.0 (0.0–0.0)	7.6 (0.0–24.5)
Eggs	3.2 (0.2–18.6)	2.9 (0.1–17.8)	10.8 (2.1–28.6)	0.0 (0.0–0.0)	2.8 (0.1–17.2)
Game	3.1 (0.2–19.3)	3.4 (0.3–17.7)	4.3 (0.3–23.4)	64.0 (14.1–92.2)	3.6 (0.3–17.3)
Lamb & goat	3.2 (0.2–19.2)	3.3 (0.2–18.4)	4.4 (0.4–23.6)	1.8 (0.1–5.7)	4.1 (0.3–18.2)
Lunchmeat	8.0 (0.7–28.2)	24.8 (11.6–46.8)	4.4 (0.3–23.2)	0.5 (0.0–2.6)	4.5 (0.4–23.9)
Nuts & seeds	3.7 (0.3–19.7)	3.0 (0.2–17.8)	6.5 (0.7–24.4)	0.0 (0.0–0.0)	5.7 (0.5–21.2)
Pork	3.5 (0.2–19.2)	6.4 (0.5–23.9)	6.0 (0.5–23.0)	31.0 (2.6–81.1)	4.1 (0.3–21.0)
Poultry	3.3 (0.2–19.5)	3.3 (0.2–18.1)	23.6 (10.1–37.8)	0.4 (0.0–2.5)	4.1 (0.3–20.9)
Produce, cooked	7.1 (0.6–27.7)	3.2 (0.2–17.8)	3.6 (0.2–23.0)	0.0 (0.0–0.0)	3.5 (0.2–20.5)
Produce, raw	34.2 (10.0–52.2)	8.9 (0.8–25.3)	7.8 (0.6–25.4)	0.0 (0.0–0.0)	11.0 (2.4–27.3)
Seafood	12.9 (4.4–31.4)	6.1 (0.5–18.8)	4.1 (0.3–22.7)	0.0 (0.0–0.0)	4.3 (0.3–20.8)
Other	3.1 (0.1–18.9)	5.0 (0.4–19.2)	4.9 (0.4–22.6)	0.6 (0.1–1.8)	2.9 (0.1–18.3)

	<i>Verotoxin-producing Escherichia coli (VTEC) O157</i>	<i>Vibrio parahaemolyticus</i>	<i>Vibrio vulnificus</i>	<i>Yersinia enterocolitica</i>
Count	8	9	5	11
Beef	47.4 (29.2–67.3)	0.0 (0.0–0.0)	0.0 (0.0–0.0)	2.5 (0.2–9.6)
Beverages	2.2 (0.2–6.6)	0.5 (0.0–1.8)	0.0 (0.0–0.0)	0.5 (0.0–2.2)
Breads & bakery	1.0 (0.1–4.3)	0.3 (0.0–1.2)	0.0 (0.0–0.0)	0.4 (0.0–2.2)
Dairy	10.4 (0.0–20.9)	0.3 (0.0–1.2)	0.0 (0.0–0.0)	8.1 (0.0–17.4)
Eggs	0.5 (0.0–2.6)	0.0 (0.0–0.0)	0.0 (0.0–0.0)	1.8 (0.1–7.0)
Game	2.8 (0.3–7.4)	0.7 (0.0–3.0)	0.0 (0.0–0.0)	1.7 (0.1–4.7)
Lamb & goat	2.1 (0.2–6.6)	0.0 (0.0–0.0)	0.0 (0.0–0.0)	3.5 (0.3–14.0)
Lunchmeat	3.9 (0.3–13.7)	0.3 (0.0–1.2)	0.0 (0.0–0.0)	5.3 (0.5–14.9)
Nuts & seeds	3.7 (0.3–11.8)	0.0 (0.0–0.0)	0.0 (0.0–0.0)	1.1 (0.1–3.8)
Pork	2.0 (0.2–6.0)	0.0 (0.0–0.0)	0.0 (0.0–0.0)	63.8 (42.9–77.2)
Poultry	1.2 (0.1–5.3)	0.0 (0.0–0.0)	0.0 (0.0–0.0)	1.2 (0.1–4.5)
Produce, cooked	1.2 (0.1–5.3)	0.0 (0.0–0.0)	0.0 (0.0–0.0)	0.7 (0.1–2.7)
Produce, raw	19.5 (8.4–37.5)	1.8 (0.2–6.1)	0.0 (0.0–0.0)	5.6 (0.5–16.6)
Seafood	0.8 (0.1–3.1)	96.2 (75.0–100.0)	100.0 (93.1–100.0)	3.3 (0.2–10.3)
Other	1.4 (0.1–5.5)	0.0 (0.0–0.0)	0.0 (0.0–0.0)	0.7 (0.1–2.8)

TABLE 2. MOST LIKELY PROPORTION AND 90% CREDIBLE INTERVAL ATTRIBUTED TO THE 10 SUBCATEGORIES ASSOCIATED WITH WATERBORNE TRANSMISSION FOR 5 OF 28 PATHOGENS FOR WHICH MORE THAN FIVE EXPERTS PRODUCED ESTIMATES

	Campylobacter <i>spp.</i>	Cryptosporidium <i>spp.</i>	Giardia <i>spp.</i>	VTEC non-O157 <sup>a</sup>	VTEC O157 <sup>b</sup>
Count	6	7	7	5	5
Recreational water	22.0 (5.3–44.8)	18.7 (6.0–36.3)	32.1 (12.7–55.1)	21.4 (7.6–39.9)	48.5 (21.3–83.2)
Private well	27.2 (13.1–44.5)	15.0 (2.4–37.7)	19.7 (8.0–34.2)	29.0 (14.4–47.2)	25.6 (5.9–47.1)
Ground water: large municipal system	5.2 (0.4–25.7)	7.4 (0.6–30.8)	1.7 (0.1–5.7)	6.1 (0.4–26.0)	0.9 (0.1–3.3)
Ground water: small municipal system	5.5 (0.5–22.6)	8.5 (0.7–31.2)	2.9 (0.2–8.1)	7.2 (1.4–26.4)	4.9 (1.3–11.0)
Surface water: large municipal system	4.5 (0.3–22.6)	8.4 (0.7–30.5)	13.0 (1.1–37.5)	4.7 (0.3–29.6)	1.6 (0.1–5.9)
Surface water: small municipal system	7.7 (0.8–23.7)	12.0 (2.1–32.2)	9.0 (0.9–23.5)	7.6 (1.5–29.4)	7.5 (2.1–16.0)
GUDI: large municipal system	4.0 (0.3–22.4)	8.0 (0.7–30.4)	6.9 (0.6–28.0)	6.4 (0.9–30.1)	2.1 (0.2–6.9)
GUDI: small municipal system	12.5 (1.1–12.5)	10.1 (1.0–10.1)	6.7 (0.6–6.7)	7.4 (1.4–7.4)	7.8 (2.1–7.8)
Bottled water	3.8 (0.2–22.5)	7.7 (0.6–30.9)	4.3 (0.3–19.0)	5.4 (0.4–26.1)	1.0 (0.1–3.1)
Other	7.6 (0.6–25.2)	4.2 (0.3–17.5)	3.8 (0.3–14.5)	4.7 (0.2–29.8)	0.0 (0.0–0.0)

<sup>a</sup>VTEC non-O157, verotoxin-producing *Escherichia coli*.

<sup>b</sup>VTEC O157, verotoxin-producing *Escherichia coli*.

GUDI, ground water under the influence of surface water.

animal contact cases were attributed to farm animals (51.2%; 90% CrI: 5.5–100.0). Farm animal contact was identified as the main pathway for bacteria (54.0%; 90% CrI: 8.5–100.0), parasites (44.3%; 90% CrI: 6.6–92.1), and viruses (51.4%; 90% CrI: 4.3–100.0) (Fig. 3).

## Discussion

This study attributes the number of enteric cases related to 15 of 28 pathogens of public health importance in Canada to the subcategories of major enteric illness transmission routes (i.e., foodborne, waterborne, and animal contact). This study is one of the first to explore such an expansive list of enteric pathogens and has considered a wide range of viruses not considered in previous studies (Supplementary Material S2). Exploring the dominant sources of exposure to enteric pathogens, across and within transmission routes, is critical for public health initiatives targeted at reducing the burden of enteric illness in Canada. Identifying existing knowledge gaps related to enteric disease transmission dynamics in Canada is one outcome of this work.

## Food


Raw produce was most frequently identified as the dominant source of foodborne transmission (7/24 pathogens) (Table 1). Of all foodborne cases, 21.8% (90% CrI: 8.1–37.5) were attributed to raw produce, 18.3% (90% CrI: 7.2–31.2) to seafood, and <10% each to the remaining subcategories (Fig. 1). For bacterial infections, the most attributed food vehicle was “other” (15.4%; 90% CrI: 3.2–28.7), followed by poultry (12.7%; 90% CrI: 2.9–24.8); raw produce was the most common food vehicle for parasitic (25.2%; 90% CrI: 4.6–54.3) and viral (29.1%; 90% CrI: 0.0–48.9) infections (Fig. 1). These results are similar to estimates from previous elicitations and from analysis of outbreak data (Supplementary Material S2).

Infection with *Trichinella spp.* is attributed largely to game (64.0%; 90% CrI: 14.1–92.2) and pork (31.0%; 90% CrI: 2.6–81.1) in the current study. Routine testing of swine as part of the Canadian Food Inspection Agency (CFIA) *Trichinella* control program (CFIA, 2013) demonstrates that domestic commercial herds are *Trichinella*-free, despite a recent

TABLE 3. MOST LIKELY PROPORTION AND 90% CREDIBLE INTERVAL ATTRIBUTABLE TO THE THREE SUBCATEGORIES ASSOCIATED WITH ANIMAL CONTACT TRANSMISSION FOR 5 OF 28 PATHOGENS FOR WHICH MORE THAN FIVE EXPERTS PRODUCED ESTIMATES

	Campylobacter <i>spp.</i>	Giardia <i>spp.</i>	Salmonella <i>spp.</i> , <i>nontyphoidal</i>	VTEC O157 <sup>a</sup>	Yersinia enterocolitica
Count	8	7	9	7	7
Domestic	27.5 (8.0–53.6)	42.5 (17.0–66.8)	39.8 (11.8–75.9)	5.3 (3.4–7.3)	23.3 (1.8–62.6)
Farm	57.6 (35.1–87.0)	48.0 (15.8–73.4)	52.6 (21.3–87.6)	83.0 (76.8–89.1)	72.5 (31.7–95.8)
Wildlife	14.9 (1.2–41.2)	9.5 (0.8–30.9)	7.6 (1.9–13.3)	11.7 (6.5–16.9)	4.2 (0.4–10.7)

<sup>a</sup>VTEC O157, Verotoxin-producing *Escherichia coli*.



**FIG. 1.** Proportion of foodborne cases attributed to 15 subcategories, by pathogen types and as a total. Error bars represent 90% credible intervals.


(2013) case of human *Trichinella* spp. infection from a pig raised on a noncommercial farm (CFIA, 2013). This estimate highlights two limitations of this research design: (1) the inability to differentiate between domestic and imported foods: given the rarity of human trichinellosis, the role of imported meats could be significant; and (2) reliance of experts on historic trends over current exposures.

Beef was highlighted as the most important food commodity for VTEC O157 infections (47.4%; 90% CrI: 29.2–67.3), followed by produce: 19.5% raw and 1.2% cooked (20.7% total). This is similar to the previous Canadian elicitation, where beef was estimated to be responsible for 54.0–57.8% of VTEC O157 infections, followed closely by produce (23.1–28.8%) (Davidson *et al.*, 2011). These results


suggest consistency between the two Canadian studies, illustrating that for some infections, experts remain in consensus about the dominant sources.

Where estimates differ between this study and other Canadian studies (e.g., for *Campylobacter* spp. *C. botulinum* and *Staphylococcus aureus*), sources of disagreement are unclear. The differences in responses may reflect a lack of data from which experts can derive estimates, highlighting a knowledge gap for future research, given the role some of these pathogens play in the overall burden of foodborne illness. Alternately, they may reflect considerations related to food categories explored.

For *Clostridium botulinum*, current estimates indicate an important role of cooked (preserved) produce, which may



**FIG. 2.** Proportion of waterborne cases attributed to 10 subcategories, by pathogen types and as a total. Error bars represent 90% credible intervals.



**FIG. 3.** Proportion of animal contact cases attributed to domestic, farm, and wildlife animal contact, by pathogen types and as a total. Error bars represent 90% credible intervals.

correspond to canned goods in the “other” category in retrospective outbreak studies (Greig and Ravel, 2009). However, in outbreak studies, seafood (Greig and Ravel, 2009) and game (Ravel *et al.*, 2009) are identified as the dominant risk factors.

*Cyclospora cayatanensis* transmission is mostly attributed to raw produce (61.7%; 95% CrI: 20.8–85.0), similar to the previous U.S. elicitation (Hoffmann *et al.*, 2006) and outbreak studies from the United States (Batz *et al.*, 2011) and internationally (Greig and Ravel, 2009). Epidemiological studies suggest that untreated drinking water and produce imported from endemic regions (particularly berries and salad greens) are significant risk factors, and the potential role of animals in transmission is unknown (Ortega and Sanchez, 2010).

Raw (unpasteurized) dairy consumption (especially milk) was identified in supplementary comments as an important transmission route for *Campylobacter* spp., VTEC O157, and VTEC non-O157. Raw milk was not specifically queried; however, dairy was implicated for *Campylobacter* spp. (9.5%), *Listeria monocytogenes* (17.0%), VTEC O157 (10.4%), and VTEC non-O157 (7.6%) (Table 1). The sale of raw milk is illegal in Canada, though cheeses made from raw milk are legal (Health Canada, 2014), and preliminary estimates indicate 3.1% of the Canadian population consumes raw milk and raw milk products.

### Water

No previously published expert elicitations have explored the waterborne transmission route in such detail. Recreational water was attributed to the most cases overall within the waterborne transmission category (23.8%; 90% CrI: 2.6–52.8), and specifically for parasites (24.2%; 90% CrI: 8.3–46.1) (Fig. 2). Private well water was most frequently implicated for bacterial (29.4%; 90% CrI: 2.4–64.9) and viral infections (23.5%; 90% CrI: 1.0–53.0) (Fig. 2).

Recreational water exposure was identified as the main risk factor for 3/5 pathogens for which estimates were provided (Table 2). Waterparks and pools have been recognized as important sources of *Cryptosporidium* spp. and *Giardia* spp. infections (Snel *et al.*, 2009; Yoder *et al.*, 2012a, b) and

pool-associated outbreaks in the United States (Lam *et al.*, 2014) and Canada (CCDR, 2004). *Cryptosporidium* spp., *C. cayatanensis*, and *Giardia* spp. are particularly resistant to chlorine, which is the most common type of disinfectant used in these settings (WHO, 2011). During the 1-day discussion, experts questioned the likelihood of sufficient persistence of other *Vibrio* spp. in Canadian water temperatures to facilitate waterborne infection (Tantillo *et al.*, 2004). Domestic waterborne transmission of *Vibrio* spp. through enteric routes (i.e., not wound infection) is not believed to play a major role in transmission (3.8–11.0% of *Vibrio* infection attributable to water) (Butler *et al.*, 2015). Less than five experts provided estimates for any of the *Vibrio* spp. explored in this study, reflecting a lack of Canadian knowledge on this particular route of transmission.

Private wells were the primary source identified for 2/5 pathogens (Table 2). Canadian studies have demonstrated the role of private well water as a risk for enteric illnesses (Uhlmann *et al.*, 2009; Galanis *et al.*, 2014). In this study, GUDI wells were implicated in viral infections more frequently than bacterial or parasitic infections (Fig. 2).

Attribution to municipal water systems was divided based on the size of population serviced by the municipal system: large municipal systems (>1000 population served, 13.3% of total waterborne cases) and small municipal systems (<1000 population served, 23.1%). Size is used as a proxy of level of treatment, resources available to support treatment adoption, and operator training (Murphy *et al.*, 2015).

Experts estimated that <1% of total waterborne infection was attributable to bottled water. Bottled water has been involved in several outbreaks in the United States; however, there is no current evidence of waterborne infections associated with bottled water in Canada (Brunkard *et al.*, 2011).

### Animal

This is the first expert elicitation to consider sources within the animal contact transmission route. Previous elicitations in Australia and the Netherlands explored transmission via direct animal contact (Vally *et al.*, 2014) and human and animal transmission (Havelaar *et al.*, 2008); the current study elucidates source attribution more deeply. Animal contact transmission was defined as relating to illness transmitted by exposure to animals (i.e., personal contact [hand or mouth] with animal/pet feed, animal/pet fur/coats, saliva, or feces). Farm animal exposure was estimated as the most likely transmission route for all five pathogens for which estimates are provided (Table 3). The majority of animal contact cases were attributed to farm animals (51.2%; 90% CrI: 5.5–100.0), and specifically for bacteria (54.0%; 90% CrI: 8.5–100.0), parasites (44.3%; 90% CrI: 6.6–92.1), and viruses (51.4%; 90% CrI: 4.3–100.0) (Fig. 3).

The likelihood of enteric pathogen transmission from household pets may be low compared to an encounter with farm animals or wildlife; however, the higher frequency of pet contact (David *et al.*, 2014) would suggest this is an important exposure. Expert discussion highlighted the need to consider the relative extent to which frequent pet contact contributes to human enteric illness compared to low frequency exposure to farm animals and wildlife, and the relative risks, identified as a current knowledge gap in Canada. A limitation of these estimates is evident for animal contact transmission rotavirus, for which group discussion suggested

zoonotic transmission was unlikely, yet animal contact transmission estimates have been produced (Table 3). This highlights the need to consider attribution estimates carefully in context of both the number of respondents and uncertainty as demonstrated by CrI width. A low response rate in this study is an indicator of a lack of knowledge for the specific pathogen and transmission route.

#### Study limitations

The limitations of this study are described in more detail in Butler *et al.* (2015). The study's greatest limitation is the low response rate, resulting in small panel sizes for some pathogen/product combinations. Wide confidence intervals for estimates of certain pathogens and transmission routes reflect uncertainty that arises from lack of knowledge and data, highlighting areas for future research. Recall bias related to large-scale or notable outbreaks that have occurred in Canada or globally could have skewed the expert estimates to reflect high impact, low likelihood events. This study lacks the ability to differentiate between risks from imported and domestic food exposures.

This study does not characterize variations in exposure between the local and national level. It can be conceptually difficult to balance the role of local or geographically specific exposure while producing nationally representative estimates, which contributes to the range of uncertainty presented for each estimate. This limits our ability to differentiate between the variability and uncertainty that influences the widths of the credible intervals presented herein.

#### Conclusions

Expert elicitation is a useful tool for answering difficult questions where data are unavailable or expensive to obtain. This study explored the source attribution of 15 of 28 enteric pathogens at a previously unexplored level of detail for sub-categories of foodborne, waterborne, and animal contact transmission. These results highlight the importance of raw produce, recreational water, and farm animal contact in transmission of enteric diseases as well as the complexity of transmission. While gaps in our understanding of source attribution remain, highlighting areas for future research, this study helps to improve our understanding of the role of food commodities and other sources in the transmission of enteric illness in Canada.

#### Acknowledgments

The researchers would like to thank all of the participating experts for their participation and engagement with this elicitation project. This research has been conducted under a National Sciences and Engineering Research Council postdoctoral research fellowship held by Ainslie Butler, working with the Centre for Food-borne, Environmental, and Zoonotic Infectious Diseases and FoodNet Canada (formerly C-EnterNet) with the Public Health Agency of Canada.

#### Disclosure Statement

No competing financial interests exist.

#### References

Batz MB, Hoffman S, Morris JGJ. Ranking the risks: The 10 pathogen-food combinations with the greatest burden on

public health. 2011;68. Available at: <http://epi.ufl.edu/wp-content/uploads/2011/04/RankingTheRisksREPORT.pdf>, accessed on December 30, 2015.

Brunkard JM, Ailes E, Roberts VA, Hill V, Hilborn ED, Craun GF, Rajasingham A, Kahler A, Garrison L, Hicks L, Carpenter J, Wade TJ, Beach MJ, Yoder JS. Surveillance for waterborne disease outbreaks associated with drinking water—United States, 2007–2008. *MMWR* 2011;60(ss12):38–68.

Butler AJ, Thomas MK, Pintar KDM. Expert elicitation as a means to attribute 28 enteric pathogens to foodborne, waterborne, animal contact and person-to-person transmission routes. *Foodborne Pathog Dis* 2015;12:335–344.

[CCDR] Canada Communicable Disease Report. An outbreak of *Cryptosporidium parvum* in a Surrey pool with detection in pool water sampling. *CCDR* 2004;30:61–66.

[CFIA] Canadian Food Inspection Agency. Trichinellosis—Fact Sheet. 2013. Available at: <http://www.inspection.gc.ca/animals/terrestrial-animals/diseases/reportable/trichinellosis/fact-sheet/eng/1330023015817/1330023110684>, accessed October 6, 2014.

Chui L, Lee M, Malejczyk K, Lim L, Fok D, Kwong P. Prevalence of Shiga toxin-producing *Escherichia coli* (STEC) as detected by enzyme-linked immunoassays and real-time PCR during the summer months in Northern Alberta, Canada. *J Clin Microbiol* 2011;49:4307–4310.

Cressey P, Lake R. Ranking Food Safety Risks: Development of NZFSA Policy 2004–2005. Client Report FW0563:23. 2005. Available at: [http://www.foodsafety.govt.nz/library/industry/Ranking\\_Food\\_Safety\\_Risks-Science\\_Research.pdf](http://www.foodsafety.govt.nz/library/industry/Ranking_Food_Safety_Risks-Science_Research.pdf), accessed on December 30, 2015.

David JM, Ravel A, Nesbitt A, Pintar K, Pollari F. Assessing multiple foodborne, waterborne and environmental exposures of healthy people to potential enteric pathogen sources: Effect of age, gender, season, and recall period. *Epidemiol Infect* 2014;142:28–39.

Davidson VJ, Ravel A, Nguyen TN, Fazil A, Ruzante JM. Food-specific attribution of selected gastrointestinal illnesses: Estimates from a Canadian expert elicitation survey. *Foodborne Pathog Dis* 2011;8:983–995.

Galanis E, Mak S, Otterstatter M, Taylor M, Zubel M, Takaro TK, Kuo M, Michel P. The association between campylobacteriosis, agriculture and drinking water: A case-case study in a region of British Columbia, Canada, 2005–2009. *Epidemiol Infect* 2014;142:2075–2084.

Greig JD, Ravel A. Analysis of foodborne outbreak data reported internationally for source attribution. *Int J Food Microbiol* 2009;130:77–87.

Havelaar AH, Galindo AV, Kurowicka D, Cooke RM. Attribution of foodborne pathogens using structured expert elicitation. *Foodborne Pathog Dis* 2008;5:649–659.

Health Canada. Food and Drug Regulations (C.R.C., c. 870). 2014. Available at: [http://laws-lois.justice.gc.ca/eng/regulations/C.R.C.%2C\\_c.\\_870/](http://laws-lois.justice.gc.ca/eng/regulations/C.R.C.%2C_c._870/), accessed September 25, 2014.

Hoffmann SA, Fischbeck PS, Krupnick AJ, McWilliams M. Attributing Foodborne Illnesses to Their Food Sources: Using Large Expert Panels to Capture Variability in Expert Judgment. Discussion Paper 06-17-REV. 2006. Available at: <http://www.rff.org/files/sharepoint/WorkImages/Download/RFF-DP-06-17-REV.pdf>, accessed on December 30, 2015.

Lam S, Sivaramalingam B, Gangodawilage H. *Cryptosporidium* outbreaks associated with swimming pools. *Environ Health Rev* 2014;57:3–8.

Murphy HM, Thomas MK, Schmidt PJ, Medeiros DT, McFadyen S, Pintar KDM. Estimating the burden of acute gastrointestinal illness due to *Giardia*, *Cryptosporidium*, *Campylobacter*, *E. coli*

- O157 and norovirus associated with private wells and small water systems in Canada. *Epidemiol Infect* 2015. doi:10.1017/S0950268815002071.
- Ortega YR, Sanchez R. Update on *Cyclospora cayentanensis*, a food-borne and waterborne parasite. *Clin Microbiol Rev* 2010;23:218–234.
- Painter JA, Ayers T, Woodruff R, Blanton E, Perez N, Hoekstra RM, Griffin PM, Braden C. Recipes for foodborne outbreaks: A scheme for categorizing and grouping implicated foods. *Foodborne Pathog Dis* 2009;6:1259–1264.
- Ravel A, Davidson VJ, Ruzante JM, Fazil A. Foodborne proportion of gastrointestinal illness: Estimates from a Canadian expert elicitation survey. *Foodborne Pathog Dis* 2010;7:1463–1472.
- Ravel A, Greig J, Tinga C, Todd E, Campbell G, Cassidy M, Marshall B, Pollari F. Exploring historical Canadian foodborne outbreak data sets for human illness attribution. *J Food Prot* 2009;72:1963–1976.
- Scallan E, Hoekstra R, Angulo F, Tauxe R, Widdowson M, Roy S, Jones J, Griffin P. Foodborne Illness acquired in the United States—Major pathogens. *Emerg Infect Dis* 2011;17:7–15.
- Snel SJ, Baker MG, Kamalesh V, French N, Learmonth J. A tale of two parasites: The comparative epidemiology of cryptosporidiosis and giardiasis. *Epidemiol Infect* 2009;137:1641–1650.
- Tantillo GM, Fontanarosa M, Di Pinto A, Musti M. Updated perspectives on emerging vibrios associated with human infections. *Lett Appl Microbiol* 2004;39:117–126.
- Thomas MK, Murray R, Flockhart L, Pintar K, Pollari F, Fazil A, Nesbitt A, Marshall B. Estimates of the burden of foodborne illness in Canada for 30 specified pathogens and unspecified agents, circa 2006. *Foodborne Pathog Dis* 2013;10:639–648.
- Uhlmann S, Galanis E, Takaro T, Mak S, Gustafson L, Embree G, Bellack N, Corbett K, Isaac-Renton J. Where's the pump? Associating sporadic enteric disease with drinking water using a geographic information system, in British Columbia, Canada, 1996–2005. *J Water Health* 2009;7:692–698.
- Vally H, Glass K, Ford L, Hall G, Kirk MD, Shadbolt C, Veitch M, Fullerton KE, Mustro J, Becker N. Proportion of illness acquired by foodborne transmission for nine enteric pathogens in Australia: An expert elicitation. *Foodborne Pathog Dis* 2014;11:727–733.
- [WHO] World Health Organization. Guidelines for drinking-water quality. 2011. Available at: [http://www.who.int/water\\_sanitation\\_health/publications/2011/dwq\\_guidelines/en/](http://www.who.int/water_sanitation_health/publications/2011/dwq_guidelines/en/), accessed on December 30, 2015.
- Yoder JS, Gargano JW, Wallace RM, Beach MJ. Giardiasis surveillance—United States, 2009–2010. *MMWR* 2012a;7:13–23.
- Yoder JS, Wallace RM, Collier SA, Beach MJ, Hlavsa MC. Cryptosporidiosis surveillance—United States, 2009–2010. *MMWR* 2012b;61:1–12.

Address correspondence to:  
Katarina D.M. Pintar, MSc, PhD  
Centre for Foodborne, Environmental,  
and Zoonotic Infectious Diseases  
Public Health Agency of Canada  
130 Colonnade Rd.  
Ottawa, ON K1A 0K9, Canada  
E-mail: katarina.pintar@phac-aspc.gc.ca