

RESEARCH ARTICLE

Characterization of the pathogenome and phylogenomic classification of enteropathogenic *Escherichia coli* of the O157:non-H7 serotypes

Fatemeh Sanjar¹, Brigida Rusconi¹, Tracy H. Hazen², Sara S.K. Koenig¹, Mark K. Mammel³, Peter C.H. Feng⁴, David A. Rasko² and Mark Eppinger^{1,*}

¹Department of Biology & South Texas Center for Emerging Infectious Diseases, University of Texas at San Antonio, San Antonio, TX 78249, USA, ²Institute for Genome Sciences (IGS), University of Maryland, School of Medicine, Department of Microbiology and Immunology, Baltimore, MD 21021, USA, ³Center for Food Safety and Applied Nutrition, U.S. Food and Drug Administration, Laurel, MD 20708, USA and ⁴Division of Microbiology, U.S. Food and Drug Administration, College Park, MD 20740, USA

*Corresponding author: Department of Biology & South Texas Center for Emerging Infectious Diseases, University of Texas at San Antonio, San Antonio, TX 78249, USA. Tel: +210-458-6276; Email: mark.eppinger@utsa.edu

One sentence summary: Whole-genome sequencing and phylogenomic analyses of 10 enteropathogenic *Escherichia coli* strains of the O157:non-H7 serotypes provide evidence for their non-monophyletic evolutionary origin and identify a heterogeneous virulence complement.

Editor: Nicholas Thomson

ABSTRACT

Escherichia coli of the O157 serogroup are comprised of a diverse collection of more than 100 O157:non-H7 serotypes that are found in the environment, animal reservoir and infected patients and some have been linked to severe outbreaks of human disease. Among these, the enteropathogenic *E. coli* O157:non-H7 serotypes carry virulence factors that are hallmarks of enterohemorrhagic *E. coli*, such as causing attaching and effacing lesions during human gastrointestinal tract infections. Given the shared virulence gene pool between O157:H7 and O157:non-H7 serotypes, our objective was to examine the prevalence of virulence traits of O157:non-H7 serotypes within and across their H-serotype and when compared to other *E. coli* pathovars. We sequenced six O157:non-H7 genomes complemented by four genomes from public repositories in an effort to determine their virulence state and genetic relatedness to the highly pathogenic enterohemorrhagic O157:H7 lineage and its ancestral O55:H7 serotype. Whole-genome-based phylogenomic analysis and molecular typing is indicative of a non-monophyletic origin of the heterogeneous O157:non-H7 serotypes that are only distantly related to the O157:H7 serotype. The availability of multiple genomes enables robust phylogenomic placement of these strains into their evolutionary context, and the assessment of the pathogenic potential of the O157:non-H7 strains in causing human disease.

Keywords: enteropathogenic *E. coli* (EPEC); O157:non-H7; pathogenome evolution; genotyping

INTRODUCTION

The enteropathogenic *Escherichia coli* (EPEC) belong to the large diarrheagenic *E. coli* (DAEC) pathovar (Nataro and Kaper 1998)

and are a major cause of infantile diarrhea and the leading cause of morbidity and mortality in infants in developing countries (Trabulsi, Keller and Tardelli Gomes 2002; Afset, Bergh and Bevanger 2003; Gomes et al. 2004; Kaper, Nataro and Mobley 2004;

Received: 29 January 2015; Accepted: 1 May 2015

© FEMS 2015. All rights reserved. For permissions, please e-mail: journals.permissions@oup.com

Alikhani, Mirsalehian and Aslani 2006; Behiry et al. 2011). The EPEC O157 serogroup is comprised of more than 100 O157:non-H7 serotypes that have been isolated from different sources including animal reservoirs and infected patients (Stephan et al. 2004; Wani et al. 2006; Feng et al. 2010); additionally, some of the O157:non-H7 strains have been associated with outbreaks of human disease (Makino et al. 1999; Yatsuyanagi et al. 2002; Feng et al. 2010). An array of virulence determinants have been identified in O157:non-H7 strains that are also present in enterohemorrhagic *E. coli* (EHEC) strains, including the highly pathogenic O157:H7 serotype (Caprioli et al. 2005; Eppinger et al. 2011a,b; Sadiq et al. 2014). One virulence feature of O157:non-H7 strains shared with O157:H7 is the locus for enterocyte effacement (LEE) pathogenicity-associated island (PAI), which mediates the formation of attaching and effacing lesions (A/E) during colonization of the gastrointestinal (GI) tract (Jerse, Gicquelais and Kaper 1991; Phillips and Frankel 2000; Blanco et al. 2006; Coburn, Sekirov and Finlay 2007). The EPEC serotypes are distinguished from Shiga toxin-producing *E. coli* (STEC), such as EHEC O157:H7, by the lack of *stx*-encoding bacteriophages (*stx1* and/or *stx2*) (Cornick et al. 2000; Eklund, Leino and Siitonen 2002). The EPEC serotypes can be further classified into typical (tEPEC) and atypical EPEC (aEPEC) strains by the presence or absence of the enteropathogenic adherence factor (EAF) plasmid-borne *bfpA* gene, which encodes bundlin, a major structural subunit of the bundle-forming pilus (BFP), used by tEPEC for aggregation and bundle formation during colonization (Blank et al. 2000). The production of BFP protein leads to localized adherence and induction of host epithelium cell death (Melo et al. 2005). Another virulence determinant of tEPEC is the EAF plasmid-encoded regulator, *perA*, which is involved in autoaggregation and elevated expression of BFP, intimin and Tir proteins during infection (Okeke et al. 2001; Iida et al. 2010).

The aim of this study was to elucidate the genomic characteristics and plasticity among the different O157:non-H7 serotypes through comparative genomics of six in-house sequenced and four publicly available O157:non-H7 strains as well as a selection of O157:H7, O157:H(-) and O55:H7 strains sequenced in this study or retrieved from public repositories. Comparative genome analysis within each H-serogroup, between different H-serogroups, as well as when compared to other aEPEC (O55:H7) and EHEC (O157:H7/H(-)) pathovar strains, identified common and unique traits in the pathogenome evolution of O157:non-H7 strains with respect to the aforementioned serotypes. Here we observed that while EHEC O157:H7 and O157:H(-) share certain virulence characteristics with O157:non-H7 serotypes, whole-genome sequence typing approaches, including multilocus sequence typing (MLST) or *eae*-genotyping, could not establish close genetic relatedness of these O157:non-H7 lineages with EHEC or other EPEC serotypes. These findings are critical in developing a refined phylogenomic framework and assessment for pathogenic potential of the extant O157:non-H7 strains, which are non-monophyletic.

METHODS

Escherichia coli strains used in this study

To capture the genomic plasticity of the O157:non-H7 serotypes, we selected 10 epidemiologically diverse O157:non-H7 strains that were isolated from water, infected patients and processed meat (Table 1) (Feng et al. 2010; Hazen et al. 2013a; Svab et al. 2013). Strains of the O55:H7, O157:H(-) and O157:H7 serotypes were included to study their genetic relatedness to

the O157:non-H7 serotypes (Table S1, Supporting Information) (Hayashi et al. 2001; Perna et al. 2001; Eppinger et al. 2011b; Rump et al. 2011; Hazen et al. 2012; Kyle et al. 2012). Representative O157:H7 strains for each of the nine phylogenetic clades (Manning et al. 2008) were used as reference to determine the relationship of O157:non-H7 to the extant O157:H7 strains.

Whole-genome sequencing and phylogeny

To study the phylogenetic relatedness of diverse O157:non-H7 strains to other *E. coli* pathotypes, we have sequenced the genomes of 14 *E. coli* strains of which six belong to O157:non-H7, one to O55:H7, four to O157:H7 and three to O157:H(-) (Table S1, Supporting Information). Genomes were subjected to Illumina sequencing using paired-end libraries with 300-bp inserts on the HiSeq2000 platform. Draft genomes were assembled using Velvet assembler (Zerbino and Birney 2008; Zerbino 2010), and annotated with the IGS Annotation Engine and Mantea for structural and functional genome annotation (Galens et al. 2011). Accession numbers are listed in Table S1 (Supporting Information). Bacterial genomes were aligned using Mugsy (Angiuoli and Salzberg 2011), and the phylogenetic tree was constructed using the maximum-likelihood method in RAxML v8.1 (-f an option) with 100 bootstraps (Stamatakis 2014), and the evolutionary relationship hypothesis was visualized using EvolView (Zhang et al. 2012). For strains G5101 (SRX702352), 493-89 (SRX702359), USDA5905 (SRX702362) and H2687 (SRX702361), the original reads deposited by Rump et al. were retrieved from the short reads archives (SRA) and *de novo* assembled with Geneious assembler v.7.1.7, (Kearse et al. 2012). Assembled contigs were queried for the presence of *stx1* and *stx2* genes using VirulenceFinder 1.2 (Joensen et al. 2014).

Molecular genotyping and subtyping

We examined the status of the β -glucuronidase enzyme *in silico* for β -glucuronidase activity (GUD) using BLASTN focusing on position +93 *uidA* gene (T:G allele) (Altschul et al. 1990; Martins et al. 1993) and functionally examined (when applicable) the ability to ferment sorbitol (SOR) on Sorbitol MacConkey agar (Oxoid, UK) (Farmer and Davis 1985). The *eae*, *perA* and *bfpA* positive O157:non-H7 strains were further subtyped into designated allele profiles (Adu-Bobie et al. 1998; Blank et al. 2000; Blanco et al. 2006; Lacher et al. 2007; Contreras et al. 2010). The Achtman MLST scheme (Wirth et al. 2006) was used for *in silico* polymorphism profiling of *E. coli* based on seven housekeeping genes.

RESULTS AND DISCUSSION

The O157:non-H7 strains selected for this study originated from diverse geographical locations and different ecological niches, such as human patients, processed meat, and water (Table S1, Supporting Information). Whole-genome screening of the virulence state and molecular subtyping of known EPEC- and EHEC-associated virulence factors revealed a highly diverse and heterogeneous genomic makeup among the O157:non-H7 serogroups studied (Table 1). Furthermore, the phylogenetic relationship of these O157:non-H7 serotypes to other well-studied pathogenic serotypes, such as EPEC (O55:H7) and EHEC (O157:H7/H(-)), was analyzed (Fig. S1, Supporting Information). Availability of high-quality whole-genome sequences enables the determination of the virulence gene state and phylogenomic grouping according to established genotypic classification methods using both *in silico* and experimental assays (Farmer and

Table 1. Genomic characteristics of EPEC O157:non-H7 strains.

| Strain | Serotype | Source | Origin | EPEC | MLST ^{a)} | GUD | SOR | <i>eae</i> | <i>tir</i> | <i>bfpA</i> | <i>perA</i> |
|-----------|----------|--------|-----------|------|--------------------|-----|-----|----------------------|------------|---------------|---------------|
| ARS4.2123 | O157:H16 | Water | USA | T | 79 | + | + | +(ϵ) | + | +(β 1) | +(β 2) |
| RN587/1 | O157:H8 | Human | Brazil | T | 725 | + | ND | +(α) | + | +(β 6) | +(β 2) |
| C844-97 | O157:H45 | Human | Japan | A | 725 | + | ND | +(α) | + | - | - |
| C639-08 | O157:H45 | Human | Denmark | A | 725 | + | ND | +(α) | + | - | - |
| TW00353 | O157:H45 | Human | USA | A | 2 | + | + | +(ϵ) | + | - | - |
| TW15901 | O157:H16 | Meat | France | A | 10 | + | + | +(ϵ) | + | - | - |
| N1 | O157:H29 | Meat | Unknown | NA | 2 | + | + | - | - | - | - |
| T22 | O157:H43 | Human | Hungary | NA | 155 | + | ND | - | - | - | - |
| 3006 | O157:H16 | Human | USA | NA | 6 | + | - | - | - | - | - |
| 7798 | O157:H39 | Human | Argentina | A | 2 | + | - | +(κ/δ) | + | - | - |

O157:non-H7 strains were genotyped for prevalence and plasticity in established EHEC- and EPEC-associated phylogenetic and virulence markers (Farmer and Davis 1985; Adu-Bobie et al. 1998; Blank et al. 2000; Okeke et al. 2001; Yoshitomi et al. 2003; Contreras et al. 2010). MLST^{a)}, Achtman ST7 MLST approach (Wirth et al. 2006); NA, Not applicable; SOR, sorbitol fermentation; GUD, β -glucourinidase activity; T, typical EPEC; A, atypical EPEC; ND, No data available.

Davis 1985; Yoshitomi, Jinneman and Weagant 2003; Wirth et al. 2006; Manning et al. 2008; Contreras et al. 2010).

Whole-genome phylogeny

The phylogenetic tree confirmed the ancestral status of aEPEC O55:H7 to EHEC serotypes (O157:H7 and O157:H(-)) and exhibited a more distant relationship to the cluster of O157:non-H7 serotypes (Fig. S1, Supporting Information) (Trabulsi, Keller and Tardelli Gomes 2002; Zhou et al. 2010; Eppinger et al. 2011b; Kyle et al. 2012). In agreement with a limited loci-based study by Feng et al. (2010), our whole-genome analysis demonstrated that the O157:non-H7 serotypes are not as closely related to the aEPEC O55:H7 when compared to the descendant EHEC O157:H7 and O157:H(-) serotypes (Feng et al. 2010; Zhou et al. 2010; Kyle et al. 2012). In contrast to the genetically highly homogenous population structure of EHEC O157:H7 and O157:H(-) serotypes (Manning et al. 2008; Eppinger et al. 2011b), whole-genome analysis revealed a heterogeneous genome composition among the O157:non-H7 group of serotypes featuring the same H-serogroup (e.g. O157:H16) (Yatsuyanagi et al. 2002; Feng et al. 2010; Bugarel et al. 2011; Eppinger et al. 2011b). As evident in the tree topology (Figs 1 and S1, Supporting Information), the clustering of the different O157:non-H7 strains did not correlate with the H-serotype, as exemplified by the O157:H16 and O157:H45 strains that are scattered within the O157:non-H7 cluster (Fig. 1).

Using EPEC-associated traits and molecular genotyping, we identified two O157:non-H7 strains as tEPEC (*perA/bfpA* positive), and the remaining eight strains were classified as aEPEC strains (Table 1). Seven of the studied O157:non-H7 strains were positive for the α , β or ϵ *eae* allele, but none carried the *eae*- γ allele, which is commonly associated with the EHEC O157:H7 serotype (Feng et al. 2010) (Table 1). Overall, these O157:non-H7 strains showed a high degree of variability in their virulence gene complement, which is consistent with the findings reported for other EPEC serotypes (Gomes et al. 2004; Hazen et al. 2013b). As shown in Table 1, seven of the O157:non-H7 strains exhibited an array of EPEC- and/or EHEC-associated virulence determinants with the notable exception of strains N1 (O157:H29), T22 (O157:H43) and 3006 (O157:H16), which lacked these loci and were not classified as tEPEC or aEPEC utilizing the currently established typing schemes (Adu-Bobie et al. 1998; Nataro and Kaper 1998; Blank et al. 2000; Okeke et al. 2001; Dulguer et al. 2003; Contreras et al. 2010), but might originate from an STEC that lost its *stx*-encoding bacteriophage. Since one of these strains was isolated from a human case with GI symptoms, its pathogenic potential was likely due to carriage of other virulence loci or the result of a coinfection

with other pathogenic *E. coli* strains (Dulguer et al. 2003). The observed combination of virulence determinants likely occurred due to lateral acquisition and cotransfer of genes packaged in virulence plasmids (e.g. EAF-encoded *perA* and *bfpA*) (Jerse, Gicquelais and Kaper 1991; Iida et al. 2010) or PAI (e.g. LEE-encoded *eae/tir*) (Bono et al. 2007; Flockhart et al. 2012). Overall, the diversity observed in these O157:non-H7 genomes can be considered significant.

MLST and phylogenetic placement

The genetically homogenous nature of the *E. coli* O157:H7 serotype was evidenced by a common Achtman MLST profile of sequence type 11 (ST11) (Wirth et al. 2006; Eppinger et al. 2011b). The non-motile O157:H(-) subgroup (clades 9 and 7) showed the same MLST profile with the exception of strain H2687 (ST587) (Fig. 1). All analyzed non-motile O157:H(-) strains belong to clade 9 and are phylogenetically clustered with the notable exception of the clade 7 strain TW14301 (Fig. 1). We screened for alterations in motility loci (*flhC*, *flhD* or *fliC*) that have been previously associated with non-motile O157:H(-) strains (Dobbin et al. 2006). Clade 9 O157:H(-) strains exhibit a 12-bp deletion in *flhC*, consistent with previous findings (Monday, Minnich and Feng 2004), while TW14301 carried the wild-type form of the respective motility genes. Our findings point to loss of motility through independent events in EHEC evolution in the non-motile O157:H(-) serotype (Wang et al. 2003; Dobbin et al. 2006; Kyle et al. 2012). MLST profiling of O55:H7 strains revealed two distinct MLST patterns of ST11 and ST335 (Wirth et al. 2006) based on the Achtman MLST scheme. As shown in Fig. 1, we found instances where MLST profiles of the O157:non-H7 did not corroborate with the corresponding phylogenomic placement. The MLST profile of O157:non-H7 strains (Wirth et al. 2006) was inconclusive in delineating their evolutionary relationship when compared to the whole-genome-based phylogenomic hypothesis (Fig. 1). For example, strains N1 (O157:H29) and TW00353 (O157:H45) or RN587/1 (O157:H8) and C844-97 (O157:H45) feature identical MLST profiles, yet differ in their H-serotype and/or *eae*/*bfpA*-genotype pattern (Table 1), as well as their placement delineated in the evolutionary tree (Fig. S1, Supporting Information).

Carriage of *stx*-encoding bacteriophages

All analyzed O157:non-H7 strains lack the *stx*-encoding bacteriophages (*stx1* and *stx2*), which is a virulence hallmark of STEC/EHEC strains (Fig. 1). For some of the strains, the presence




Figure 1. Whole-genome-based phylogeny of O157:non-H7 EPEC strains and select EPEC and EHEC strains. The tree topology demonstrates that O157:non-H7 serogroups form a heterogeneous and non-monophyletic group, as evident from the phylogenetic placement of strains belonging to the same H-serogroup (e.g. H16 and H45). Inferred MLST profiles do not agree with the phylogenetic placement delineated from whole genome alignment. Our data confirm that O157:non-H7 strains are not closely related to the EHEC O157:H7/H(-) lineage or its ancestral aEPEC O55:H7 serotype. For phylogenetic and evolutionary distances, refer to Fig. S1 (Supporting Information).

of *stx1* and *stx2* genes had to be assessed from a *de novo* assembly from the SRA files, as the published contigs were *stx* negative. However, among the O55:H7 strains we identified two *stx2*-positive strains, 5905 and USDA5905, which carried a lambda-like *stx2d*-encoding bacteriophage at the *yecE* locus. This phage is closely related to the enterobacteria phage BP-4795 found in the EHEC strain RM13516 of the O145:H28 serotype (De Schrijver et al. 2008; Buvens et al. 2011). This particular serotype is related to serotypes O157 and O55 (EHEC1/EPEC1 lineage) (Cooper et al. 2014). We note here that several strains of the O55:H7 serotype

have been reported as *stx*-positive (Feng et al. 1998a,b; Kyle et al. 2012); however, to determine the distinct evolutionary relationship of *stx*-positive O55:H7 strains compared to EHEC O157:H7 and O157:H(-) serotypes, further phylogenetic studies on larger genomic data sets are needed.

Metabolism

All of the EPEC O157:non-H7 and O55:H7 strains carried the T allele at position +93 *uidA* and were functionally GUD

positive, while all of the O157:H7 and O157:H(-) serotypes carried the G allele and lacked GUD metabolic activity. In analogy to a subset of O157:H7 strains, two of the O157:non-H7 strains (e.g. 3006 and 7798) also did not have the ability to ferment sorbitol (Table 1) (Kyle et al. 2012). Although these two metabolic features are closely associated with the highly pathogenic EHEC O157:H7 lineage, while basal EHEC and ancestral O55:H7 strains are GUD/SOR positive, there are atypical strains of O157:H7 that are GUD positive and the O157:H(-) group is positive for both sorbitol and GUD (Feng et al. 2007).

CONCLUSIONS

The availability of high-quality annotated draft genome sequences for EPEC O157:non-H7 strains provides a critical resource in further understanding the pathogenome evolution and relatedness among different *E. coli* pathotypes (Dugan et al. 2014; Franz et al. 2014). Our analysis revealed the heterogeneous nature of O157:non-H7 serotypes by studying their genomic inventory and plasticity in the core genome and laterally acquired regions. Overall, the genotypic data for majority of the O157:non-H7 strains we studied (Table 1) did not corroborate with the phylogenomic clustering, as delineated from the whole-genome analysis (Fig. 1), and as suggested by previous publications (Steyert et al. 2012; Hazen et al. 2013b). Even among the same H-serotype strains (e.g. H45 or H16), we observed a non-monophyletic origin that accounts for their diverse genomic content (Fig. 1). Contrary to previous reports of a close relationship of aEPEC and STEC strains in terms of genetic characteristics (Trabulsi, Keller and Tardelli Gomes 2002), we observed a more distant evolutionary relationship between the different tEPEC and aEPEC O157:non-H7 serotypes and the EHEC O157:H7 serotype, as compared to the EPEC O55:H7 (Fig. S1, Supporting Information). In general, aEPEC strains are reported to carry a more heterogeneous virulence profile in comparison to tEPEC strains, and their degree of pathogenicity in the absence of the EAF plasmid remains more elusive (Levine et al. 1978; Trabulsi, Keller and Tardelli Gomes 2002). Here we report that even strains within the same aEPEC O157:non-H7 serotype remain genotypically distinct, and that the reported variability could be attributed to polymorphisms within MLST and established virulence markers. On the other hand, the aEPEC O55:H7 exhibits a more homogenous genotype with few exceptions (Fig. 1), which highlights the degree of diversity found within and among different EPEC serotypes. Further studies are needed to elucidate the evolutionary origin and emergence of EPEC O157:non-H7, which appear to be only distantly related to aEPEC O55:H7 and the highly pathogenic EHEC O157:H7 and H(-) serotypes.

SUPPLEMENTARY DATA

Supplementary data is available at FEMSPD online.

FUNDING

This work received support from the South Texas Center of Emerging Infectious Diseases (STCEID), Department of Biology and Computational System Biology Core at the University of Texas at San Antonio, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Department of Health and Human Services under contract HHSN272200900007C, the High Performance Computing Center/Computational Bioinformatics Initiative (HPC/CBI) under

contract 2G12RR013646-12. This project is supported by the Army Research Office of the Department of Defense under Contract No. W911NF-11-1-0136. FS is supported by the South Texas Center for Emerging Infectious Diseases (STCEID) and an University Teaching Fellowship (UTF). BR is supported by the Swiss National Science Foundation (SNSF) Early.Postdoc.Mobility Fellowship (P2LAP3-151770).

Conflict of interest. None declared.

REFERENCES

- Adu-Bobie J, Frankel G, Bain C, et al. Detection of intimins alpha, beta, gamma, and delta, four intimin derivatives expressed by attaching and effacing microbial pathogens. *J Clin Microbiol* 1998;36:662–8.
- Afset JE, Bergh K, Bevanger L. High prevalence of atypical enteropathogenic *Escherichia coli* (EPEC) in Norwegian children with diarrhoea. *J Med Microbiol* 2003;52:1015–9.
- Alikhani MY, Mirsalehian A, Aslani MM. Detection of typical and atypical enteropathogenic *Escherichia coli* (EPEC) in Iranian children with and without diarrhoea. *J Med Microbiol* 2006;55:1159–63.
- Altschul SF, Gish W, Miller W, et al. Basic local alignment search tool. *J Mol Biol* 1990;215:403–10.
- Angiuoli SV, Salzberg SL. Mugsy: fast multiple alignment of closely related whole genomes. *Bioinformatics* 2011;27:334–42.
- Behiry IK, Abada EA, Ahmed EA, et al. Enteropathogenic *Escherichia coli* associated with diarrhea in children in Cairo, Egypt. *ScientificWorldJournal* 2011;11:2613–9.
- Blanco M, Blanco JE, Dahbi G, et al. Typing of intimin (*eae*) genes from enteropathogenic *Escherichia coli* (EPEC) isolated from children with diarrhoea in Montevideo, Uruguay: identification of two novel intimin variants (*muB* and *xiR/beta2B*). *J Med Microbiol* 2006;55:1165–74.
- Blank TE, Zhong H, Bell AL, et al. Molecular variation among type IV pilin (*bfpA*) genes from diverse enteropathogenic *Escherichia coli* strains. *Infect Immun* 2000;68:7028–38.
- Bono JL, Keen JE, Clawson ML, et al. Association of *Escherichia coli* O157:H7 *tir* polymorphisms with human infection. *BMC Infect Dis* 2007;7:98.
- Bugarel M, Martin A, Fach P, et al. Virulence gene profiling of enterohemorrhagic (EHEC) and enteropathogenic (EPEC) *Escherichia coli* strains: a basis for molecular risk assessment of typical and atypical EPEC strains. *BMC Microbiol* 2011;11:142.
- Buvens G, Posse B, De Schrijver K, et al. Virulence profiling and quantification of verocytotoxin-producing *Escherichia coli* O145:H28 and O26:H11 isolated during an ice cream-related hemolytic uremic syndrome outbreak. *Foodborne Pathog Dis* 2011;8:421–6.
- Caprioli A, Morabito S, Brugere H, et al. Enterohaemorrhagic *Escherichia coli*: emerging issues on virulence and modes of transmission. *Vet Res* 2005;36:289–311.
- Coburn B, Sekirov I, Finlay BB. Type III secretion systems and disease. *Clin Microbiol Rev* 2007;20:535–49.
- Contreras CA, Ochoa TJ, Lacher DW, et al. Allelic variability of critical virulence genes (*eae*, *bfpA* and *perA*) in typical and atypical enteropathogenic *Escherichia coli* in Peruvian children. *J Med Microbiol* 2010;59:25–31.
- Cooper KK, Mandrell RE, Louie JW, et al. Comparative genomics of enterohemorrhagic *Escherichia coli* O145:H28 demonstrates a common evolutionary lineage with *Escherichia coli* O157:H7. *BMC Genomics* 2014;15:17.

- Cornick NA, Matise I, Samuel JE, et al. Shiga toxin-producing *Escherichia coli* infection: temporal and quantitative relationships among colonization, toxin production, and systemic disease. *J Infect Dis* 2000;**181**:242–51.
- De Schrijver K, Buvens G, Posse B, et al. Outbreak of verocytotoxin-producing *E. coli* O145 and O26 infections associated with the consumption of ice cream produced at a farm, Belgium, 2007. *Euro Surveill* 2008;**13**:pii:8041.
- Dobbin HS, Hovde CJ, Williams CJ, et al. The *Escherichia coli* O157 flagellar regulatory gene *flhC* and not the flagellin gene *fliC* impacts colonization of cattle. *Infect Immun* 2006;**74**:2894–905.
- Dugan VG, Emrich SJ, Giraldo-Calderon GI, et al. Standardized metadata for human pathogen/vector genomic sequences. *PLoS One* 2014;**9**:e99979.
- Dulguer MV, Fabbriotti SH, Bando SY, et al. Atypical enteropathogenic *Escherichia coli* strains: phenotypic and genetic profiling reveals a strong association between enteroaggregative *E. coli* heat-stable enterotoxin and diarrhea. *J Infect Dis* 2003;**188**:1685–94.
- Eklund M, Leino K, Siitonen A. Clinical *Escherichia coli* strains carrying stx genes: stx variants and stx-positive virulence profiles. *J Clin Microbiol* 2002;**40**:4585–93.
- Eppinger M, Mammel MK, Leclerc JE, et al. Genome signatures of *Escherichia coli* O157:H7 isolates from the bovine host reservoir. *Appl Environ Microb* 2011a;**77**:2916–25.
- Eppinger M, Mammel MK, Leclerc JE, et al. Genomic anatomy of *Escherichia coli* O157:H7 outbreaks. *P Natl Acad Sci USA* 2011b;**108**:20142–7.
- Farmer JJ, 3rd, Davis BR. H7 antiserum-sorbitol fermentation medium: a single tube screening medium for detecting *Escherichia coli* O157:H7 associated with hemorrhagic colitis. *J Clin Microbiol* 1985;**22**:620–5.
- Feng P, Lampel KA, Karch H, et al. Genotypic and phenotypic changes in the emergence of *Escherichia coli* O157:H7. *J Infect Dis* 1998a;**177**:1750–3.
- Feng P, Sandlin RC, Park CH, et al. Identification of a rough strain of *Escherichia coli* O157:H7 that produces no detectable O157 antigen. *J Clin Microbiol* 1998b;**36**:2339–41.
- Feng PC, Keys C, Lacher D, et al. Prevalence, characterization and clonal analysis of *Escherichia coli* O157: non-H7 serotypes that carry *eae* alleles. *FEMS Microbiol Lett* 2010;**308**:62–7.
- Feng PC, Monday SR, Lacher DW, et al. Genetic diversity among clonal lineages within *Escherichia coli* O157:H7 stepwise evolutionary model. *Emerg Infect Dis* 2007;**13**:1701–6.
- Flockhart AF, Tree JJ, Xu X, et al. Identification of a novel prophage regulator in *Escherichia coli* controlling the expression of type III secretion. *Mol Microbiol* 2012;**83**:208–23.
- Franz E, Delaquis P, Morabito S, et al. Exploiting the explosion of information associated with whole genome sequencing to tackle Shiga toxin-producing *Escherichia coli* (STEC) in global food production systems. *Int J Food Microbiol* 2014;**187C**:57–72.
- Galens K, Orvis J, Daugherty S, et al. The IGS standard operating procedure for automated prokaryotic annotation. *Stand Genomic Sci* 2011;**4**:244–51.
- Gomes TA, Irino K, Girao DM, et al. Emerging enteropathogenic *Escherichia coli* strains? *Emerg Infect Dis* 2004;**10**:1851–5.
- Hayashi T, Makino K, Ohnishi M, et al. Complete genome sequence of enterohemorrhagic *Escherichia coli* O157:H7 and genomic comparison with a laboratory strain K-12. *DNA Res* 2001;**8**:11–22.
- Hazen TH, Sahl JW, Fraser CM, et al. Draft genome sequences of three O157 enteropathogenic *Escherichia coli* isolates. *Genome Announc* 2013a;**193**:2058–9.
- Hazen TH, Sahl JW, Fraser CM, et al. Refining the pathovar paradigm via phylogenomics of the attaching and effacing *Escherichia coli*. *P Natl Acad Sci USA* 2013b;**110**:12810–5.
- Hazen TH, Sahl JW, Redman JC, et al. Draft genome sequences of the diarrheagenic *Escherichia coli* collection. *J Bacteriol* 2012;**194**:3026–7.
- Iida M, Okamura N, Yamazaki M, et al. Classification of *perA* sequences and their correlation with autoaggregation in typical enteropathogenic *Escherichia coli* isolates collected in Japan and Thailand. *Microbiol Immunol* 2010;**54**:184–95.
- Jerse AE, Gicquelais KG, Kaper JB. Plasmid and chromosomal elements involved in the pathogenesis of attaching and effacing *Escherichia coli*. *Infect Immun* 1991;**59**:3869–75.
- Joensen KG, Scheutz F, Lund O, et al. Real-time whole-genome sequencing for routine typing, surveillance, and outbreak detection of verotoxigenic *Escherichia coli*. *J Clin Microbiol* 2014;**52**:1501–10.
- Kaper JB, Nataro JP, Mobley HL. Pathogenic *Escherichia coli*. *Nat Rev Microbiol* 2004;**2**:123–40.
- Kearse M, Moir R, Wilson A, et al. Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 2012;**28**:1647–9.
- Kyle JL, Cummings CA, Parker CT, et al. *Escherichia coli* serotype O55:H7 diversity supports parallel acquisition of bacteriophage at Shiga toxin phage insertion sites during evolution of the O157:H7 lineage. *J Bacteriol* 2012;**194**:1885–96.
- Lacher DW, Steinsland H, Blank TE, et al. Molecular evolution of typical enteropathogenic *Escherichia coli*: clonal analysis by multilocus sequence typing and virulence gene allelic profiling. *J Bacteriol* 2007;**189**:342–50.
- Levine MM, Bergquist EJ, Nalin DR, et al. *Escherichia coli* strains that cause diarrhoea but do not produce heat-labile or heat-stable enterotoxins and are non-invasive. *Lancet* 1978;**1**:1119–22.
- Makino S, Asakura H, Shirahata T, et al. Molecular epidemiological study of a mass outbreak caused by enteropathogenic *Escherichia coli* O157:H45. *Microbiol Immunol* 1999;**43**:381–4.
- Manning SD, Motiwala AS, Springman AC, et al. Variation in virulence among clades of *Escherichia coli* O157:H7 associated with disease outbreaks. *P Natl Acad Sci USA* 2008;**105**:4868–73.
- Martins MT, Rivera IG, Clark DL, et al. Distribution of *uidA* gene sequences in *Escherichia coli* isolates in water sources and comparison with the expression of beta-glucuronidase activity in 4-methylumbelliferyl-beta-D-glucuronide media. *Appl Environ Microb* 1993;**59**:2271–6.
- Melo AR, Lasunskaja EB, de Almeida CM, et al. Expression of the virulence factor, *BfpA*, by enteropathogenic *Escherichia coli* is essential for apoptosis signalling but not for NF-kappaB activation in host cells. *Scand J Immunol* 2005;**61**:511–9.
- Monday SR, Minnich SA, Feng PC. A 12-base-pair deletion in the flagellar master control gene *flhC* causes nonmotility of the pathogenic German sorbitol-fermenting *Escherichia coli* O157:H- strains. *J Bacteriol* 2004;**186**:2319–27.
- Nataro JP, Kaper JB. Diarrheagenic *Escherichia coli*. *Clin Microbiol Rev* 1998;**11**:142–201.
- Okeke IN, Borneman JA, Shin S, et al. Comparative sequence analysis of the plasmid-encoded regulator of enteropathogenic *Escherichia coli* Strains. *Infect Immun* 2001;**69**:5553–64.
- Perna NT, Plunkett G, 3rd, Burland V, et al. Genome sequence of enterohaemorrhagic *Escherichia coli* O157:H7. *Nature* 2001;**409**:529–33.

- Phillips AD, Frankel G. Intimin-mediated tissue specificity in enteropathogenic *Escherichia coli* interaction with human intestinal organ cultures. *J Infect Dis* 2000;**181**:1496–500.
- Rump IV, Strain EA, Cao G, et al. Draft genome sequences of six *Escherichia coli* isolates from the stepwise model of emergence of *Escherichia coli* O157:H7. *J Bacteriol* 2011;**193**:2058–9.
- Sadiq SM, Hazen TH, Rasko DA, et al. EHEC genomics: past, present, and future. *Microbiol Spectrum* 2014;**2**:1–13.
- Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 2014;**30**:1312–3.
- Stephan R, Borel N, Zweifel C, et al. First isolation and further characterization of enteropathogenic *Escherichia coli* (EPEC) O157:H45 strains from cattle. *BMC Microbiol* 2004;**4**:10.
- Steyert SR, Sahl JW, Fraser CM, et al. Comparative genomics and stx phage characterization of LEE-negative Shiga toxin-producing *Escherichia coli*. *Front Cell Infect Microbiol* 2012;**2**:133.
- Svab D, Horvath B, Szucs A, et al. Draft genome sequence of an *Escherichia coli* O157:H43 strain isolated from cattle. *Genome Announc* 2013;**1**:e00263-13.
- Trabulsi LR, Keller R, Tardelli Gomes TA. Typical and atypical enteropathogenic *Escherichia coli*. *Emerg Infect Dis* 2002;**8**:508–13.
- Wang L, Rothmund D, Curd H, et al. Species-wide variation in the *Escherichia coli* flagellin (H-antigen) gene. *J Bacteriol* 2003;**185**:2936–43.
- Wani SA, Samanta I, Munshi ZH, et al. Shiga toxin-producing *Escherichia coli* and enteropathogenic *Escherichia coli* in healthy goats in India: occurrence and virulence properties. *J Appl Microbiol* 2006;**100**:108–13.
- Wirth T, Falush D, Lan R, et al. Sex and virulence in *Escherichia coli*: an evolutionary perspective. *Mol Microbiol* 2006;**60**:1136–51.
- Yatsuyanagi J, Saito S, Sato H, et al. Characterization of enteropathogenic and enteroaggregative *Escherichia coli* isolated from diarrheal outbreaks. *J Clin Microbiol* 2002;**40**:294–7.
- Yoshitomi KJ, Jinneman KC, Weagant SD. Optimization of a 3'-minor groove binder-DNA probe targeting the uidA gene for rapid identification of *Escherichia coli* O157:H7 using real-time PCR. *Mol Cell Probes* 2003;**17**:275–80.
- Zerbino DR. Using the Velvet de novo assembler for short-read sequencing technologies. *Curr Protoc Bioinformatics* 2010;**31**:11.5.1–11.5.12.
- Zerbino DR, Birney E. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res* 2008;**18**:821–9.
- Zhang H, Gao S, Lercher MJ, et al. EvolView, an online tool for visualizing, annotating and managing phylogenetic trees. *Nucleic Acids Res* 2012;**40**:W569–72.
- Zhou Z, Li X, Liu B, et al. Derivation of *Escherichia coli* O157:H7 from its O55:H7 precursor. *PLoS One* 2010;**5**:e8700.