

Bacteriological Quality and Incidence of Some Pathogenic Bacteria in Fresh White Cheese Sold in Tripoli, Libya

Yahia S Abujnah, Liala S El Magdoli, Said O Gnan, Mufida K Eljabali and Rabya A Lahmer*

Department of Food Science and Technology, Faculty of Agriculture, University of Tripoli, Tripoli, Libya

Abstract

The aim of this study was to evaluate the bacteriological quality and safety of fresh white cheese sold in Tripoli, Libya. The study lasted for approximately 7 months (November 2011-May 2012), during this period 87 fresh white cheese samples were collected from seven different areas (4 to 5 factories from each area with the rate of 3 duplicates). The samples were tested for temperature at receiving, pH and acidity, total aerobic counts, total coliform counts and the detection of incidence of some pathogenic bacteria including: *Escherichia coli*, *Escherichia coli* O157:H7, *Staphylococcus aureus*, *Salmonella* spp., *Aeromonas hydrophila* and *Listeria monocytogenes*. Results indicated that the average temperature, % acidity, and pH of the tested samples were ($15.80^{\circ}\text{C} \pm 5.8$, $0.21 \pm 0.02\%$ and 5.81 ± 0.06) respectively. Data indicated that 70.1% of the samples exceeded the Libyan standard for white cheese No. 366 in respect to temperature, while pH of all samples was within the limits of such standard. On the other hand, means of total aerobic counts, total coliform counts, and the numbers of *Staphylococcus aureus* were (38×10^7 , 74×10^5 , 35×10^4 and 53×10^3 cfu/g) respectively of the study.

Keywords: White cheese; Bacteriological quality; Pathogenic bacteria; *E. coli*; *Staphylococcus aureus*

Introduction

Cheese are mostly divided according to moisture content as very hard, hard, semi soft and soft cheese. Fresh white cheese is considered soft cheese and according to moisture content and nutrients it is subject to the growth of different microorganisms and pathogens. The presence of *E. coli* in fresh white cheese is an indication of low quality milk, poor production practices or contamination after processing [1-3]. Thus, Identification and enumeration of *E. coli* is widely used as an indicator for evaluating the safety of fresh white cheese. The presence of *E. coli* in fresh white cheese was confirmed by O'Brien et al. [1] in Ireland, Tekinsen and Ozdemir [2], Aygun et al. [4] in Turkey, Araujo et al. [5] in Brazil, Hamid and El-Owni [6] in Sudan and Abo-aza et al. [7] in Syria. Obviously, the magnitude of its presence differs from region to another. Hamid and El-Owni [6] and Karim et al. [8] found that fresh white cheese contains 2.3×10^6 cfu/g and 1.5×10^5 cfu/g, in Sudan and Syria respectively. Similarly, In Antakya, Turkey; Aygun et al. [7] found *E. coli* in fresh white cheese at a rate of 1.02×10^4 cfu/g. Moreover, in a study in Brazil [5] indicated that the range of *E. coli* was 1.0×10^3 to 1.8×10^8 cfu/g. Milk and milk products are important cause of out breaks by *Staphylococcus aureus* in many areas around the world and soft white cheese was reported to be the primary source for such pathogen yet its presence differs from study to another. The percentage of its presence in white cheese varied from 4%, O'Brien et al. [1] to 30.9% Moraes, et al. [9]. Furthermore, it was isolated from 20.6%, 16.3%, 15%, and 14.16% of white cheese by Araujo et al. [5]; De Luca, et al. [10], Karim et al. [8] and Al-abidy and Sharif [11] respectively. Thus, different studies were oriented to the enumeration of *Staph. aureus* in fresh cheese in different countries. For instance, in Turkey, Irkin [12], Aygun et al. [7] and Tekinsen and Ozdemir [2] reported an average of 1.5×10^3 cfu/g, 2.5×10^3 cfu/g and 1.2×10^6 cfu/g, respectively. On the other hand, Abo-aza et al. [1] and Hamid and El-owni [6] reported the presence of such pathogen in white cheese In Syria and Sudan up to 4.8×10^4 cfu/g and 1.1×10^4 cfu/g

respectively. *Salmonella* is considered the main causative of food poisoning, which contribute to about 90% of food outbreaks. Different investigators in many areas pointed the presence of *Salmonella* in fresh white cheese [2,4,6,13,14]. *Aeromonas hydrophila* is usually destroyed by pasteurization and its presence in white soft cheese is mostly an indication of post pasteurization contamination. Yet, *Aeromonas* spp. were detected in 17.7% of fresh white cheese samples from Rio de Janeiro, Brazil, 54.4% of those isolates were reported to be *A. hydrophila* [5]. Listeriosis is a very serious food poisoning illness caused by *Listeria* spp. especially *L. monocytogenes*, which was reported to be responsible for about 30% fetal, cases [15-18]. The main objective of this study was to evaluate the bacteriological quality and the presence of *E. coli*, *E. coli* O157: H7, *Staph. aureus*, *Salmonella* spp., *A. hydrophila* and *Listeria monocytogenes* in fresh white cheese sold in Tripoli, Libya. Additionally to investigate the potential effect of sample source on the existence of those microorganisms.

Materials and Methods

The study lasted for approximately 7 months (Nov. 2011-May 2012). During that period 87 fresh white cheese samples were collected from local small factories in 7 areas around Tripoli, Libya (4-5 factories in each area) and transferred aseptically in cooled ice box to the laboratory for further analysis. Fresh white cheese in the studied areas were produced from raw unpasteurized milk by adding rennet enzyme at 40°C and remove the whey immediately after coagulation

*Corresponding author: Rabya A Lahmer, Department of Food Science and Technology, Faculty of Agriculture, University of Tripoli, Tripoli, Libya, Tel: +218 21-4627901; E-mail: rabyalahmer@yahoo.co.uk

Received May 30, 2016; Accepted June 24, 2016; Published July 04, 2016

Citation: Abujnah YS, El Magdoli LS, Gnan SO, Eljabali MK, Lahmer RA (2016) Bacteriological Quality and Incidence of Some Pathogenic Bacteria in Fresh White Cheese Sold in Tripoli, Libya. J Microb Biochem Technol 8: 307-311. doi: [10.4172/1948-5948.1000301](http://dx.doi.org/10.4172/1948-5948.1000301)

Copyright: © 2016 Abujnah YS, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

and subjected to selling. The samples were subjected to measuring temperature at receiving, pH (WTW, 300) at 25°C as mentioned by Graham [19] and titratable acidity as specified by Hooi et al [20]. Different microorganisms were enumerated using standard methods for the examination of dairy products [19-21].

Detection of Pathogenic Bacteria

E. coli was isolated and detected as specified by Davidson et al. [22], Hary and Pall [23]. Meanwhile, *E. coli* O157:H7 was detected as specified by Meng et al. [21], Hary and Pall [23]. *Salmonella* were detected using API 20 E System and identified using BD Phoenix, ISO 6579, Hary and Pall [23]. *Staphylococcus aureus*: *Staphylococcus aureus* was detected as specified by Lancette and Bennett [24]. *A. hydrophila* was detected and identified using API 20 E System as specified by Palumbo et al. [25] and ISO 6579. *L. monocytogenes* was isolated and detected as specified by Gasanov et al. [26], Hary and Pall [23].

Statistical Analysis

The data obtained were analyzed statistically using SAS System as specified by Abosalah and Awad [27].

Results and Discussion

Results in Table 1 showed that overall means at receiving point in the factories of temperature, pH, and titratable acidity were $15.8 \pm 5.9^\circ\text{C}$, 5.81 ± 0.007 and $0.21 \pm 0.03\%$ respectively. The relatively high temperature of receiving of some samples coincides with high acidity and low pH, beside low quality unpasteurized milk used and the possibility of contamination with spoilage and pathogenic microorganisms including mastitis causing types is consider hazard to consumer. The effect of sample source on temperature at receiving as well as pH, and titratable acidity could be clearly seen from data presented in Table 2. Results indicated that source Z was the highest in temperature ($16.7 \pm 6.81^\circ\text{C}$) and source A was the lowest $14 \pm 6.74^\circ\text{C}$. Yet, statistical analysis indicated that source of samples has no significant effect ($P < 0.05$) on temperature of receiving. Concerning pH and % of titratable acidity statistical analysis showed that samples source has significant effect ($P < 0.05$) on acidity and pH. Source H was the lowest in acidity ($0.18 \pm 0.01\%$) and highest in pH (6.10 ± 0.02) while source Z showed the highest in acidity ($0.24 \pm 0.01\%$) and lowest in pH (5.83 ± 0.02). This could be related to high temperature of receiving since source Z was the highest in temperature of receiving and could

Test	Means \pm S.D
Temperature ($^\circ\text{C}$)	15.80 ± 5.9
Acidity (%)	0.21 ± 0.03
pH	5.81 ± 0.07

Table 1: Means and (SD) of temperature, acidity and pH of all samples.

Test Source	No. of Samples	Temperature ($^\circ\text{C}$)	Acidity (%)	pH
A	13	14.0 ± 6.74^a	0.20 ± 0.02^b	5.93 ± 0.1^b
B	15	15.8 ± 5.93^a	0.19 ± 0.02^c	5.95 ± 0.03^b
C	12	16.5 ± 5.54^a	0.21 ± 0.01^b	5.88 ± 0.05^c
D	11	16.5 ± 5.25^a	0.23 ± 0.02^a	5.83 ± 0.03^c
H	12	15.0 ± 6.0^a	0.18 ± 0.01^c	6.10 ± 0.02^a
E	12	16.5 ± 5.38^a	0.19 ± 0.01^c	5.96 ± 0.03^a
Z	12	16.7 ± 6.81^a	0.24 ± 0.01^a	5.83 ± 0.02^c
Over all mean		15.80 ± 5.8	0.21 ± 0.03	5.81 ± 0.06

^{abc}LS Means with the different letter within the attribute are significantly different

Table 2: Means and (SD) of temperature, acidity and pH by sources.

be of highly contaminated with spoilage bacteria. Results of the present study are in agreement with data reported by Karim et al. [8] in Syria. Table 3 presents means and variation of total aerobics, total coliform and *Staphylococcus aureus* in all samples. The mean number of total aerobic counts, coliforms, most probable number of thermo-tolerance and *Staphylococcus aureus* were 38×10^7 cfu/g, 74×10^5 cfu/g, 35×10^4 cfu/g and 53×10^3 cfu/g respectively. The highest number of *E. coli* in this study could be related to low quality of milk used, unhygienic production during manufacturing, contamination after processing and the use of unpasteurized milk. Results of the present study are relatively lower than those reported by Moraes et al. [9] of 18×10^6 cfu/g and higher than those reported by Oksuz and Aici [28] of 10^2 - 10^5 cfu/g. The overall mean of *Staphylococcus aureus* of all samples studied were 53×10^3 cfu/g. Results of this study were close to the finding of Irkin [12] of 15×10^3 cfu/g, but relatively higher than the results of Karim et al. [8] of 2×10^3 cfu/g and lower than the results of Aygun et al. [4] of 25×10^5 cfu/g. Figure 1 presents the number and percentage of pathogenic bacteria in all samples. The number and percentage of samples which contain *E. coli*, *E. coli* O157:H7, *Salmonella*, *Staphylococcus aureus*, *A. hydrophila* and *Listeria monocytogenes* were 72 (82.7%), 31 (35.6%), 7 (8%), 62 (71.2%), 21 (24.1%) and zero consequently. The high percentage of *E. coli* (82.75%, $P < 0.05$) in this study could be related to the use of low quality milk, unhygienic production, use of unpasteurized milk and contamination after processing. Generally speaking, results of the present study were relatively better than the finding of O'Brien et al. [1] and Araujo et al. [5], who reported that 97% and 97.7% of their samples contained *E. coli* respectively and higher than the results of Karim [8], which showed only 70%. *E. coli* O157:H7 were present in 31 (35.6%) of samples studied. Results of this study were higher than the results of Mora et al. [29], which indicated that only 7.8% from the fresh white cheese samples studied in Lima, Peru contain *E. coli* O157:H7. *Salmonella* was isolated from 7 samples (8.04%) of all samples studied, one of the isolates was identified as being *S. cholera* and the other was *S. paratyphi* A (Tables 4 and 5). Results of this study were relatively lower ($P < 0.05$) than the findings of Todd [30] in Mexico, and Karim et al. [8] in Syria who reported that *Salmonella* was isolated from 13% and 18% of cheese samples respectively. On the other hand, these results

Test	Means	Range	
		Minimum	Maximum
c (cfu/g)Total aerobi	38×10^7	29×10^4	10×10^9
Total Coliforms (cfu/g)	74×10^5	15×10^3	14×10^7
(cfu/g)	53×10^3	50×10^0	87×10^5

Table 3: Means and ranges of total aerobic, total coliform, of thermo-tolerant and in all samples.




Figure 1: Percentages of pathogenic bacteria in samples.

Salmonella type	Total isolates	% of Isolates
	5	71.4
	1	14.3
	1	14.3

Table 4: Numbers and percentage of spp. in samples.

Parameter	Maximum Permitted Limit by Libyan Standards	Number/(%) out of Libyan Standards
% acidity	Not more than (1.9%)	0/(0.00)
Temperature (°C)	4-10	61/(70.1)
Total Coliforms (cfu/g)	Not more than 5 cfu/g	87/100
Pathogenic bacteria (cfu/g)	Zero	87/100

Table 5: Numbers and percentage of fresh cheese samples out of the Libyan standards for % acidity and temperature and Pathogenic bacteria.

were higher than those reported by AL-Abidy and Sharif [11] in Iraq and Tekinsen and Ozdemir [2] in Turkey of 5% and 6% respectively. *Staphylococcus aureus* was isolated from 62 (71.2%) of the samples studied. These results were relatively higher than those reported by Karim [8] in Syria (5%) and Araujo et al. [5] in Brazil (20%). *Aeromonas hydrophila* was found in 21 samples (24.1%) studied. These results were very similar compared with the study of Effat [31] in Egypt, who isolated *A. hydrophila* from 23.1% of samples of Demiate cheese and less in Garies cheese. *Listeria monocytogenes* was not detected in any of the samples studied. Results of this study were relatively similar to the finding of Moraes et al. [9], Hamid and Elowni [6]. However O'Brien et al. [1] and Colak et al. [13] reported the presence of *L. monocytogenes* in 5% and 4.8% of soft white cheese produced in Ireland and Turkey, respectively.

The Effect of Sample Source on Pathogenic Bacteria

Figure 2 showed that sources A and Z were the highest in *E. coli* (100%) source E was the lowest (41.6%) and sources B, C and D were almost the same. At the same time source Z was the highest in *E. coli* O157:H7 (58.3%) and source H and E were the lowest (16.6%) (Figure 3). The effect of sample source on *Salmonella* is shown in Figure 4. The source Z was the highest of 25% and in source B, D and E were 14.2%, 9.8% and 9.8% and not isolated in sources A, D and H. Figure 5 showed the effect of sample source on *Staphylococcus aureus*. The source E and Z were the highest (100%), source C and D (90.6%) and source H was the lowest. In the case of *A. hydrophila* source B was the highest (40%) and source E was the lowest (8%) (Figure 6). Statistical analysis indicated that the source of samples has no significant effect at 5% level on the presence of *E. coli* O157:H7 and *A. hydrophila* while it has significant effect on *E. coli* and *Staphylococcus aureus*. Several factors could contribute to these differences between samples sources including source of raw milk used in manufacturing of fresh white cheese, acidity of milk, temperature of transportation and storage, hygienic status at production point, use of milk from mastitis animals, the use of unpasteurized milk, contamination during and/or after processing. These factors were also confirmed by Gilmour and Row [32], Bramley and Mckinnon [33], Karim et al. [8], and O'Brien et al. [8], Shata and El-Mgdob [33,34]. Finally, the results of this study were evaluated according to Libyan standards of fresh white cheese [35-37]. As presented in Table 5, data of acidity and temperature at receiving point showed 100 and 70.1% agreement with Libyan standards, respectively [38]. On the other hand, total coliforms and other pathogenic bacteria were 100% out of standards. This is a clear indicator for a very low bacteriological quality of fresh white cheese produced at that period. Mostly source Z was the worst of all sources.




Figure 2: Percentages of *E. coli* in samples according to source.




Figure 3: Percentages of *E. coli* O157:H7 in samples according to source.




Figure 4: Percentages of *Salmonella* in samples according to source.

This could be due the area is crowded, undeveloped, poor management, poor sanitation procedures and further far from milk supply.

Conclusion

The relatively high microbial count accompanied with a rise in cooling temperature and the higher incidence of several pathogenic bacteria in the tested cheese samples reflecting the poor sanitation and hygienic practices in those sources and signaling an alarming health threatening problem with the potential occurrence of infection and food poisoning outbreaks among white cheese consumers in Tripoli during the period of this study.




Figure 5: Percentages of *Staphylococcus aureus* in samples according to source.




Figure 6: Percentages of *A. hydrophila* in samples according to source.

References

- O'Brien M, Hunt K, Mc Sweeney S, Jordan K (2009) Occurrence of foodborne pathogens in Irish farmhouse cheese. *International Journal of Food Microbiology* 26: 910-914.
- Tekinsen KK, Ozdemir Z (2006) Prevalence of foodborne pathogens in Turkish Van otlu (Herb) cheese. *Food Control* 17: 707-711.
- Quinto EJ, Cepeda A (1997) Incidence of toxigenic *Escherichia coli* in soft cheese made with raw or pasteurized milk. *Lett Appl Microbiol* 24: 291-295.
- Aygun O, Aslantas O, Oner S (2005) A survey on the microbiological quality of Carra, a traditional Turkish cheese. *Journal of Food Engineering* 66: 401-404.
- Araujo VS, Pagliares VA, Queiroz MLP, Freitas-Ameida AC (2002) Occurrence of *Staphylococcus* and Enteropathogens in soft cheese commercialized in the city of Rio de Janeiro, Brazil. *Journal of Applied Microbiology* 92: 1172-1177.
- Hamid OA, Owni OA (2007) Microbiological properties and sensory characteristics of white cheese (Gibnabayda) Collected in Zalingei Area, West Darfur State. *Journal of Animal and Veterinary Sciences* 2: 61-65.
- Aboaza S, Hadel A, Habiba F (2010) The effect of fresh white cheese processing on microbial quality. *University of Damascus Agriculture Journal Syria* 26: 247-260.
- Karim Y, Abaaja S, Salig S (2007) Study of some properties of fresh white cheese produced from fresh milk in Syria. *University of Damascus Agriculture Journal* 23: 299-315.
- Moraes PM, Viosa GN, Yamazi AK, Ortolani MBT, Nero LA (2009) Foodborne pathogens and microbiological characteristics of raw milk soft cheese produced and on retail sale in Brazil. *Foodborne Pathogens and Disease* 6: 245-249.
- De Luca G, Zanetti F, Stampi S (1997) *Staphylococcus aureus* in dairy products in the Bologna area. *Int J Food Microbiol* 35: 267-270.
- AL-abidy S, Sharif A (2008) The microbial contamination of fresh white cheese produced in Nenawy, Iraq. *Rafidain Journal of Science* 19: 115-129.
- Irkin R (2010) Determination of microbial contamination sources for use in quality management of cheese industry: "Dil" cheese as an example. *Journal of Consumer Protection and Food Safety* 5: 91-96.
- Colak H, Hampikyan BE, Uisoy B (2007) Prevalence of *L. monocytogenes* and *Salmonella* spp. in Tulum cheese. *Food Control* 18: 576-579.
- Ellis A, Preston M, Borczyk A, Miller B, Stone P, et al. (1998) A community outbreak of *Salmonella berta* associated with a soft cheese product. *Epidemiol Infect* 120: 29-35.
- Ashour M (2005) Bacterial contamination of dairy and its relation with human health. (1st edn), Almaaref Publishing, Alexandria, Egypt.
- Donnelly CW (2004) Growth and survival of microbial pathogens in cheese. In: "Cheese: Chemistry, Physics and Microbiology. (3rd Edn), General Aspects, Elsevier Ltd.
- Food and Agricultural Organisation (FAO) Risk Assessment of *Listeria monocytogenes* in ready-to-eat Foods: Technical Report. World Health Organisation and Food and Agricultural Organisation of the United Nations.
- Arsian S, Özdemir F (2008) Prevalence and antimicrobial resistance of *Listeria* spp. in homemade white cheese. *Food Control* 19: 360-363.
- Graham T (2004) Sampling dairy and related products. In: "Standard Methods for the Examination of Dairy Products". (17th Edn) American Public Health Association, Washington, DC.
- Hooi R, Barbano DM, Bradley RL, Budde D, Bulthaus M, et al. (2004) Chemical and physical methods. In: "Standard Methods for Examination of Dairy Products". (17th Edn), American Public Health Association, Washington, USA.
- Meng J, Feng P, Doyle MP (2001) Pathogenic *Escherichia coli* In: "Compendium of Methods for the Microbiological Examination of foods". (4th Edn), American Public Health Association, Washington, DC, USA.
- Davidson PM, Roth LA, Gambrel-Lenarz SA (2004) Coliform and other indicator bacteria. In: "Standard Methods for the Examination of Dairy Products". American Public Health Association, Washington, DC, USA.
- Hary OS, Pall GF (1992) Laboratory manual for microorganisms. Translated by Abdalhafid AM and Mobarak MAS. Alarabia Publishing, (2nd Edn), Cairo, Egypt.
- Lancette GA, Bennett BW (2001) *Staphylococcus aureus* and *Staphylococcal enterotoxins*. In: "Compendium of Methods for the Microbiological Examination of foods". American Public Health Association, Washington, DC, USA.
- Palumbo S, Abcyta C, Stelma G (2001) *Aeromonas*, *Arcobacter* and *Plesiomonas*. In: "Compendium of methods for the microbiological examination of foods". (4th Edn), American Public Health Association, Washington, DC, USA.
- Gasarov U, Hughes D, Hansbro PM (2005) Methods for the isolation and identification of *Listeria* spp. and *Listeria monocytogenes*: A review. *FEMS Microbiology Reviews* 29: 851-875.
- Abosalah MS, Awad AM (2008) Introduction to Statistics. (2nd edn) Dar-Almasira Publishing, Amman Jordan.
- Oksuz O, Aici M (2004) Incidence of *Escherichia coli* O157 in raw milk and white pickled cheese manufactured from raw milk in Turkey. *Food Control* 15: 453-456.
- Mora A, Leon SL, Blanco M (2007) Phage types, virulence genes and PFGE profiles of shiga toxin-producing *Escherichia coli* O157:H7 isolated from raw beef, soft cheese and vegetables in Lima (Peru). *International Journal of Food Microbiology* 114: 204-210.
- Todd ECD (2011) The international risk governance council framework and its application to *Listeria monocytogenes* in soft cheese made from unpasteurised Milk. *Food Control* 10: 1-27.
- Effat BA, Hosny M, Dabiza NM (2000) Occurrence of *Aeromonas hydrophila* and its growth in Egyptian soft cheese. *Egyptian Journal of Dairy Science* 28: 1-12.
- Gilmour A, Row MT (1990) Micro-organisms associated with milk. In: "Dairy Microbiology". (2nd Edn) Applied Science Publishers, London and New York.
- Bramley AJ, McKinnon CH (1990) The microbiology of raw milk in "Dairy microbiology. (2nd Edn), Applied Science Publishers, London and New York.

34. Shata AS, EL-Magdob MN (2003) Microbiological quality of dairy farms and Plants. (1st Edn) Enahada Publishing, Cairo, Egypt.
35. International Organization for Standardization (ISO). Microbiology of food and animal feeding stuffs: Horizontal method for the detection of Salmonellae sops ISO, Geneva, Switzerland.
36. International Organization for Standardization (ISO). Microbiology of food and animal feeding stuffs: horizontal method for the detection of *Escherichia coli* ISO, Geneva, Switzerland.
37. Libyan Standard for fresh white cheese, National Center for Standardization and Meteorology, Tripoli, Libya.
38. Kliver DO (2002) Diseases transmitted by food. (1st Edn) King Faisal University, Saudi Arabia.

Citation: Abujnah YS, El Magdoli LS, Gnan SO, Eljabali MK, Lahmer RA (2016) Bacteriological Quality and Incidence of Some Pathogenic Bacteria in Fresh White Cheese Sold in Tripoli, Libya. J Microb Biochem Technol 8: 307-311. doi: [10.4172/1948-5948.1000301](https://doi.org/10.4172/1948-5948.1000301)

OMICS International: Open Access Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700+ Open Access Journals
- 50,000+ editorial team
- Rapid peer review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled for better prominence and citations
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Best discounts for your subsequent articles

Submit your manuscript at: <http://www.editorialmanager.com/jmbt>