

Antimicrobial Resistance, Resistance Genes and Virulence Genes in Salmonella Isolates From Chicken

Yahong Liu, Yurong Yang, Xiaoping Liao, Liang Li,
Chunyin Lei, Lulu Li, Jain Sun and Baotao Liu

Guangdong Provincial Key Laboratory of Veterinary Pharmaceutics Development and
Safety Evaluation, College of Veterinary Medicine, South China Agriculture University,
Wushan Road, Tianhe District, 510642 Guangzhou, China

Abstract: Ninety eight *Salmonella* isolates (53 isolates from 2007 and 45 from 2009) from diseased chicken were examined for antimicrobial susceptibility to 15 antimicrobials, possession of resistance and virulence genes and Pulsed Field Gel Electrophoresis (PFGE) patterns. The 82 (84%), all 45 from 2009 and 37 from 2007 were resistant to 2 or more antimicrobials. A single isolate from 2009 was resistant to 13 of the 15 antimicrobials tested. The isolates from 2009 exhibited significantly greater resistance to streptomycin, florfenicol, tetracycline, doxycycline and nalidixic acid than that from 2007. Resistance genes *sul3* and *aadA1* were the most prevalent being found in 19 (36%) and 14 (31%) isolates from 2007 and 2009, respectively. All 98 isolates carried *invA*; in comparison with the isolates from 2007, the isolates from 2009 exhibited significantly lower rates of carrying *spvC*, *sopE* and *iroB*. Of the 98 isolates, 75 isolates were successfully typed, resulting in 49 different PFGE patterns with a difference of at least seven bands. This study shows that the majority of *Salmonella* strains from Guangdong display resistance to multiple antimicrobial compounds and carry multiple resistance genes and virulence genes.

Key words: Antimicrobial agents, resistance, resistance genes, virulence genes, salmonella

INTRODUCTION

Salmonella sp. is a gram negative, rod shaped, motile and facultative anaerobe bacterium that normally resides in the gut of wild and domestic animals (Pang *et al.*, 2011). *Salmonella enterica* serovar Enteritidis (*S. enteritidis*) is a major cause of food borne gastroenteritis in humans worldwide. Poultry and poultry products are considered the major vehicles of transmission to humans (Shah *et al.*, 2011). *Salmonella* contamination continues to be one of the major concerns for the microbiological safety of raw poultry products. The US Department of Agriculture's Food Safety Inspection Service (USDA-FSIS) has estimated that in 2007 poultry products accounted for approximately 60% of the food borne illnesses originating from *Salmonella* (Benli *et al.*, 2011). *Salmonella* was the common contamination of poultry products with the prevalence in chicken meat of 51.7% in Tunisia, 54% in China and 66% in Thailand (Abbassi-Ghozzi *et al.*, 2011; Yang *et al.*, 2010). *S. enterica* can easily cross contaminate other ready to eat foods exposed to these

surfaces, posing a risk for foodborne illness outbreaks. The FDA recommended practice of washing kitchen implements with soap, hot water and vigorous mechanical scrubbing can remove *S. enterica* effectively and hence reduce cross contamination (Ravishankar *et al.*, 2010).

The ability of chickens to carry *Salmonella* without displaying disease symptoms is responsible for *Salmonella* propagation in poultry stocks and for subsequent human contamination through the consumption of contaminated eggs or meat. The selection of animals more resistant to carrier state might be a way to decrease the propagation of *Salmonella* in poultry stocks and its transmission to humans (Calenge *et al.*, 2009). Wisner *et al.* (2010) reported that the *S. enteritidis* SPI-2 T3SS facilitates invasion and systemic spread in chickens although alternative mechanisms for these processes appear to exist (Wisner *et al.*, 2010). Pulsed Field Gel Electrophoresis (PFGE) has been successfully used for typing of *S. enteritidis*, *S. typhi* and *S. typhimurium* furthermore which has been accepted as the gold standard for *Salmonella* molecular typing by Pulse

Corresponding Author: Yahong Liu, Guangdong Provincial Key Laboratory of Veterinary Pharmaceutics Development and Safety Evaluation, College of Veterinary Medicine, South China Agriculture University, Wushan Road, Tianhe District, 510642 Guangzhou, China

Net International, the international molecular subtyping network for foodborne disease surveillance (Chen *et al.*, 2011; Hur *et al.*, 2011a, b).

The objective this study was to determine the prevalence of antimicrobial resistance and their associated genes, virulence associated genes and to analyze the PFGE patterns of Salmonella strains isolated from chicken with diarrhea. The isolates were divided into 2 groups by collection period (2007 or 2009) to investigate trends over time. Researchers also analyzed the variance in strain characteristics between the 2 groups.

MATERIALS AND METHODS

Bacterial isolates: Ninety eight Salmonella were collected from diseased chicken in Guangdong province, China, in 2007 and 2009. Salmonella strains were mainly isolated from fecal swabs taken from diseased chicken with white diarrhea. Each isolate was taken from an individual animal.

Antimicrobial susceptibility testing: Antimicrobial susceptibility testing was performed on all 98 Salmonella isolates using the agar dilution method according to the guidelines of the Clinical and Laboratory Standards Institute (CLSI) in 2008. The following antimicrobials were used: Ampicillin (AMP), Ceftiofur (CEF), Ceftriaxone (CRO), Streptomycin (STR), Gentamicin (GEN), Apramycin (APM), Chloramphenicol (CHL), Florfenicol (FEN), Tetracycline (TET), Doxycycline (DOX), Sulfamethoxazole (SMZ), Nalidixic Acid (NA), Ciprofloxacin (CIP), Enrofloxacin (ERO), Levofloxacin (LEF). The reference strain, *E. coli* ATCC 25922 was used as a quality control strain for determining the minimum inhibitory concentrations of the 15 antimicrobial agents.

Resistance and virulence genes: All isolates were screened for 23 resistance genes and 17 virulence genes with PCR as reported earlier (Del Cerro *et al.*, 2003; Karasova *et al.*, 2009).

Pulsed Field Gel Electrophoresis (PFGE): PFGE was used to analyze the genomic relatedness among Salmonella isolates from diseased chicken. The method used was basically according to Chen *et al.* (2011). PFGE of chromosomal DNA digested with the restriction enzyme XbaI was carried out according to a standard protocol using a CHEF-MAPPER System (Bio-Rad Laboratories, Hercules, CA). The gels were run at 6.0 V cm⁻¹ with an angle of 12°C at 14°C for 20 h and the results were interpreted according to the criteria of Tenover *et al.* (1995) Salmonella ser. Braenderup H9812 standards served as size markers.

Statistical analysis: Differences in the year by year rates of antimicrobial resistance were assessed using Fisher’s exact tests (SPSS 17.0). A p<0.05 was considered significant.

RESULTS

Antimicrobial resistance phenotypes: The results of antimicrobial susceptibility tests were shown in Table 1. None of the Salmonella isolates were resistant to CRO and LEF, none of the Salmonella isolates collected from 2007 were resistant to GEN and ERO. In comparison with the isolates from 2007, the isolates from 2009 exhibited significantly greater resistance to STR, FEN, TET, DOX and NA (p<0.05 to p<0.01). Of the 98 isolates, 82 (84%) all 45 from 2009 and 70% of those from 2007 were resistant to 2 or more antimicrobials. A single isolate from 2009 was resistant to 13 of the 15 antimicrobials tested.

Resistance genes: Seven of the 27 resistance genes (*bla cmy-2*, *rmtB*, *tetB*, *tetC*, *sul2*, *qnrS* and *qepA*) were not detected in any of the isolates. The results of PCR identification of the other genes associated with antimicrobial resistance were shown in Table 2. Sul3 and aadA1 was the most prevalent which were found in 19 (36%) and 14 (31%) isolates from 2007 and 2009, respectively. In comparison with the isolates from 2007, the isolates from 2009 exhibited significantly higher rates of carrying aph(3’)-VII, aadA1 and aadA2 (p<0.05 to p<0.01).

Virulence genes: The results of PCR identification of virulence genes were shown in Table 3. All 98 isolates carried *invA*; in comparison with the isolates from 2007, the isolates from 2009 exhibited significantly lower rates of carrying *spvC*, *sopE* and *iroB* (p<0.05 to p<0.01).

Table 1: Antimicrobial resistance rates of 98 Salmonella isolates collected from Guangdong province

Antimicrobial	Collection period, number and percentage of resistant	
	2007 (n = 53)	2009 (n = 45)
Ampicillin (AMP)	30 (57)	27 (60)
Ceftiofur (CEF)	24 (45)	20 (44)
Ceftriaxone (CRO)	0 (0)	0 (0)
Streptomycin (STR)	18 (34)	29 (64) ^a
Gentamicin (GEN)	0 (0)	4 (9)
Apramycin (APM)	1 (2)	6 (13)
Chloramphenicol (CHL)	4 (8)	9 (20)
Florfenicol (FEN)	7 (13)	30 (67) ^a
Tetracycline (TET)	4 (8)	28 (62) ^a
Doxycycline (DOX)	4 (8)	15 (33) ^a
Sulfamethoxazole (SMZ)	34 (64)	34 (76)
Nalidixic Acid (NA)	5 (9)	24 (53) ^a
Ciprofloxacin (CIP)	2 (4)	6 (13)
Enrofloxacin (ERO)	0 (0)	7 (16)
Levofloxacin(LEF)	0 (0)	0 (0)

^aSignificantly different from the rate in 2007 (p<0.05)

Table 2: Presence of genes associated with antimicrobial resistance in the 98 isolates

Resistance genes	Collection period, number and percentage of resistant	
	2007 (n = 53)	2009 (n = 45)
<i>bla_{TEM}</i>	11 (21)	0 (0)
<i>bla_{SHV}</i>	1 (2)	2 (4)
<i>bla_{DHA}</i>	0 (0)	3 (7)
<i>aac (3)-IV</i>	0 (0)	4 (9)
<i>aph (3')-VII</i>	3 (6)	13 (29) ^a
<i>aadA1</i>	6 (11)	14 (31) ^a
<i>aadA2</i>	3 (6)	12 (27) ^a
<i>cat1</i>	0 (0)	1 (2)
<i>cat2</i>	4 (8)	2 (4)
<i>cmlA</i>	1 (2)	7 (16)
<i>cmlB</i>	0 (0)	2 (4)
<i>floR</i>	7 (13)	4 (9)
<i>tetA</i>	5 (9)	10 (22)
<i>sul1</i>	8 (15)	10 (22)
<i>sul3</i>	19 (36)	7 (16)
<i>qnrA</i>	3 (6)	0 (0)
<i>qnrB</i>	11 (21)	1 (2)

^aSignificantly different from the rate in 2007 (p<0.05)

Table 3: Presence of virulence genes in the 98 isolates

Virulence genes	Collection period, number and percentage of resistant	
	2007 (n = 53)	2009 (n = 45)
<i>spvB</i>	7 (13)	11 (24)
<i>spvC</i>	12 (23) ^a	1 (2)
<i>spvD</i>	5 (9)	5 (11)
<i>invA</i>	53 (100)	45 (100)
<i>sopE</i>	14 (26) ^a	1 (2)
<i>phoP/Q</i>	47 (89)	35 (78)
<i>stn</i>	22 (42)	23 (51)
<i>sodCI</i>	24 (45)	30 (67)
<i>sodCII</i>	49 (92)	43 (96)
<i>iroB</i>	24 (45) ^a	5 (11)
<i>hin/H2</i>	36 (68)	29 (64)
<i>repFIIA</i>	6 (11)	6 (13)
<i>shvA</i>	50 (94)	41 (91)

^aSignificantly different from the rate in 2009 (p<0.05)

Genetic relatedness by PFGE: All Salmonella isolates were analyzed for their genetic relatedness by using PFGE. Of the 98 isolates, 75 isolates were successfully typed, resulting in 49 different PFGE patterns with a difference of at least seven bands. This suggests that dissemination of the Salmonella isolates might not be due to the spread of a specific clone. However, in a small number of cases, isolates from the same farms or from different farms were found to have identical PFGE patterns.

DISCUSSION

Resistance phenotypes: Salmonella colonizes the gastrointestinal tracts of a wide range of wild and domestic animals including poultry raised for food (Lestari *et al.*, 2009). Antimicrobial resistance in Salmonella isolated from both food and veterinary clinical

sources appears to be increasing in many countries and regions (Dutil *et al.*, 2010; Iwabuchi *et al.*, 2011; Yan *et al.*, 2010; Yang *et al.*, 2010). In the present study, there was a high rate of resistance to particular antimicrobials, notably AMP, SMZ, regardless of collection period in agreement with earlier reports (Hur *et al.*, 2011b; Yang *et al.*, 2010). To some antimicrobials the rate was significantly increased by year which might be the result of the widely use of the antimicrobials. In addition to most of the antimicrobials, the antimicrobial resistance rates were higher in isolates from 2009, through there was no statistical significance.

Ceftiofur is the only cephalosporin approved for systemic use in food producing animals since, 2002 in China and it is highly effective against Salmonella isolates. The rate of resistance to CEF was higher in the study than in earlier studies (Dutil *et al.*, 2010; Lestari *et al.*, 2009; Wang *et al.*, 2010) presumably as a consequence of the increasing use of cephalosporins on animal farms. Prudent use of antimicrobials in veterinary practice is therefore fundamental to the reduction of resistance development. However, much higher rates were also reported such as 85.7% by Lu *et al.* (2011)'s report.

Fortunately, MIC assays in the present study indicated that >90% of the 98 isolates were within the susceptibility ranges of several antimicrobials including CRO, CIP, ERO and LEF. Thus, these antimicrobials are still potentially effective against Salmonella.

Resistance genes: Though many resistance genes were identified in different sources of Salmonella (Garcia-Fernandez *et al.*, 2009; Kozak *et al.*, 2009; Lu *et al.*, 2011; Rayamajhi *et al.*, 2010; Yang *et al.*, 2010) however few data are available on prevalence of 27 resistance genes in Salmonella from diseased chicken origin.

Resistance to ampicillin and cephalosporins in Gram negative bacteria is primarily mediated by β-Lactamases (BLAs) which hydrolyze the β-lactam ring and thus inactivate the antibiotic. Many different BLAs have been described such as TEM-, SHV-, CTX-M-, OXA- and CMY-type BLAs (Bradford, 2001). In the present study, researchers investigated the presence of BLAs encoding genes by using a set of primers for the conserved regions of common BLAs genes. PCR and DNA sequencing results showed that the gene *blaTEM-1* was identified in 11(21%) of Salmonella isolates from 2007 while none of isolates from 2009. In addition, researchers identified the *blaDHA-1* and *blaSHV-1* gene in three Salmonella isolates, respectively.

Lin *et al.* (2009) first examined the ciprofloxacin resistance level in the Salmonella strains isolated from

animal sources but researchers checked out the qnr resistance gene mediated by plasmid which can transfer to the recipient *E. coli* DH5a strain (Garcia-Fernandez *et al.*, 2009)

Virulence genes: In the present study virulence associated genes were examined involved of SPIs and the chromosomally encoded stn (Salmonella enterotoxin gene), phoP/Q (two component global regulator) and iroB (Parvathi *et al.*, 2011) and plasmid in Salmonella isolates from chicken. Salmonella virulence genes play important role in the pathogenicity of the organism and Salmonella pathogenesis is dictated by a group of genes responsible for colonization (Thiagarajan *et al.*, 1996) while Data showed that the deletion of SPI-1 does not affect cecal colonization in 1 week old chicken but causes a milder and delayed systemic infection (Desin *et al.*, 2009) in addition, SPI-1 genes are highly expressed at early and late stages of infection in cultured epithelial cells according to Hautefort's report (Hautefort *et al.*, 2008). In the present study, >90% of the 98 isolates from chicken carried the virulence genes of *sodCII* and *slyA*, >60% with *phoP/Q* and *hin/H2* and >40% with *stn* and *iroB* and all carried *invA* regardless of collection period. It is interesting that the rate of isolates carried virulence genes of *sopE* and *iroB* from 2009 was significantly lower than that from 2007 while the isolates from 2009 exhibited significantly greater resistance to some of the antimicrobials and higher rates of carrying some resistance genes. Whether the virulence is weakened with the resistance further studies are needed to examine this possibility.

PFGE: Pulsed field gel electrophoresis has been widely used to determine strain relatedness, confirm bacterial disease outbreaks and identify the sources of strains (Chen *et al.*, 2011; Gaul *et al.*, 2007; Lu *et al.*, 2011). In this study, the PFGE results indicated a genetically diverse Salmonella population whereas several indistinguishable PFGE patterns were shared among isolates obtained from different farms. The majority of these isolates exhibited similar resistance profile. This suggests that dissemination of the Salmonella isolates might not be due to the spread of a specific clone.

CONCLUSION

The results of this study showed that the majority of Salmonella strains display resistance to multiple antimicrobial compounds and carry multiple resistance genes and virulence genes. These findings indicate that a surveillance program is needed to employ effective control measures to reduce Salmonella contaminations and the levels of antimicrobial resistant Salmonella in poultry products.

ACKNOWLEDGEMENTS

This research was supported by the National Science Fund for Distinguished Young Scholars (Grant No. 31125026), the Special Fund for Agro-Scientific Research in the Public Interest (Grant No. 201203040), the National Natural Science Foundation of China (Grant No. U0631006 and U1031004).

REFERENCES

- Abbassi-Ghozzi, I., A. Jaouani, S. Hammami, J. Martinez-Urtaza, A. Boudabous and M. Gtari, 2011. Molecular analysis and antimicrobial resistance of *Salmonella* isolates recovered from raw meat marketed in the area of Grand Tunis. Tunisia. *Pathol. Biol.*, 60: 49-54.
- Benli, H., M.X. Sanchez-Plata and J.T. Keeton, 2011. Efficacy of epsilon-polylysine, lauric arginate, or acidic calcium sulfate applied sequentially for *Salmonella* reduction on membrane filters and chicken carcasses. *J. Food Protect.*, 74: 743-750.
- Bradford, P.A., 2001. Extended-spectrum beta-lactamases in the 21st century: Characterization, epidemiology and detection of this important resistance threat. *Clin. Microbiol. Rev.*, 14: 933-951.
- Calenge, F., F. Lecerf, J. Demars, K. Feve and F. Vignoles *et al.*, 2009. QTL for resistance to *Salmonella* carrier state confirmed in both experimental and commercial chicken lines. *Anim. Genet.*, 40: 590-597.
- Chen, M.H., W.Z. Wang, S.W. Wang, Y.C. Shih and H.Y. Tsen, 2011. Pulsed Field Gel Electrophoresis (PFGE) analysis for multidrug resistant *Salmonella enterica* serovar Schwarzengrund isolates collected in six years (2000-2005) from retail chicken meat in Taiwan. *Food Microbiol.*, 28: 399-405.
- Del Cerro, A., S.M. Soto and M.C. Mendoza, 2003. Virulence and antimicrobial-resistance gene profiles determined by PCR-based procedures for *Salmonella* isolated from samples of animal origin. *Food Microbiol.*, 20: 431-438.
- Desin, T.S., P.K.S. Lam, B. Koch, C. Mickael and E. Berberov *et al.*, 2009. *Salmonella enterica* serovar enteritidis pathogenicity island 1 is not essential for but facilitates rapid systemic spread in chickens. *Infect. Immun.*, 77: 2866-2875.
- Dutil, L., R. Irwin, R. Finley, L.K. Ng and B. Avery *et al.*, 2010. Ceftiofur resistance in *Salmonella enterica* serovar Heidelberg from chicken meat and humans, Canada. *Emerg. Infect. Dis.*, 16: 48-54.
- Garcia-Fernandez, A., D. Fortini, K. Veldman, D. Mevius and A. Carattoli, 2009. Characterization of plasmids harbouring *qnrS1*, *qnrB2* and *qnrB19* genes in *Salmonella*. *J. Antimicrob. Chemother.*, 63: 274-281.

- Gaul, S.B., S. Wedel, M.M. Erdman, D.L. Harris, I.T. Harris, K.E. Ferris and L. Hoffman, 2007. Use of pulsed-field gel electrophoresis of conserved XbaI fragments for identification of swine *Salmonella* serotypes. J. Clin. Microbiol., 45: 472-476.
- Hautefort, I., A. Thompson, S. Eriksson-Ygberg, M.L. Parker and S. Lucchini *et al.*, 2008. During infection of epithelial cells *Salmonella enterica* serovar typhimurium undergoes a time-dependent transcriptional adaptation that results in simultaneous expression of three type 3 secretion systems. Cell. Microbiol., 10: 958-984.
- Hur, J., J.H. Kim, J.H. Park, Y.J. Lee and J. H. Lee, 2011b. Molecular and virulence characteristics of multi-drug resistant *et al* *Salmonella enteritidis* strains isolated from poultry. Vet. J., 189: 306-311.
- Hur, J., Y. Y. Choi, J.H. Park, B.W. Jeon, H.S. Lee, A.R. Kim and J.H. Lee, 2011a. Antimicrobial resistance, virulence-associated genes and pulsed-field gel electrophoresis profiles of *Salmonella enterica* subsp. *Enterica* serovar typhimurium isolated from piglets with diarrhea in Korea. Can. J. Vet. Res., 75: 49-56.
- Iwabuchi, E., S. Yamamoto, Y. Endo, T. Ochiai and K. Hirai, 2011. Prevalence of *Salmonella* isolates and antimicrobial resistance patterns in chicken meat throughout Japan. J. Food Protect., 74: 270-273.
- Karasova, D., H. Havlickova, F. Sisak and I. Rychlik, 2009. Deletion of *sodCI* and *spvBC* in *Salmonella enterica* serovar Enteritidis reduced its virulence to the natural virulence of serovars Agona, Hadar and Infantis for mice but not for chickens early after infection. Vet. Microbiol., 139: 304-309.
- Kozak, G.K., D.L. Pearl, J. Parkman, R.J. Reid-Smith, A. Deckert and P. Boerlin, 2009. Distribution of sulfonamide resistance genes in *Escherichia coli* and *Salmonella* isolates from swine and chickens at abattoirs in Ontario and Quebec, Canada. Appl. Environ. Microbiol., 75: 5999-6001.
- Lestari, S.I., F. Han, F. Wang and B. Ge, 2009. Prevalence and antimicrobial resistance of *Salmonella* serovars in conventional and organic chickens from Louisiana retail stores. J. Food Protect., 72: 1165-1172.
- Lin, C.C., T.H. Chen, Y.C. Wang, C.C. Chang, S.L. Hsuan, Y.C. Chang and K.S. Yeh, 2009. Analysis of ciprofloxacin-resistant *Salmonella* strains from swine, chicken and their carcasses in Taiwan and detection of *parC* resistance mutations by a mismatch amplification mutation assay PCR. J. Food Protect., 72: 14-20.
- Lu, Y., C.M. Wu, G.J. Wu, H.Y. Zhao and T. He *et al.*, 2011. Prevalence of antimicrobial resistance among *Salmonella* isolates from chicken in China. Foodborne Pathog. Dis., 8: 45-53.
- Pang, E., C. Tien-Lin, M. Selvaraj, J. Chang and K.J. Wang, 2011. Deletion of the *aceE* gene (encoding a component of pyruvate dehydrogenase) attenuates *Salmonella enterica* serovar Enteritidis. FEMS Immunol. Med. Microbiol., 63: 108-118.
- Parvathi, A., J. Vijayan, G. Murali and P. Chandran, 2011. Comparative virulence genotyping and antimicrobial susceptibility profiling of environmental and clinical *Salmonella enterica* from Cochin, India. Curr. Microbiol., 62: 21-26.
- Ravishankar, S., L. Zhu and D. Jaroni, 2010. Assessing the cross contamination and transfer rates of *Salmonella enterica* from chicken to lettuce under different food-handling scenarios. Food Microbiol., 27: 791-794.
- Rayamajhi, N., B.Y. Jung, S.B. Cha, M.K. Shin and A. Kim *et al.*, 2010. Antibiotic resistance patterns and detection of *blaDHA-1* in *Salmonella* species isolates from chicken farms in South Korea. Appl. Environ. Microbiol., 76: 4760-4764.
- Shah, D.H., X. Zhou, T. Addwebi, M.A. Davis and L. Orfe *et al.*, 2011. Cell invasion of poultry-associated *Salmonella enterica* serovar enteritidis isolates is associated with pathogenicity, motility and proteins secreted by the type III secretion system. Microbiol., 157: 1428-1445.
- Tenover, F.C., R.D. Arbeit, R. V. Goering, P.A. Mickelsen, B.E. Murray, D.H. Persing and B. Swaminathan, 1995. Interpreting chromosomal DNA restriction patterns produced by pulsed-field electrophoresis: Criteria for bacterial strain typing. J. Clin. Microbiol., 33: 2233-2239.
- Thiagarajan, D., M. Saeed, J. Turek and E. Asem, 1996. *In vitro* attachment and invasion of chicken ovarian granulosa cells by *Salmonella enteritidis* phage type 8. Infect. Immun., 64: 5015-5021.
- Wang, Y.C., Y.C. Chang, H.L. Chuang, C.C. Chiu and K.S. Yeh *et al.*, 2010. Antibiotic resistance, integrons and *Salmonella* genomic island 1 among *Salmonella* Schwarzengrund in broiler chicken and pig. Afr. J. Microbiol. Res., 4: 677-681.
- Wisner, A.L., T.S. Desin, B. Koch, P.K. Lam and E.M. Berberov *et al.*, 2010. *Salmonella enterica* subspecies enterica serovar Enteritidis *Salmonella* pathogenicity island 2 type III secretion system: Role in intestinal colonization of chickens and systemic spread. Microbiol., 156: 2770-2781.
- Yan, H., L. Li, M.J. Alam, S. Shinoda, S. Miyoshi and L. Shi, 2010. Prevalence and antimicrobial resistance of *Salmonella* in retail foods in northern China. Int. J. Food Microbiol., 143: 230-234.
- Yang, B., D. Qu, X. Zhang, J. Shen and S. Cui *et al.*, 2010. Prevalence and characterization of *Salmonella* serovars in retail meats of marketplace in Shaanxi, China. Int. J. Food Microbiol., 141: 63-72.