



Coagulase gene polymorphism of *Staphylococcus aureus* isolates: A study on dairy food products and other foods in Tehran, Iran

Mohammad Mehdi Soltan Dallal^{a,b,*}, Mohammad Reza Khoramizadeh^{c,d}, Solmaz Agha Amiri^b, Ali Akbar Saboor Yaraghi^{b,e}, Ramin Mazaheri Nezhad Fard^a

^a Division of Food Microbiology, Department of Pathobiology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

^b Food Microbiology Research Center, Tehran University of Medical Sciences, Tehran, Iran

^c Department of Medical Biotechnology, School of Advanced Technologies in Medicine, Iran

^d Biosensor Research Center, Endocrinology and Metabolism Cellular and Molecular Science Institute, Tehran University of Medical Sciences, Iran

^e Division of Immunology, Department of Pathobiology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

Received 24 March 2016; received in revised form 13 July 2016; accepted 8 September 2016

Available online 18 October 2016

Abstract

Coagulase is considered as a major determinant factor for the identification of *Staphylococcus aureus* strains. The 3'-end coding region of the coagulase (*coa*) gene contains a series of 81-bp tandem repeats, which differ in the number and location of enzymatic restriction sites among different isolates. *coa* PCR-RFLP has been used widely to type *S. aureus* isolates in epidemiological studies. The current study was conducted to investigate the coagulase gene polymorphisms in *S. aureus* isolated from various food samples using an *in house* PCR-RFLP method. A total of 100 strains of *S. aureus* were isolated from food samples. Isolates were typed by PCR-RFLP analysis using *NdeI* restriction digestion of the coagulase gene PCR products. Results showed that amplification of coagulase genes from *S. aureus* produced different PCR products. The isolates were grouped into 18 genotypes using RFLP analysis results of the genes. In this study, the *S. aureus* isolates have been shown to include more than one coagulase genotype, but only had a few *coa* genotypes predominated.

© 2016 Beijing Academy of Food Sciences. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Keywords: *Staphylococcus aureus*; PCR-RFLP; Coagulase gene; Food samples

1. Introduction

Staphylococcus aureus is an important food-borne pathogen [1]. It is among the most significant pathogens that cause various diseases in humans and animals. In humans, nosocomial and community acquired infections are the most frequently reported

problems caused by *S. aureus* [2,3]. The bacterium is one of the most significant pathogens causing intramammary infections (IMI) in dairy ruminants [4]. The primary reservoirs for *S. aureus* are the skin and mucous membranes, especially of the nasopharyngeal region of birds and mammals. This microorganism is found in 30%–80% of the human population, thus, unhygienic processing of foods has to be considered as a major risk of contamination [5–7]. Staphylococcal food poisoning (SFP) is considered to be one of the leading causes of all food-borne diseases [1]. In the last few decades, SFP has been reported as the third cause of food-borne infections in the world [5,8,9]. Milks, dairy products and meats, especially in traditional foods, play an important role in SFP; from which, *S. aureus* strains have been isolated frequently [1,10,11]. Genetic heterogeneity is considerable in natural population of *S. aureus* [12]. Relatively, many molecular techniques such as random amplified polymorphic

* Corresponding author at: Food Microbiology Research Center/Division of Food Microbiology, Department of Pathobiology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran.

E-mail addresses: soltanda@sina.tums.ac.ir, msoltandallal@gmail.com (M.M. Soltan Dallal).

Peer review under responsibility of Beijing Academy of Food Sciences.



Production and hosting by Elsevier

DNA (RAPD), ribotyping, multilocus enzyme electrophoresis (MLEE), plasmid profiling and coagulase gene polymorphism have been used for the identification and characterization of *S. aureus* in epidemiological studies [13,14]. There is no information on genetic diversity of *S. aureus* isolated from foods in Iran since the bacterial routine identification is carried on by conventional methods such as Gram staining and catalase, clumping factor, DNase and mannitol fermentation tests [15]. However, use of molecular techniques as rapid tools in microbiology research and diagnosis has been increased recently. One of these molecular techniques, PCR-based coagulase genotyping by RFLP analysis (*coa*-RFLP) of the 3' end of the gene encoding staphylococcal coagulase has been suggested as a simple and effective method for typing *S. aureus* isolates in epidemiological studies [16,17]. Numerous studies based on the coagulase gene polymorphism have been carried out for genotyping of *S. aureus* isolated from bovine mastitic milks and other foods [12,15,18,19]. The purpose of the current study was to identify *S. aureus* subtypes isolated from food samples, using *coa* gene polymorphism profile.

2. Materials and methods

2.1. Bacterial isolates

A total number of 100 *S. aureus* strains was isolated from food samples (78 isolates from dairy products, 16 isolates from meat products and six isolates from other foods). Samples were diluted with normal saline and then homogenized and seeded onto *Staphylococcus* selective media (Merck, Germany) and Baird–Parker agar (Merck, Germany). Colonies showing typical aspect of coagulase-positive staphylococci were identified by conventional methods, including Gram staining and catalase, clumping factor, DNase and mannitol fermentation tests. Isolates identified as *S. aureus* were frozen in skimmed milk containing 15% (v/v) glycerol at -20°C until use.

2.2. DNA extraction

DNA was extracted using QIAamp DNA mini kit (Qiagen, Germany), according to the manufacturer's instructions for Gram-positive bacteria.

2.3. PCR amplification

PCR of the *coa* gene was carried out using primers COAG₂: 5'-CGAGACCAAGATTCAACAAG-3' and COAG₃: 5'-AAAGAAAACCACTCATCA-3' described by Raimundo et al. in 1999 [20]. Reactions were prepared in a final volume of 25 μL using HotStarTaq Plus master mix kit (Qiagen, Germany), containing 12.5 μL of HotStarTaq Plus master mix, 2.5 μL of 10X buffer, 5 μL of RNase-free water, 20 pmol of each primer and 3 μL of the template. The amplification program included an initial denaturation step of 3 min at 94°C followed by 30 cycles of 1 min at 94°C , 1 min at 55°C and 1 min at 72°C and a final extension step of 5 min at 72°C . The amplicon size varied in various strains. *Staphylococcus*

epidermidis ATCC 12228 and *S. aureus* COL were used as controls.

2.4. Restriction enzyme digestion

Generally, 8.5 μL of the PCR product was incubated with 10 U of *Nde*I endonuclease enzyme (Fermentas, USA), 2.5 μL of restriction buffer and 13 μL of distilled water for 3 h at 37°C .

2.5. Agarose gel electrophoresis

Digested fragments and PCR products were separated in 1% and 2% agarose gels (Gibco, USA), respectively. A 100-bp ladder (Fermentas, USA) was used as a standard molecular marker for the calculation of the sizes of the *coa* and *Nde*I-generated *coa* fragments. Gels were visualized under UV light after staining with fluorescent dyes.

2.6. Specificity testing

For the specificity of the primer pair test, the DNA of *S. epidermidis* ATCC 12228 and *S. aureus* COL strain was analyzed.

2.7. Reproducibility testing

For the reproducibility of PCR, 5 selected isolates were chosen randomly and tested by twice submitting 3 different PCR products to *Nde*I digestion.

2.8. Data analysis

The software was used for the size of PCR and RFLP products. Numeric codes were assigned to the PCR genotypes and RFLP patterns.

3. Results and discussion

All isolates produced PCR amplicon with the COAG₂ and COAG₃ primers (Fig. 1). The agarose gel analysis of the digestion products showed ten different sizes, ranging from approximately 500 to 1000 bp. The product sizes of 800, 900 and 850 ± 20 bp were the most frequent sizes, reported for 24, 20 and 19% of the isolates, respectively. There was no amplification product of the DNA from *S. epidermidis*. As summarized in Table 1, *Nde*I restriction enzyme digestion of the PCR products generated 18 different *Nde*I restriction patterns for all isolates. Except for the products of 500, 600 and 1000 bp, amplicons of the same size generated different patterns, with the number of fragment varying from two to three and molecular sizes from approx. 80 to 330 bp (Fig. 2). Types 9, 11 and 14 were the most common patterns and seen in 50% of the isolates (Table 2). Type 1 was the most frequent pattern and reported in 31.25% of meat product samples, while Type 9 was the most frequent pattern in other foods (50%).

Coagulase protein is a main virulence factor in *S. aureus* [21]. The 3' end of the coagulase gene contains a series of 81-bp tandem repeats, which is different between *S. aureus* strains [22].




Fig. 1. PCR products of *S. aureus coa* coagulase genes isolated from food samples. Lanes 1 and 12, 100-bp ladder; Lane 6, *S. epidermidis* ATCC 12228 negative control; Lane 13, *S. aureus* COL positive control; Lanes 2–5, 7–11, PCR products.

Table 1
NdeI RFLP patterns and frequency of coagulase genotypes of *S. aureus* isolated from food samples.

Type	PCR (bp)	RFLP (bp)	*
1	500	80-170-330	5
2	600	80-250-330	2
3	650 ± 20	80-170-330	1
4	650 ± 20	80-250-330	6
5	700	80-170-330	1
6	700	80-250-330	2
7	750 ± 20	80-170-330	6
8	750 ± 20	80-250-330	2
9	800	80-170-330	22
10	800	80-250-330	2
11	850 ± 20	80-170-330	12
12	850 ± 20	80-250-330	4
13	850 ± 20	80-330	3
14	900	80-170-330	16
15	900	80-250-330	4
16	950 ± 20	80-170-330	6
17	950 ± 20	80-250-330	1
18	1000	80-170-330	5
Total			100

* Distribution of RFLP type in isolates (%).

Classification based on the RFLP of *coa* gene, has been considered as a simple and accurate method for typing *S. aureus* isolated from various sources [17,19]. Using this method, different genotypes were seen among the studied isolates, which

Table 2
Distribution of *NdeI* RFLP patterns and frequencies of coagulase based genotypes of *S. aureus* isolated from food samples.

Origin	n	Genotype																	
		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
Dairy products	78	–	2	1	3	1	2	4	2	18	2	12	3	3	13	2	4	1	5
Meat products	16	5	–	–	3	–	–	1	–	1	–	–	1	–	3	1	1	–	–
Other foods	6	–	–	–	–	–	–	1	–	3	–	–	–	–	–	1	1	–	–




Fig. 2. Electrophoretic profile of *NdeI* restriction fragments of the *coa* gene PCR products. L to R: Lane 1, 100-bp ladder; Lanes 2, 4, 6 and 8, PCR products; Lanes 2, 4, 6 and 8, PCR products; Lane 3, PCR-RFLP Type-1 fragments (80-170-330 bp); Lane 9, PCR-RFLP Type-2 fragments (80-250-330 bp); Lanes 5 and 7, PCR-RFLP Type-13 fragments (80-330 bp).

suggest that *S. aureus* has a certain degree of heterogeneity in the food samples. The ten different PCR types and the 18 *NdeI* RFLP patterns suggest that the isolated *S. aureus* strains harbor more than one variant of the *coa* gene. Results also showed that however different genotypes were detected; only a few predominated. These findings were similar to findings published by researchers who tested milk of bovine mastitis in various countries [12,19,21–23] and showed that several coagulase gene types were responsible for the majority of bovine mastitis cases; only some were predominated in each country. El Bayomi et al. by Genotyping using *spa* PCR-RFLP showed identical restriction banding patterns of MRSA isolates of human and chicken meat origin, indicating the genetic relatedness of the isolates [19]. Aslantas et al. typed 80 *S. aureus* isolated from subclinical bovine mastitis milks and distinguished five PCR and nine RFLP types, with two most common genotypes reported for 73.8% of the isolates [22]. Guler et al. tested 125 *S. aureus* isolates from bovine clinical mastitis in Turkey and found four PCR products of 1000, 900, 800 and 700 bp, with a 1000 bp product being the predominant product [23]. Blaiotta et al. showed in some strains of *Staphylococcus xylosus*, *Staphylococcus saprophyticus*, and *Staphylococcus equorum*, two catalase genes, *katA* and *katB*, were found [24]. Kizerwetter-Swida et al. showed

coagulase-positive staphylococci (CoPS) are opportunistic veterinary pathogens, of which *S. aureus*, *Staphylococcus delphini* and *Staphylococcus intermedius* can be isolated from pigeons. The biochemical identification of *S. delphini* and *S. intermedius* isolates may be incorrect, because of their phenotypic similarity. A total number of 31 isolates of CoPS was obtained, 15 were identified as *S. delphini* group B, six as *S. aureus*, four as *S. delphini* group A, three as *S. intermedius* and three as *Staphylococcus schleiferi* subsp. coagulans. PFGE restriction patterns of *S. delphini* group A and *S. delphini* group B form separate clusters, demonstrating their genetic heterogeneity [25]. Da Silva et al. tested 33 *S. aureus* from milk of goats with clinical and sub-clinical mastitis, belonging to two regions from Brazil and found 11 genotypes. One predominant type was found in each region in most of the isolates [26]. According to Hennekinne et al. Staphylococcal food poisoning (SFP) is one of the most common food-borne diseases caused by the ingestion of staphylococcal enterotoxins (SEs) produced in foods by enterotoxigenic *S. aureus* [27]. Similarly, Moon et al. typed *S. aureus* isolates from animal and vegetable sources in Korea and distinguished 12 genotypes, varied with the source of the microorganisms. However, only a few genotypes prevailed in each source [28]. This distribution might be explained by the coevolution of the hosts and the pathogens, and also differences in reservoirs and imply that the successful transfer of bacteria between bovine mastitis milk, raw meat and vegetables is not a frequent occurrence [22,27].

These characteristics show the necessity of comprehensive studies on epidemiological and ecological profiles of a specific origin before applying food-hygiene control measures.

4. Conclusion

In conclusion, results from the current study showed that however the food samples were contaminated by *S. aureus* harboring more than one coagulase genotype, only a few *coa* genotypes were predominate. Further studies are needed to determine the common characteristics of the predominate strains. This information can be used to better develop control measures for foods contaminated by *S. aureus*.

Acknowledgment

This work was supported by a grant from Vice-Chancellor for Research (grant no. 5615) of Tehran University of Medical Sciences, Tehran, Iran.

References

- [1] E. Shin, H. Hong, J. Park, Y. Oh, J. Jung, Y. Lee, Characterization of *Staphylococcus aureus* fecal isolates associated with food-borne disease in Korea, J. Appl. Microbiol. (2016), <http://dx.doi.org/10.1111/jam.13133> (Epub ahead of print).
- [2] S. Umesh, H.M. Manukumar, Advanced molecular diagnostic techniques for detection of food-borne pathogens; current applications and future challenges, Crit. Rev. Food Sci. Nutr. (2016) (Epub ahead of print).
- [3] F. Rahimi, M. Bouzari, Biochemical fingerprinting of methicillin-resistant *Staphylococcus aureus* isolated from sewage and hospital in Iran, Jundishapur J. Microbiol. 8 (7) (2015) e19760, <http://dx.doi.org/10.5812/jjm.19760v2>, eCollection 2015.
- [4] M. Mazzilli, R. Piccinini, F. Scali, A. Zecconi, Pattern characterization of genes involved in non-specific immune response in *Staphylococcus aureus* isolates from intramammary infections, Res. Vet. Sci. 103 (2015) 54–59, <http://dx.doi.org/10.1016/j.rvsc.2015.09.007> (Epub 2015 Sep 12).
- [5] Z. Aminzadeh, M. Molla Abedin, M.M. Soltan Dallal, L. Gatchkar, Examination of changes of oropharynx flora related to hospitalization, Iran. J. Public Health 32 (2003) 16–19.
- [6] M.M. Soltan Dallal, Z. Salehipour, S. Eshraghi, J. Fallah Mehrabadi, R. Bakhtiari, Occurrence and molecular characterization of *Staphylococcus aureus* strains isolated from meat and dairy products by PCR-RFLP, Ann. Microbiol. 60 (2010) 189–196.
- [7] D. Gutiérrez, S. Delgado, D. Vázquez-Sánchez, B. Martínez, M.L. Cabo, A. Rodríguez, et al., Incidence of *Staphylococcus aureus* and analysis of associated bacterial communities on food industry surfaces, Appl. Environ. Microbiol. 78 (24) (2012) 8547–8554.
- [8] S. Morandi, M. Brasca, R. Lodi, P. Cremonesi, B. Castiglioni, Detection of classical enterotoxins and identification of enterotoxin genes in *Staphylococcus aureus* from milk and dairy products, J. Vet. Microbiol. 124 (2007) 66–72.
- [9] S.K. OH, N. Lee, Y.S. Cho, D.B. Shin, S.Y. Choi, M. Koo, Occurrence of toxigenic *Staphylococcus aureus* in ready-to-eat food in Korea, J. Food Prot. 5 (2006) 1153–1158.
- [10] H. Hasman, A. Moodley, L. Guardabassi, M. Stegger, R.L. Skov, F.M. Aarestrup, Spa type distribution in *Staphylococcus aureus* originating from pigs, cattle and poultry, Vet. Microbiol. 141 (3–4) (2010) 326–331.
- [11] U. Tsegmed, G. Normanno, M. Pringle, K. Krovacek, Occurrence of enterotoxigenic *Staphylococcus aureus* in raw milk from yaks and cattle in Mongolia, J. Food Prot. 70 (2007) 1726–1729.
- [12] M. Kizerwetter-Swida, D. Chrobak-Chmiel, M. Rzewuska, A. Antosiewicz, B. Dolka, A. Ledwon, Genetic characterization of coagulase-positive staphylococci isolated from healthy pigeons, Pol. J. Vet. Sci. 18 (3) (2015) 627–634.
- [13] H. Khoshkaram-Roodmajani, J. Sarvari, A. Bazargani, M.R. Kandekar-Ghahraman, A. Nazari-Alam, M. Motamedifar, Molecular typing of methicillin-resistant and methicillin-susceptible *Staphylococcus aureus* isolates from Shiraz teaching hospitals by PCR-RFLP of coagulase gene, Iran. J. Microbiol. 6 (4) (2014) 246–252.
- [14] C.E. Okolie, K.G. Wooldridge, D.P. Turner, A. Cockayne, R. James, Development of a new pentaplex real-time PCR assay for the identification of poly-microbial specimens containing *Staphylococcus aureus* and other staphylococci, with simultaneous detection of staphylococcal virulence and methicillin resistance markers, Mol. Cell. Probes 29 (3) (2015) 144–150.
- [15] C.J. Shanholtzer, L.R. Peterson, Clinical laboratory evaluation of the thermonuclease test, Am. J. Clin. Pathol. 77 (5) (1982) 587–591.
- [16] M.M. Soltan-Dallal, Z. Salehipour, J. Fallah Mehrabadi, Molecular epidemiology of *Staphylococcus aureus* in food samples based on the protein A gene polymorphic region DNA sequence, Can. J. Microbiol. 56 (1) (2010) 18–21.
- [17] M. Bukowski, K. Polakowska, W.M. Ilcyszyn, A. Sitarska, K. Nytko, M. Kosecka, et al., Species determination within *Staphylococcus* genus by extended PCR-restriction fragment length polymorphism of *saoC* gene, FEMS Microbiol. Lett. 362 (1) (2015) 1–11.
- [18] A. Cosandey, R. Boss, M. Luini, K. Artursson, M. Bardiau, F. Breitenwieser, et al., *Staphylococcus aureus* genotype B and other genotypes isolated from cow milk in European countries, J. Dairy Sci. 99 (1) (2016) 529–540.
- [19] R.M. El Bayomi, H.A. Ahmed, M.A. Awadallah, R.A. Mohsen, A.E. Abd El-Ghafar, M.A. Abdelrahman, Occurrence, virulence factors, antimicrobial resistance, and genotyping of *Staphylococcus aureus* strains isolated from chicken products and humans, Vector Borne Zoonotic Dis. 16 (3) (2016) 157–164.
- [20] O. Raimundo, M. Deighton, J. Capstick, N. Gerraty, Molecular typing of *Staphylococcus aureus* of bovine origin by polymorphisms of the coagulase gene, J. Vet. Microbiol. 66 (1999) 275–284.
- [21] H. Motamedi, S.S. Rahmat Abadi, S.M. Moosavian, M. Torabi, The association of *panT*-*val* leukocidin and *mecA* genes in methicillin-

- resistant *Staphylococcus aureus* isolates from patients referred to educational hospitals in Ahvaz, Iran, Jundishapur J. Microbiol. 8 (8) (2015) e22021.
- [22] O. Aslantas, C. Demir, H. Turutoglu, Z. Cantekin, Y. Ergun, G. Dogruer, Coagulase gene polymorphism of *Staphylococcus aureus* isolated from subclinical bovine mastitis, Turk. J. Vet. Anim. Sci. 31 (2006) 253–257.
- [23] L. Guler, U. Ok, K. Gunduz, Y. Gulcu, H. Hadimli, Antimicrobial susceptibility and coagulase gene typing of *Staphylococcus aureus* isolated from bovine clinical mastitis cases in Turkey, J. Dairy Sci. 88 (2005) 3149–3154.
- [24] G. Blaiotta, V. Fusco, D. Ercolini, O. Pepe, S. Coppola, Diversity of *Staphylococcus* species strains based on partial kat (catalase) gene sequences and design of a PCR-restriction fragment length polymorphism assay for identification and differentiation of coagulase-positive species (*S. aureus*, *S. delphini*, *S. hyicus*, *S. intermedius*, *S. pseudintermedius*, and *S. schleiferi* subsp. *coagulans*), J. Clin. Microbiol. 48 (1) (2010) 192–201.
- [25] M. Kizerwetter-Swida, D. Chrobak-Chmiel, M. Rzewuska, A. Antosiewicz, B. Dolka, A. Ledwon, et al., Genetic characterization of coagulase-positive staphylococci isolated from healthy pigeons, Pol. J. Vet. Sci. 18 (3) (2015) 627–634.
- [26] E.R. Da Silva, J.V.D. Boechat, N. Da Silva, Coagulase gene polymorphism of *Staphylococcus aureus* isolated from goat mastitis in Brazilian dairy herds, Lett. Appl. Microbiol. 42 (2005) 30–34.
- [27] J.A. Hennekinne, M.L. De Buyser, S. Dragacci, *Staphylococcus aureus* and its food poisoning toxins: characterization and outbreak investigation, FEMS Microbiol. Rev. 36 (2012) 815–836.
- [28] J.S. Moon, A.R. Lee, S.H. Jaw, H.M. Kang, Y.S. Joo, Y.H. Park, et al., Comparison of antibiogram, staphylococcal enterotoxin productivity, and coagulase genotypes among *Staphylococcus aureus* isolated from animal and vegetable source in Korea, J. Food Prot. 70 (2007) 2541–2548.