

SCIENTIFIC REPORTS



OPEN

Inactivation Mechanism of *Escherichia coli* Induced by Slightly Acidic Electrolyzed Water

Zhangying Ye¹, Shuo Wang¹, Tao Chen², Weishan Gao¹, Songming Zhu¹, Jinsong He³ & Zhiying Han¹

Foodborne disease outbreak caused by food microbiological contamination is a serious public health problem. Slightly acidic electrolyzed water (SAEW), a new ultra-high effect and wide-spectrum disinfectant that is colourless, odourless, and harmless to humans and the environment, is directly used on food surfaces in Japan and America. However, the underlying inactivation mechanism remains unknown. In this study, biochemical and cellular changes were observed to investigate the bactericidal mechanism of SAEW against *Escherichia coli* (*E. coli*). The results indicated that SAEW with a pH of 6.40, an oxidation-reduction potential (ORP) of 910 mV, an available chlorine concentration (ACC) of 60 mg/L, and a volume ratio of 20:1, produced the most effective sterilization action. A fluorescence-based live-dead assay was further used to demonstrate the sterilized effect and the cell esterase activity damage caused by SAEW. During the observation period, within 10 min, the cell morphology changed, which was characterized by cell expansion, cell elongation and increased membrane permeability. Meanwhile, reactive oxygen substances (ROS) were released in the bacterial cells. *E. coli* inactivation and apoptosis induced by SAEW were observed. Our findings illustrate that the bactericidal effects of SAEW against *E. coli* occurred through cellular and biochemical mechanisms of cell necrosis and apoptosis.

Foodborne diseases have continued to be a widespread and growing health problem, in both developed and developing countries¹. A research study estimated that each year, 31 major pathogens that are acquired in the United States cause 9.4 million episodes of foodborne illness². Developing countries, which bear the brunt of the problem due to the presence of a wide range of foodborne diseases, are even more alarming³. Each year, almost 2.2 million people, mostly children, die from diarrhoeal diseases in developing countries. A considerable proportion of these diseases are probably transmitted through unsafe food⁴. In China, pathogenic microorganisms have been reported to be responsible for over 50% of foodborne illnesses since 2006⁵.

Escherichia coli, which is widely distributed in intestinal environments⁶, have been recognized as a cause of serious clinical illnesses and mortality in foodborne disease outbreaks that involve an enormous variety of foods⁷. As a vital treatment process, effective sterilization and disinfection of food play an important role in controlling disease outbreaks. Chlorine⁸, hydrogen peroxide⁹, ozone¹⁰, and organic acids¹¹ are chemical sanitizers that are frequently used in the food industry to produce high-quality, microbiologically safe food for human consumption. Nevertheless, it is forbidden to use some chemical sanitizers in several European countries because of the potential formation of carcinogenic halogenated disinfection by-products¹², which do harm to the environment as well as to human health. In recent years, there have been many investigations on food decontamination approaches, which have focused on the search for alternative sanitizers based on ensuring food quality and safety¹³.

Slightly acidic electrolyzed water (SAEW) is a novel disinfectant produced by the electrolysis of dilute sodium chloride or hydrochloric acid solutions or both in an electrolytic cell without a separating membrane. Its high bactericidal effect is due to the available chlorine compounds, including ClO⁻, HClO, and Cl₂^{14,15}. A schematic representation of a SAEW generator is shown in Fig. 1.

The main reaction equations are described as below:

Anode side:

¹School of Biosystems Engineering and Food Science, Zhejiang University, Hangzhou, 310058, China. ²School of Environmental & Resource Sciences, Zhejiang University, Hangzhou, 310058, China. ³College of Food Science and Technology, Yunnan Agricultural University, Kunming, 650201, China. Correspondence and requests for materials should be addressed to Z.H. (email: hanzhiying@zju.edu.cn)




Figure 1. Schematic diagram of a slightly acidic electrolyzed water generator



Cathode side:



The main effective chlorine compound form in SAEW (pH 5.0–6.5) is hypochlorous acid (HClO; 97%), which has strong antimicrobial activity¹⁶. As a sanitizer, hypochlorous acid is 80 times more effective than an equivalent concentration of the hypochlorite ion (ClO^-) for inactivating *Escherichia coli*¹⁷. The application of SAEW at a near-neutral pH minimizes human health and safety issues from Cl_2 off-gassing while maximizing the utilization of the hypochlorous acid species¹⁸. Many studies have investigated the significant effectiveness of the antimicrobial activity of SAEW against different foodborne pathogens, including *Listeria monocytogenes*, *E. coli* O157:H7, *Staphylococcus aureus*, *Salmonella typhimurium*, *Campylobacter jejuni* and *Vibrio parahaemolyticus*^{14, 19–22}. Meanwhile, many studies have demonstrated the potential use of SAEW as an alternative sanitizer to reduce microbial contamination on vegetables^{23–25}; cutting boards²⁶; poultry²⁷ and sea foods^{20, 28}. At the same time, SAEW causes less irritation on the human body and less corrosive effects on food preparation appliances^{29, 30}. It was identified as a legal permitted food detergent that can be directly used on food surfaces in Japan and America^{31, 32}.

To date, abundant reports of the antibacterial mechanism of SAEW were focused on the physic and chemical properties on SAEW itself. There have been very few studies on the physiological and biological changes of the bacteria by using SAEW. Biochemical and morphological hallmarks of apoptosis and necrosis are popular in eukaryotic cell. Recently, the characteristic phenomena of apoptosis was also observed in *E. coli* which induced by different kinds of bactericidal antibiotic treatment³³. In this experiment, we devoted to study the cellular and biochemical properties in *E. coli* induced by SAEW, which may illustrate the deep-seated disinfection mechanisms of SAEW.

Results

The viability of *E. coli* was inhibited by SAEW. The physicochemical parameters of SAEW were pH 6.40, an ORP of 900 mV, and an ACC of 60 mg/L in all experiments. The SAEW solution reduced the *E. coli* population levels on LB agar medium, which is shown in Fig. 2A. When the volume ratio of SAEW and bacteria solution was 9:1, the colony counts fell from $8.36 \pm 0.10 \log$ (CFU/mL) to $2.45 \pm 0.07 \log$ (CFU/mL) after being treated by SAEW for 5 minutes, and they fell to $1.33 \pm 0.11 \log$ (CFU/mL) after 10 minutes. Moreover, when the volume ratio of SAEW and bacteria solution was 20:1, the *E. coli* was eliminated after only 5 minutes of SAEW treatment. A higher bactericidal effect was observed at the volume ratio of 20:1 than at the 9:1 ($P < 0.05$, $n = 3$). However, no significant differences were observed between the 5- and 10-minute treatment for the 9:1 and 20:1 volume ratio, respectively ($P > 0.05$).

Colony count was used to estimate the cloning potential of *E. coli* by treating with SAEW. To further elucidate the immediate effect of SAEW on the *E. coli*, MTT assay was used. As shown in Fig. 2B, the OD value was decreased significantly both at the ratio of 9:1 and 20:1. A higher level of *E. coli* inhibition was also observed with




Figure 2. Inhibition of cell proliferation induced by SAEW was determined by colony count and MTT assays. (A) Statistical *E. coli* colony data were measured following 9:1 and 20:1 volume ratios under SAEW treatments at 0, 5 and 10 min. (B) OD absorbance values and (C) cell proliferation were measured following 9:1 and 20:1 volume ratios under treatments at 0, 5 and 10 min (n = 4).

the volume ratio of 20:1 at both 5 ($P = 0.004$) and 10 minutes ($P = 0.01$, Fig. 2C). The results of plate dilution colony counting and MTT colorimetric method showed that with an increased volume ratio, the inhibitory effect on *E. coli* was more effective.

The change of the proportion of living *E. coli* and dead *E. coli* induced by SAEW was detected.

FDA-PI double dye was used to demonstrate the influence process of SAEW on *E. coli* survival state. During the observation duration, the number of cells dyeing immediately with FDA(+), PI(-) decreased gradually (Fig. 3A and B), and dyeing immediately with FDA(+), PI(+) increased gradually by treating with SAEW (Fig. 3A and C). These phenomena showed more obviously by dyeing after incubation. Very few cells were stained with FDA(-), PI(+) (Fig. 3A and D) and FDA(-), PI(-).

SAEW induced cell morphology and permeability changes in the *E. coli*. The above data indicated that treating of SAEW with less than 5 min led to the viability increase and cell membrane integrity destroy. Next, cell morphological changes and cell permeability were studied. As shown in Fig. 4A, compared with the control condition, SAEW induced cell swelling and bloating, with lengths increasing at 1, 5 and 10 min. The relative PI average fluorescence intensity via flow cytometry detection is shown in Fig. 4B. The bacterial cell membrane permeability was enhanced, and the cells were stained by PI. The relative PI average fluorescence intensities of the *E. coli* treated with SAEW were significantly higher than those in the control group (n = 3, $P < 0.05$). The relative average fluorescence intensity was strongest in the 5 min SAEW treated samples (Fig. 4C).

SAEW induced reactive oxygen species production by *E. coli*. The bacterial DCF fluorescence intensity was proportional to the amount of ROS in the cells^{34,35}. As shown in Fig. 5A, compared with the control conditions, the intracellular DCF fluorescence increased 3–4 fold at 1, 5 and 10 min compared with the 0 min group (n = 3, $P < 0.05$; Fig. 5A and B) following SAEW treatment. These observations indicated that SAEW induced ROS production.

***E. coli* exhibit characteristic markers of necrosis and apoptosis induced by SAEW.** Phosphatidylserine (PS) is normally located inside the cell membrane. Externally exposed PS is a typical biochemical marker of apoptosis³⁶, which can be observed by Annexin V-FITC/PI apoptosis detection. Dwyer, D. et al.³³ observed this phenomenon when *E. coli* were treated with antibiotics through Annexin V-FITC/




Figure 3. The live-dead proportion of *E. coli* following SAEW treatment was estimated with a FDA-PI fluorescence assay. (A) The representative data were measured by flow cytometry under treatment of SAEW with the volume ratio of 20:1 at 0, 1, 5 and 10 min. Cell numbers dyeing with (B) FDA (+), (C) FDA-PI(+) and (D) PI(+) following the volume ratio of 20:1 within 0, 1, 5 and 10 min.

PI apoptosis detection. In this study, no obvious fluorescence was observed in control conditions under a fluorescence microscope (As shown in Fig. 6A control). A small proportion of the cells with Annexin V-FITC/PI double dyeing which indicated the apoptosis can be observed (As shown in Fig. 6A yellow arrow). However, most of the cell stained only with PI (red fluorescence) represented that the necrosis or cell damage was the main manifestation induced by SAEW. Figure 6B supplied a little more evidence that SAEW induced apoptosis and necrosis in the bacteria. As shown in Fig. 6B, the apoptosis rate and the necrosis rate increased to around 9% (B2 + B4) and around 15.2% (B2) was induced treatment with SAEW at 10 min, respectively.

Discussion

In this study, the bactericidal effect of SAEW on *E. coli*, which manifested features of apoptosis and necrosis, was estimated. First, the traditional method of counting colony numbers showed that the bactericidal effect was more obvious at the volume ratio of 20:1 (SAEW to bacteria solution) than at the volume ratio of 9:1. The MTT method is not only a popular colorimetric assay for appraising metabolic activity of mammalian cell but can also be used as a quantitative evaluation of viable organisms^{37, 38}. The results also demonstrate highly bactericidal effects of SAEW on *E. coli* within several minutes. It is worth noting that the inhibition of SAEW at the ratio of 20:1 on *E. coli* was incomplete by using MTT assay, while colony formation assay showed a complete elimination. These data manifested that the proliferation potential of a small proportion of *E. coli* by treating with SAEW (20:1) was lost even though they still had mitochondrial succinate dehydrogenase.

Fluorescence-based live-dead assays can be performed with FDA and PI to evaluate viable and dead cells. FDA fluorescence can be converted by esterases that exist in live cells from non-fluorescent FDA. However, PI reaches




Figure 4. SAEW induced *E. coli* morphology and permeability changes. (A) The representative cell morphology data were observed under a microscope. (B) The representative cell permeability data were observed under a fluorescence microscope and by flow cytometry. (C) The statistical data of the relative PI fluorescence intensity were measured by flow cytometry ($n = 4$, $**P < 0.01$ vs 0 min).

into the cell and intercalates with the DNA double helix through disordered or dead cell membranes. In this study, the FDA fluorescence decreased and FDA-PI fluorescence increased significantly but completely in the cells that dyed immediately indicated that instantly effect of SAEW may not cause the *E. coli* dead totally, which correlated with the MTT assay. But the potential for growth was stunted with the observation in the cells that dyed after incubation. The results of FDA-PI double dye indicated that the cell esterase activity and cell membrane integrity were damaged by SAEW application³⁹.

During the bactericidal process, morphological changes often can be induced by antibiotics or disinfectants³³. Here, we found that the *E. coli* elongated and were bloated by SAEW. SAEW could change the *E. coli* morphology but still maintain their cell shape, which indicated that the cells may not disrupt or spilt following SAEW treatment. However, the cell shape existence does not imply the cellular function still existence. We found the cell permeability increased, which means the cell membrane integrity was destroyed. In terms of changes of cell membrane permeability, our postulation was verified when the PI fluorescence increased following SAEW treatment. These phenomena can also be observed with different stimuli, such as heat and high pressure sterilization on bacteria with irreversible loss of membrane integrity, as indicated by PI uptake^{40,41}. Wenwei Tang *et al.*⁴² found that electrochemical oxidizing water (EOW) could strengthen membrane permeability, improve the conductivity of suspension and cause leakage of K^+ and protein out of *Bacillus subtilis* cells. They found that the cell wall and membrane were damaged, which provides a potential explanation for how SAEW may act.

ROS is known to induce apoptotic cell death in various cell types⁴³. The bacterial DCF fluorescence intensity was proportional to the amount of ROS in the cells⁴⁴. In this study, the relative ROS contents reached a maximum value around 5 minutes and maintained to 30 minutes. The relative ROS contents in the *E. coli* increased significantly after treatment with SAEW ($P < 0.05$), which produced free radicals, such as superoxide anion ($O_2^{\cdot-}$), hydrogen peroxide (H_2O_2), and hydroxyl radical ($\cdot OH$) that could oxidize the lipids, glycolipids, and proteins in the cytomembrane into peroxides^{45,46}. Jeong, J. *et al.*⁴⁷ examined the role of ROS in electrochemical disinfection, and they found that $\cdot OH$ was the major lethal species responsible for *E. coli* inactivation in the chloride-free electrochemical disinfection process.

Annexin V-FITC/PI double dye fluorescence is often used to quantify apoptosis⁴⁸. In this study, under a fluorescence microscope, the green fluorescent label by Annexin V-FITC inferred early apoptosis, and the red fluorescent label by PI inferred late apoptosis. The flow cytometry results also demonstrated that SAEW induced apoptotic characteristics in the *E. coli* cells. These results are similar to the conclusions of a study reported by Dwyer, D. *et al.*³³ SAEW damaged bacteria rather than causing immediate death.




Figure 5. SAEW induced ROS production in *E. coli*, as measured by flow cytometry. (A) The representative and (B) the statistical DCF fluorescence data were observed following SAEW treatment with the volume ratio of 20:1 ($n = 4$, ** $P < 0.01$ vs 0 min).

Conclusion

In a whole, SAEW showed an effective bactericidal effect on the *E. coli* which associated with the physiological and biological changes of apoptosis and necrosis induced by SAEW. Our investigation may further illustrate the disinfection mechanisms and expand the applications of SAEW.

Materials and Methods

Preparation and physicochemical property measurements of SAEW. SAEW was produced by electrolysis of 6% HCl solution using a flow-type electrolysis apparatus equipped with an electrolytic cell without a separating membrane between the anode and cathode at a voltage of 220 V (HD-240L, Wangpu Trading Co., Ltd, Shanghai, China). The SAEW was prepared on the day of the experiments and used within 1 h of production. The physicochemical properties of SAEW were determined immediately after generation. The pH and ORP were determined with a multi-functional pH/ORP meter (LE-438, METTLER TOLEDO, Shanghai, China) bearing a pH electrode and an ORP electrode. The available chlorine concentration was evaluated by the standard iodometric titration method⁴⁹.

Preparation of bacterial suspension. The strain, *Escherichia coli* ATCC25922, was obtained from the China General Microbiological Culture Collection Center (CGMCC). Prior to each experiment, The stock cultures were transferred into 100 mL nutrient broth (Huankai Microbial Sci. & Tech. Co., Ltd., Guangdong, China) and incubated in an incubator shaker (SHA-CA, Honghua Instrument Factory, Jintan, China) for 24 h at 150 r/min, 37 °C. A 5-ml volume of the enriched culture was pooled into sterile centrifuge tube and subsided in a refrigerated centrifuge (5424R, Eppendorf China Ltd., Shanghai, China) at 5000 rpm, 4 °C for 10 min. The resulting cell pellet was washed twice and resuspended in 5 mL of sterile phosphate buffer solution (PBS) to obtain a final bacterial cell density of approximately 10^9 CFU/mL which was confirmed by plating 1 mL portions of appropriately diluted *E. coli* suspension on plate count agar plates and then incubated at 37 °C for 24 h.

Effect of SAEW on microbiota inactivation. The SAEW (pH: 6.40, ORP: 910 mV, ACC: 60 mg/L) used in this study was generated by electrolysis. In SAEW treatment experiments, SAEW and *E. coli* cell suspensions that were prepared as previously described were mixed at ratios of 9:1 and 20:1 for 5 and 10 min. At the end of each treatment time, 1 ml of each sample was transferred to a sterile tube containing 9 ml neutralizing buffer solution (0.5% $\text{Na}_2\text{S}_2\text{O}_3$ containing 0.03 M phosphate buffer solution) to halt disinfection by SAEW. After 5 min of neutralization, treated and control (untreated) samples were then serially diluted in sterile PBS. Subsequently, viable colonies were enumerated by the plate dilution colony counting and MTT colorimetric methods (Sigma, America).




Figure 6. SAEW induced *E. coli* necrosis and apoptosis, as measured with fluorescence microscopy and flow cytometry. (A) The representative data of the FITC-PI label were observed with a fluorescence microscope and (B) flow cytometry.

MTT (3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide) colorimetric method. MTT that is water soluble, is converted to insoluble formazan by cleavage of the tetrazolium ring by succinate dehydrogenase in the presence of live cells⁵⁰. Since formazan is purple, the change in colour of the solution provides an indication of cellular activity. MTT allows for cellular activity to be quantified by measuring the absorbance of the solution at certain wavelengths^{33, 37, 38}. After treatment with SAEW (and PBS as a blank control), samples were centrifuged at 5000 rpm for 5 minutes in a refrigerated centrifuge (5424 R, Eppendorf China Ltd., Shanghai, China). The supernatant was removed and then re-suspended in 1 mL of sterile broth. Bacteria liquid was added into 24-well culture plates (100 μ L of each sample per well). Each group was established in three holes. Ten microliters of MTT was added to each well, which was then incubated at 37 °C for 4 h without light. Then, 100 μ L of dimethyl sulfoxide (DMSO) was added to each well, and the bacterial liquid was vibrated in an incubator shaker for 10 minutes until the crystalline DMSO dissolved. Afterwards, the absorbance of each sample was measured at an OD of 570 nm with an ELISA instrument. Inhibition rate of *E. coli* proliferation was calculated by OD values.

Determination of *E. coli* morphologic change. Two hundred microlitres of bacterium suspension was drizzled in the polylysine coated slides in 6-well plates. Then, the suspension was submerged in 2 mL of sterile broth at 37 °C for 3 h to make it stick on the glass for easy observation. The excess broth was then removed, and 4 mL of SAEW (PBS as a blank control) was added to the bacteria for the pre-set times. At each timepoint, approximately 10^6 cells were collected. The cells were washed once and observed under a fluorescence microscope (ECLIPSE Ti-s, Nikon, Japan).

Determination of *E. coli* reactive oxygen species content. Reactive Oxygen Detection Kits (Blue skies biotechnology, China) were used for detection of *E. coli* ROS levels by fluorescence probe (DCFH-DA, (2,7-Dichlorodi-hydrofluorescein diacetate)). The fluorescence intensity was proportional to the levels of ROS. After treatment with SAEW (PBS as blank control) for pre-set times, the samples were centrifuged at 5000 rpm for 5 minutes in a refrigerated centrifuge. They were then washed once and resuspended in 200 μ L of DCFH-DA working solution and incubated at 37 °C in the dark for 20 minutes. To detect the ROS contents by BD flow cytometry (GALLIOS, BECKMAN COULTER, USA), approximately 10^6 cells were collected for each sample.

Fluorescein diacetate (FDA)-propidium iodide (PI) double dye analysis to determine the influence of SAEW on cell activity. The influence of SAEW on cell activity was detected by FDA (West Asia reagent, China) and PI (Blue skies biotechnology, China). FDA is a non-polar, hydrophobic, non-fluorescent esterified compound, which readily permeates the cell membrane and is hydrolyzed by non-specific esterases producing a fluorescein. FDA was used to indicate the presence of active esterase. Meanwhile, PI was utilized to assess cell membrane integrity. Cells with an intact cell membrane and inactive esterase cannot be stained with FDA or PI. After treatment with SAEW (PBS as a blank control) for pre-set times, samples were centrifuged at 5000 rpm for 5 minutes in a refrigerated centrifuge and washed once. Then, the samples were divided into two groups: those that were dyed immediately and those that were dyed after a 4-h incubation. In the immediately dyed group, cells were resuspended in one hundred microlitres of binding buffer. Ten microlitres of FDA working solution and 2.5 μL of PI were added to each sample, which were then incubated in the dark for 10 minutes. In the group that was dyed after the 4-h incubation, cells were resuspended in 1 mL of sterile broth and incubated for 4 h. Then, the samples were centrifuged at 5000 rpm for 5 minutes in a refrigerated centrifuge and washed once before the staining step. Approximately 10^6 cells were collected for each sample. The treated cells were detected with a BD flow cytometer.

PI single-staining to analyse the influence of SAEW on membrane permeability. Two kinds of detection methods to determine PI single-staining were utilized: fluorescence microscope observation and flow cytometry.

Fluorescence microscope observation. Two hundred microlitres bacterium suspension was drizzled onto poly-lysine coated slides in 6-well plates. Then, the bacterial suspension was submerged in 2 mL of sterile broth at 37°C for 3 h so that it would stick on the glass for easy observation. Then, the excess broth was removed, and 4 mL of SAEW (PBS as a blank control) was added to the bacteria for the pre-set times. The cells were washed once, and then 100 μL of PI diluent (100 μL of binding buffer + 2.5 μL of PI) was added to each sample, which were then incubated in the dark for 10 minutes at room temperature. At each timepoint, approximately 10^6 cells were collected. The cells were washed once and observed under a fluorescence microscope.

Flow cytometry. After treatment with SAEW (PBS as a blank control) for the pre-set times, the samples were centrifuged at 5000 rpm for 5 minutes in a refrigerated centrifuge, washed once and resuspended in 100 μL of binding buffer. Then, 2.5 μL of PI was added to each sample, which were then incubated in the dark for 10 minutes. Approximately 10^6 cells were collected for each sample. The treated cells were detected with a BD flow cytometer.

Annexin V-FITC/PI apoptosis detection. Apoptosis was detected with an Annexin V-FITC/PI apoptosis kit (Blue skies biotechnology, China). Annexin V is a Ca^{2+} -dependent phospholipid-binding protein with high affinity for phosphatidylserine (PS). PS externalization is a relatively early event in apoptosis and occurs before plasma membrane integrity. Therefore Annexin V labeled with fluorescein isothiocyanate (FITC) can be used as a sensitive probe for PS exposure upon the cell membrane. In late apoptotic cells and dead cells, PI is able to pass through the cell membrane to make the cell stained. Thus cells can be distinguished at different stages of apoptosis through Annexin V-FITC/PI apoptosis kits. Furthermore, two kinds of detection methods were utilized to determine apoptosis: fluorescence microscope observations and flow cytometry.

The method to affix the bacteria to slides and treat with SAEW was in accordance with the operating procedure of the *E. coli* morphologic change determinations as mentioned above. The cells were washed once, and then 100 μL of Annexin V-FITC diluent (100 μL of binding buffer + 5 μL of Annexin V-FITC) was added to each sample, which was then incubated in the dark for 20 minutes at room temperature. The cells were washed once followed by the addition of 100 μL of PI diluent (100 μL of binding buffer + 2.5 μL of PI). At each timepoint, approximately 10^6 cells were collected. The cells were washed again and then observed under a fluorescence microscope.

Flow cytometry: After treatment with SAEW (PBS as a blank control) for pre-set times, the samples were centrifuged at 5000 rpm for 5 minutes in a refrigerated centrifuge, washed once and resuspended in 100 μL of binding buffer. Ten microlitres of Annexin V-FITC was added to each sample. Then, 2.5 μL of PI was added to each sample, which was then incubated in the dark for 10 minutes. Approximately 10^6 cells were collected for each sample. The treated cells were detected with a BD flow cytometer.

Statistical analysis. For each treatment, data from independent replicate trials were pooled. The flow cytometry data were analysed with CellQuest software. Data integration and drawing processes were conducted with the Origin Version 8 software (OriginLab Corp., USA). One-way analysis of variance (ANOVA) and Tukey's test for independent replicates were performed using SPSS 19.0 (Statistical Package for the Social Sciences; SPSS, Inc, Chicago, IL) software. Significant differences were defined at $P = 0.05$. Data were expressed as means \pm standard errors.

References

1. World Health Organization. Global Foodborne Infections Network (GFN) (2011).
2. Scallan, E. *et al.* Foodborne illness acquired in the united states-major pathogens. *Emerging Infectious Diseases* **17**, 7–15 (2011).
3. World Health Organization. Food safety and foodborne illness. *Nutrition Newsletter* (2003).
4. Kuchenmüller, T., Abela-Ridder, B., Corrigan, T. & Tritscher, A. World Health Organization initiative to estimate the global burden of foodborne diseases. *Revue Scientifique Et Technique* **32**, 459–467 (2013).
5. Xue, J. & Zhang, W. Understanding China's food safety problem: An analysis of 2387 incidents of acute foodborne illness. *Food Control* **30**, 311–317 (2013).

6. Joseph, S. W., Ingram, D. T. & Kaper, J. B. The epidemiology, pathogenicity and microbiology of foodborne *Escherichia coli* O157:H7. *Reviews in Medical Microbiology* **13**, 53–62 (2002).
7. Bell, C. Approach to the control of enterohaemorrhagic *Escherichia coli* (EHEC). *International Journal of Food Microbiology* **78**, 197–216 (2002).
8. Tomás-Callejas, A. *et al.* Chlorine dioxide and chlorine effectiveness to prevent *Escherichia coli* O157: H7 and *Salmonella* cross-contamination on fresh-cut red chard. *Food Control* **23**, 325–332 (2012).
9. Alexander, E. M. C., Brandao, T. R. S. & Silva, C. L. M. Assessment of the impact of hydrogen peroxide solutions on microbial loads and quality factors of red bell peppers, strawberries and watercress. *Food Control* **27**, 362–368 (2012).
10. Alexandre, E. M. C., Santos-Pedro, D. M., Brandao, T. R. S. & Silva, C. L. M. Influence of aqueous ozone, blanching and combined treatments on microbial load of red bell peppers, strawberries and watercress. *Journal of Food Engineering* **105**, 277–282 (2011).
11. Huang, Y. X. & Chen, H. Q. Effect of organic acids, hydrogen peroxide and mild heat on inactivation of *Escherichia coli* O157: H7 on baby spinach. *Food Control* **22**, 1178–1183 (2011).
12. Ölmez, H. & Kretzschmar, U. Potential alternative disinfection methods for organic fresh-cut industry for minimizing water consumption and environmental impact. *LWT-Food Science and Technology* **42**, 686–693 (2009).
13. Mansur, A. R. & Oh, D. H. Combined effects of thermosonication and slightly acidic electrolyzed water on the microbial quality and shelf life extension of fresh-cut kale during refrigeration storage. *Food Microbiology* **51**, 154–162 (2015).
14. Cao, W., Zhu, Z., Shi, Z., Wang, C. & Li, B. Efficiency of slightly acidic electrolyzed water for inactivation of *Salmonella enteritidis* and its contaminated shell eggs. *International Journal of Food Microbiology* **130**, 88–93 (2009).
15. Zheng, W. *et al.* Bactericidal activity of slightly acidic electrolyzed water produced by different methods analyzed with ultraviolet spectrophotometric. *International Journal of Food Engineering* **8**, Article 41 (2012).
16. Yoshifumi, H. Improvement of the electrolysis equipment and application of slightly acidic electrolyzed water for dairy farming. *Journal of the Japanese Society of Agricultural Machinery* **65**, 27–29 (2003).
17. Tango, C. N., Mansur, A. R., Kim, G. H. & Oh, D. H. Synergetic effect of combined fumaric acid and slightly acidic electrolysed water on the inactivation of food-borne pathogens and extending the shelf life of fresh beef. *Journal of Applied Microbiology* **117**, 1709–1720 (2014).
18. Guentzel, J. L., Lam, K. L., Callan, M. A., Emmons, S. A. & Dunham, V. L. Reduction of bacteria on spinach, lettuce, and surfaces in food service areas using neutral electrolyzed oxidizing water. *Food Microbiology* **25**, 36–41 (2008).
19. Al-Holy, M. A. & Rasco, B. A. The bactericidal activity of acidic electrolyzed oxidizing water against *Escherichia coli* O157:H7, *Salmonella Typhimurium*, and *Listeria monocytogenes* on raw fish, chicken and beef surfaces. *Food Control* **54**, 317–321 (2015).
20. McCarthy, S. & Burkhardt, W. Efficacy of electrolyzed oxidizing water against *Listeria monocytogenes* and *Morganella morganii* on conveyor belt and raw fish surfaces. *Food Control* **24**, 214–219 (2012).
21. Park, C. M., Hung, Y. C. & Brackett, R. E. Antimicrobial effect of electrolyzed water for inactivating *Campylobacter jejuni* during poultry washing. *International Journal of Food Microbiology* **72**, 77–83 (2002).
22. Wang, J. J. *et al.* Fate of *Vibrio parahaemolyticus* on shrimp after acidic electrolyzed water treatment. *International Journal of Food Microbiology* **179**, 50–56 (2014).
23. Issa-Zacharia, A., Kamitani, Y., Miwa, N., Muhimbula, H. & Iwasaki, K. Application of slightly acidic electrolyzed water as a potential non-thermal food sanitizer for decontamination of fresh ready-to-eat vegetables and sprouts. *Food Control* **22**, 601–607 (2011).
24. Koseki, S., Yoshida, K., Isobe, S. & Itoh, K. Decontamination of lettuce using acidic electrolyzed water. *Journal of Food Protection* **64**, 652 (2001).
25. Forghani, F. & Oh, D. H. Hurdle enhancement of slightly acidic electrolyzed water antimicrobial efficacy on Chinese cabbage, lettuce, sesame leaf and spinach using ultrasonication and water wash. *Food Microbiology* **36**, 40–5 (2013).
26. Venkitanarayanan, K. S., Ezeike, G. O. I., Hung, Y. C. & Doyle, M. P. Inactivation of *E. coli* O157:H7 and *L. monocytogenes* on plastic kitchen cutting boards by electrolyzed oxidizing water. *Journal of Food Protection* **62**, 857–860 (1999).
27. Mansur, A. R., Tango, C. N., Kim, G. H. & Oh, D. H. Combined effects of slightly acidic electrolyzed water and fumaric acid on the reduction of foodborne pathogens and shelf life extension of fresh pork. *Food Control* **47**, 277–284 (2015).
28. Kim, W. T. *et al.* Use of electrolyzed water ice for preserving freshness of pacific saury (*Cololabis saira*). *Journal of Food Protection* **69**, 2199–204 (2006).
29. Hricova, D., Stephan, R. & Zweifel, C. Electrolyzed water and its application in the food industry. *Journal of Food Protection* **71**, 1934–1947 (2008).
30. Eda, H. & Seckin, A. M. The effect of different electrolyzed water treatments on the quality and sensory attributes of sweet cherry during passive atmosphere packaging storage. *Postharvest Biology & Technology* **102**, 32–41 (2015).
31. Katayose, M., Yoshida, K., Achiwa, N. & Eguchi, M. Safety of electrolyzed seawater for use in aquaculture. *Aquaculture* **264**, 119–129 (2007).
32. Yoshida, K., Achiwa, N. & Katayose, M. Application of electrolyzed water for food industry in Japan. Las Vegas: Proceedings of 2004 Institute Food Technologist Annual Meeting. (2004).
33. Dwyer, D., Camacho, D., Kohanski, M., Callura, J. & Collins, J. Antibiotic-induced bacterial cell death exhibits physiological and biochemical hallmarks of apoptosis. *Molecular Cell* **46**, 561–572 (2012).
34. Kim, J. A., Lau, E. K., Pan, L. & De Blanco, E. J. NF- κ B inhibitors from *Brucea javanica* exhibiting intracellular effects on reactive oxygen species. *Anticancer Research* **30**, 3295–3000 (2010).
35. Liu, X. *et al.* Determination of a threshold dose to reduce or eliminate CdTe-induced toxicity in L929 cells by controlling the exposure dose. *PLOS ONE* **8**(4), e59359 (2013).
36. Fadok, V. A. *et al.* Exposure of phosphatidylserine on the surface of apoptotic lymphocytes triggers specific recognition and removal by macrophages. *Journal of Immunology* **148**, 2207–2216 (1992).
37. Freimoser, F. M., Jakob, C. A., Aebi, M. & Tuor, U. The MTT [3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide] assay is a fast and reliable method for colorimetric determination of fungal cell densities. *Applied & Environmental Microbiology* **65**, 3727–3729 (1999).
38. Montoro, E. *et al.* Comparative evaluation of the nitrate reduction assay, the MTT test, and the resazurin microtitre assay for drug susceptibility testing of clinical isolates of *Mycobacterium tuberculosis*. *Journal of Antimicrobial Chemotherapy* **55**, 500–505 (2005).
39. Shuona Chen *et al.* Influence of co-existed benzo[a]pyrene and copper on the cellular characteristics of *Stenotrophomonas maltophilia* during biodegradation and transformation. *Bioresource Technology* **158**, 181–187 (2014).
40. Ananta, E. *et al.* Assessment of high pressure induced damage on *Lactobacillus rhamnosus* GG by flow cytometry. *Food Microbiology* **21**, 567–577 (2004).
41. Klotz, B., Mañas, P. & Mackey, B. M. The relationship between membrane damage, release of protein and loss of viability in *Escherichia coli* exposed to high hydrostatic pressure. *International Journal of Food Microbiology* **137**, 214–220 (2010).
42. Wenwei Tang *et al.* Disinfection effect and its mechanism of electrolyzed oxidizing water on spores of *Bacillus subtilis* var. *niger*. *Food Science and Biotechnology* **20**, 889–895 (2011).
43. Dimmeler, S. & Zeiher, A. M. Reactive oxygen species and vascular cell apoptosis in response to angiotensin II and pro-atherosclerotic factors. *Regulatory Peptides* **90**, 19–25 (2000).
44. Ohashi, T. *et al.* Rapid oxidation of dichlorodihydrofluorescein with heme and hemoproteins: formation of the fluorescein is independent of the generation of reactive oxygen species. *FEBS Letters* **511**, 21–27 (2002).

45. Halliwell, B. & Gutteridge, J. M. C. Free radicals in biology and medicine, third edition. *Free Radical Biology and Medicine* **10**, 449–450 (1999).
46. Voziyan, P. A. & Yazlovitskaya, E. M. Reactive Oxygen Species. *J Bioequiv Availab* **6**, e57 (2014).
47. Jeong, J., Kim, J. Y. & Yoon, J. The role of reactive oxygen species in the electrochemical inactivation of microorganisms. *Environmental Science & Technology* **40**, 6117–6122 (2006).
48. Olszowski, T. *et al.* The effects of cadmium at low environmental concentrations on THP-1 macrophage apoptosis. *International Journal of Molecular Sciences* **16**, 21410–27 (2015).
49. APHA, AWWA, WEF. Standard methods for the examination of water and wastewater. 22nd ed. American Public Health Association, American Water Works Association and Water Environment Federation, Washington, DC (2012).
50. Gerlier, D. & Thomasset, N. Use of MTT colorimetric assay to measure cell activation. *Journal of Immunological Methods* **94**, 57–63 (1986).

Acknowledgements

This research was financially supported by the National Key R&D Program of China (Project number: 2017YFD0701703), ZHEJIANG Province Key R&D Project (Project number: 2015C02010) and the Key Laboratory of Equipment and Informatization in Environment Controlled Agriculture, Ministry of Agriculture, P.R. China. Any opinions, findings, and conclusions expressed in this publication are those of the authors and do not necessarily reflect the views of Zhejiang University.

Author Contributions

Zhangying Ye and Zhiying Han conceived the experiments. Zhangying Ye, Shuo Wang and Tao Chen conducted the experiments. Weishan Gao and Tao Chen analysed the data. Songming Zhu and Jinsong He provided technical experience. Zhangying Ye, Shuo Wang and Zhiying Han wrote the manuscript. All authors reviewed the manuscript.

Additional Information

Competing Interests: The authors declare that they have no competing interests.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2017