

**Foodborne illness source attribution
estimates for 2013 for *Salmonella*,
Escherichia coli O157,
Listeria monocytogenes, and
Campylobacter using multi-year outbreak
surveillance data,
United States**

**The Interagency Food Safety Analytics Collaboration
(IFSAC)**

December 2017



Acknowledgments

The methods used for this report were developed by members of the Interagency Food Safety Analytics Collaboration (IFSAC), which includes the Centers for Disease Control and Prevention (CDC), the U.S. Food and Drug Administration (FDA), and the U.S. Department of Agriculture's Food Safety and Inspection Service (USDA-FSIS).

The following persons (listed alphabetically) contributed substantially to this report:

CDC

Beau B. Bruce, MD, PhD
R. Michael Hoekstra, PhD
LaTonia Richardson, PhD

FDA

Michael Batz, MSc
Michael Bazaco, PhD
Cary Chen Parker, MPH
Christopher Waldrop, MPH

FSIS

Kristin Holt, PhD
Gabrielle Johnston, MPH
Joanna Zablotsky-Kufel, PhD, MPH

IFSAC Steering Committee:

CDC

Patricia M. Griffin, MD
Robert V. Tauxe, MD, MPH

FDA

Katherine Vierk Brill, MPH
Sherri McGarry, MS
Beverly Wolpert, PhD, MS

FSIS

Christopher Alvares, MS
David Goldman, MD, MPH

Suggested Citation

Interagency Food Safety Analytics Collaboration. Foodborne illness source attribution estimates for 2013 for *Salmonella*, *Escherichia coli* O157, *Listeria monocytogenes*, and *Campylobacter* using multi-year outbreak surveillance data, United States. GA and D.C.: U.S. Department of Health and Human Services, CDC, FDA, USDA-FSIS. 2017.

Executive Summary

Each year in the United States an estimated 9 million people get sick, 56,000 are hospitalized, and 1,300 die of foodborne disease caused by known pathogens. These estimates help us understand the scope of this public health problem. However, to develop effective prevention measures, we need to understand the types of foods contributing to the problem.

The Interagency Food Safety Analytics Collaboration (IFSAC) is a tri-agency group created by the Centers for Disease Control and Prevention (CDC), the U.S. Food and Drug Administration (FDA), and the U.S. Department of Agriculture's Food Safety and Inspection Service (USDA-FSIS). Based on a new method, IFSAC developed foodborne illness source attribution estimates for 2012 using outbreak data from 1998 through 2012 for four priority pathogens, *Salmonella*, *Escherichia coli* O157, *Listeria monocytogenes*, and *Campylobacter*. IFSAC described this method and the estimates for 2012 in a [report](#) and at a [public meeting](#).

IFSAC derived the estimates for 2013 using the same method used for the 2012 estimates, with some modifications. The data came from 1,043 foodborne disease outbreaks that occurred from 1998 through 2013 and for which each confirmed or suspected implicated food fell into a single food category. The method relies most heavily on the most recent five years of outbreak data (2009–2013). Foods are categorized using a scheme IFSAC created to classify foods into 17 categories that closely align with the U.S. food regulatory agencies' classification needs.

***Salmonella* illnesses came from a wide variety of foods.**

Salmonella illnesses were broadly attributed across multiple food categories. More than 75% of *Salmonella* illnesses were attributed to seven food categories: Seeded Vegetables (such as tomatoes), Eggs, Chicken, Other Produce (such as nuts), Pork, Beef, and Fruits.

***E. coli* O157 illnesses were most often linked to Vegetable Row Crops (such as leafy greens) and Beef.** More than 75% of illnesses were linked to these two categories.

***Listeria monocytogenes* illnesses were most often linked to Fruits and Dairy products.**

More than 75% of illnesses were attributed to these two categories, but the rarity of *Listeria monocytogenes* outbreaks makes these estimates less reliable than those for other pathogens.

Non-Dairy *Campylobacter* illnesses were most often linked to Chicken.

Almost 80% of non-Dairy foodborne illnesses were attributed to Chicken, Other Seafood (such as shellfish), Seeded Vegetables, Vegetable Row Crops, and Other Meat/Poultry (such as lamb or duck). An attribution percentage for Dairy is not included because, among other reasons, most foodborne *Campylobacter* outbreaks were associated with unpasteurized milk, which is not widely consumed, and we think these over-represent Dairy as a source of *Campylobacter* illness. Removing Dairy illnesses from the calculations highlights important sources of illness from widely consumed foods, such as Chicken.

This collaborative effort to provide annual attribution estimates continues IFSAC's work to improve foodborne illness source attribution, which can help inform efforts to prioritize food safety initiatives, interventions, and policies for reducing foodborne illnesses. These consensus estimates allow all three agencies to take a consistent approach to identifying food safety priorities to protect public health. For more information on IFSAC projects visit <http://www.cdc.gov/foodsafety/ifsac/index.html>.

Introduction

Each year in the United States, an estimated 9 million people get sick, 56,000 are hospitalized, and 1,300 die of foodborne disease caused by known pathogens—these estimates help us understand the scope of this public health problem.¹ However, to develop effective prevention measures, we need to understand the percentage of foodborne illnesses associated with specific foods; we call this work foodborne illness source attribution.

With the creation of the Interagency Food Safety Analytics Collaboration (IFSAC) in 2011, the Centers for Disease Control and Prevention (CDC), the U.S. Food and Drug Administration (FDA), and the U.S. Department of Agriculture's Food Safety and Inspection Service (USDA-FSIS) agreed to improve data and methods used to estimate foodborne illness source attribution and provide timely estimates of the food sources of four priority foodborne pathogens: *Salmonella*, *Escherichia coli* O157, *Listeria monocytogenes* (*Lm*), and *Campylobacter*. IFSAC considers these priority pathogens because of the frequency (estimated 1.9 million illnesses each year) and severity of illness they cause, and because targeted interventions can significantly reduce these illnesses.

IFSAC developed a method for analyzing outbreak data to estimate which foods are responsible for illnesses related to the four priority pathogens, using a scheme IFSAC created to classify foods into 17 categories that closely align with the U.S. food regulatory agencies' classification needs.² IFSAC described this method and the resulting estimates for the year 2012 in a report³ and at a public meeting.⁴ IFSAC derived the estimates for 2013 using the same method, with some modifications. IFSAC plans to provide annual estimates of the sources of foodborne illness for the priority pathogens while continuing to work on methods to further improve these estimates.

Consensus among CDC and the regulatory agencies on methods and attribution estimates can help in efforts to prioritize food safety initiatives, interventions, and policies for reducing foodborne illnesses. The 2012 and 2013 estimates achieve IFSAC's goals of using improved methods to develop estimates of foodborne illness source attribution for priority pathogens and of achieving consensus that these are the best current estimates for the agencies to use in their food safety activities. These estimates can also help scientists; federal, state, and local policy-makers; the food industry; consumer advocacy groups; and the public assess whether prevention measures are working.

Methods

We analyzed data extracted from CDC's Foodborne Disease Outbreak Surveillance System (FDOSS) ^{5,6} (www.cdc.gov/foodsafety/fdoss) on outbreaks that were confirmed or suspected to be caused by the four priority pathogens from 1998 through 2013. We excluded outbreaks that occurred in a U.S. territory, were caused by multiple etiologies, or had no identified food vehicle or contaminated ingredient.

Each outbreak was assigned to a single food category using the IFSAC food categorization scheme² based on confirmed or suspected implicated foods and ingredients (i.e., a single ingredient was confirmed or suspected to be implicated or all ingredients in the food were assigned to the same food category). We excluded outbreaks that could not be assigned to a single food category, usually because the food was complex (i.e., composed of ingredients belonging to more than one category) or the contaminated ingredient could not be identified.

We developed pathogen-specific analysis of variance (ANOVA) models using our previously described method³ to mitigate the impact of large outbreaks and control for epidemiological factors. We estimated the number of log-transformed illnesses associated with each outbreak based on three factors deemed to be important based on our exploratory analyses: food category, type of preparation location (e.g., restaurant, home), and whether the outbreak occurred in one or more states.

These model estimates were then back-transformed and down-weighted with a function that declines exponentially for outbreaks older than the most recent five years (2009–2013) because we considered foods more recently implicated to be most relevant for estimating current attribution.

We used the resulting down-weighted model-estimated illnesses to calculate each estimated attribution percentage: the sum of illnesses associated with a pathogen-food category pair was divided by the sum of illnesses associated with that pathogen across all food categories. After down-weighting, 67% of overall information came from the most recent five years, 28% from the next most recent five years (2004–2008), and 5% from the oldest data (1998–2003). Among the 176 *Campylobacter* outbreaks during the study period, the 116 assigned to Dairy were excluded from the final calculation of the attribution percentage due to the challenges of estimating *Campylobacter* attribution using outbreak data. Thus, the *Campylobacter* attribution point estimates primarily reflect data from 60 outbreaks.

In the graphs and tables, food categories appear in descending order of their estimated attribution percentage, and those that contributed to a cumulative attribution of at least 75% of illnesses are indicated.


Results

We identified 2,919 outbreaks that occurred from 1998 through 2013 and that were confirmed or suspected to be caused by *Salmonella*, *E. coli* O157, *Lm*, or *Campylobacter*. Of these, we excluded 88 outbreaks with multiple confirmed or suspected etiologies. We further excluded 1,083 outbreaks without a confirmed or suspected implicated food, 702 outbreaks for which the food vehicle could not be assigned to one of the 17 food categories, and three that occurred in a U.S. territory.

The resulting dataset included 1,043 outbreaks in which the confirmed or suspected implicated food or foods could be assigned to a single food category: 638 caused or suspected to be caused by *Salmonella*, 203 by *E. coli* O157, 26 by *Lm*, and 176 by *Campylobacter*. Due to down-weighting, the last five years of outbreaks provide the majority of information for the estimates; outbreaks from 2009 through 2013 provide 71% of model-estimated illnesses used to calculate attribution for *Salmonella*, 68% for *E. coli* O157, 85% for *Lm*, and 63% for *Campylobacter*.

The overall results and those for each pathogen are shown in Figures 1 through 5.

Figure 1: Estimated percentage (with 90% credibility intervals) for 2013 of *Salmonella*, *Escherichia coli* O157, *Listeria monocytogenes*, and *Campylobacter* illnesses attributed to 17 food categories, based on multi-year outbreak data,*† United States. Download data tables for graph at <https://www.cdc.gov/foodsafety/ifsac/files/P19Fig1Data.csv>




*Based on a model using outbreak data that gives equal weight to each of the most recent five years of data (2009–2013), and exponentially less weight to each earlier year (1998–2008).

† *Campylobacter* estimates exclude results derived from Dairy outbreak data.

Overall Key Results

- The results are based on 638 outbreaks caused or suspected to be caused by *Salmonella*, 203 by *E. coli* O157, 26 by *Lm*, and 60 by *Campylobacter* (after outbreaks due to Dairy were excluded).
- Estimated *Salmonella* and *Campylobacter* illnesses were more widely distributed across food categories than illnesses from *E. coli* O157 and *Lm*; most of the illnesses for the latter two pathogens were attributed to two food categories.
- For most pathogens, the credibility intervals overlap for categories with the highest attribution percentages, indicating no statistically significant difference among them.

Figure 2: Estimated percentage of *Salmonella* illnesses (with 90% credibility intervals) for 2013, in descending order, attributed to each of 17 food categories, based on multi-year outbreak data,*United States




*Based on a model using outbreak data that gives equal weight to each of the most recent five years of data (2009–2013) and exponentially less weight to each earlier year (1998–2008).

Salmonella Key Results

- Over 75% of illnesses were attributed to seven food categories: Seeded Vegetables (such as tomatoes), Eggs, Chicken, Other Produce, Pork, Beef, and Fruits.
- No statistically significant differences in the estimated attribution percentages were found among most food categories.

Figure 3: Estimated percentage of *Escherichia coli* O157 illnesses (with 90% credibility intervals) for 2013, in descending order, attributed to each of 17 food categories, based on multi-year outbreak data,* United States




*Based on a model using outbreak data that gives equal weight to each of the most recent five years of data (2009–2013) and exponentially less weight to each earlier year (1998–2008).

***E. coli* O157 Key Results**

- Eighty percent of *E. coli* O157 illnesses were attributed to Beef and Vegetable Row Crops (such as leafy vegetables).
- The credibility intervals for the Beef and Vegetable Row Crops categories overlapped, indicating no statistically significant difference between their estimated attribution percentages.
- No illnesses were attributed to Eggs, Grains-Beans, Oils-Sugars, Other Seafood, or Pork.

Figure 4: Estimated percentage of *Listeria monocytogenes* illnesses (with 90% credibility intervals) for 2013, in descending order, attributed to each of 17 food categories, based on multi-year outbreak data,* United States




*Based on a model using outbreak data that gives equal weight to each of the most recent five years of data (2009–2013) and exponentially less weight to each earlier year (1998–2008).

Listeria monocytogenes Key Results

- Nearly 90% of illnesses were attributed to Fruits and Dairy.
- The credibility intervals for the Fruits and Dairy categories were quite wide, partly due to the small number of total outbreaks (26). The credibility intervals overlapped each other, and the intervals for the Fruit category overlapped those for food categories with much smaller estimated attribution percentages, such as Sprouts.
- The high point estimate for Fruits reflects the impact of a single large outbreak linked to cantaloupes in 2011.
- No illnesses were attributed to Eggs, Fish, Game, Grains-Beans, Oils-Sugars, Other Meat/Poultry, Other Seafood, or Seeded Vegetables.

Figure 5: Estimated percentage of *Campylobacter* illnesses (with 90% credibility intervals) for 2013, in descending order, attributed to each of 16 food categories, based on multi-year outbreak data,*† United States



*Based on a model using outbreak data that gives equal weight to each of the most recent five years of data (2009–2013) and exponentially less weight to each earlier year (1998–2008).

† *Campylobacter* estimates exclude results derived from Dairy outbreak data

***Campylobacter* Key Results**

- Almost 80% of non-Dairy illnesses were attributed to Chicken, Other Seafood, Seeded Vegetables, Vegetable Row Crops, and Other Meat/Poultry.
- No statistically significant differences in the estimated attribution percentages were found among most food categories.
- No illnesses were attributable to Eggs, Grains-Beans, or Sprouts.
- An attribution percentage for Dairy is not presented partly because most foodborne *Campylobacter* outbreaks were associated with unpasteurized milk, which is not widely consumed (The unadjusted Dairy attribution percentage was 68.3%, and was less than 10% for all other food categories). The adjusted Chicken attribution percentage increased from 9.3% to 29.2% after removing Dairy.

Discussion

This report uses data from 1998 through 2013 to provide outbreak-based attribution estimates for 2013 of the percentage of illnesses caused by four priority pathogens, assigning illnesses to each of 17 food categories. Data from foodborne disease outbreaks are the foundation of many foodborne illness source attribution analyses, in part because outbreak investigations often link illnesses to a specific food and the data are captured nationally. An IFSAC study found that outbreak and sporadic infections caused by the four priority pathogens were generally demographically similar; this supports the use of foodborne outbreaks for source attribution.⁷ These estimates can inform food safety decision-making and provide pathogen-specific direction for reducing foodborne illness.

The attribution of *Salmonella* illnesses to multiple food categories suggests that interventions designed to reduce illnesses from these pathogens need to target a variety of food categories. In contrast, *E. coli* O157 and *Lm* illnesses were attributed to fewer food categories: Beef and Vegetable Row Crops for *E. coli* O157 and Dairy and Fruits for *Lm*. The data suggest that interventions for *E. coli* O157 focusing on these two food categories may be most effective in reducing illnesses. However, the occurrence of outbreaks due to novel pathogen-food category pairs, such as an *E. coli* O157 outbreak due to SoyNut Butter in 2017,⁸ mandates vigilance in seeking unrecognized food sources of outbreaks and illnesses.

A single large outbreak linked to cantaloupes in 2011 had a major influence on the *Lm* attribution estimates. While novel at the time, more *Lm* outbreaks have now been linked to Fruit. Moreover, although the limited number of outbreaks and wide credibility intervals dictate caution in interpreting the attribution percentage for Dairy for *Lm*, the risk to pregnant women and persons with weakened immune systems of consuming soft cheese made from unpasteurized milk or in unsanitary conditions is well-recognized.⁹

Like *Salmonella*, *Campylobacter* illnesses were broadly attributed across multiple food categories. The attribution percentages for Dairy are not presented in the figures for *Campylobacter* for several reasons. Most *Campylobacter* outbreaks included in the database were associated with unpasteurized milk, which is not widely consumed by the general population. Moreover, an analysis of 38 case-control studies of sporadic campylobacteriosis found a much smaller percentage of illnesses attributable to consumption of raw milk than chicken.¹⁰ For example, a U.S. FoodNet case-control study attributed 1.5% of campylobacteriosis cases to consumption of unpasteurized milk, compared with 24% to consumption of chicken prepared in a restaurant.¹¹ Structured expert judgment studies estimate about 8–10% of foodborne campylobacteriosis to be attributable to dairy products (principally raw milk), compared with 33–72% to chicken.^{12–14} Thus, *Campylobacter* outbreaks in the Dairy food category appear to over-represent Dairy as a source of *Campylobacter* illness. After removing the Dairy outbreaks, the Chicken attribution increased to 29.2%, which is consistent with published literature.^{10, 16}

Our approach addresses a number of issues with outbreak-based foodborne illness source attribution, yet limitations associated with generalizing outbreak data to include sporadic illnesses remain and are well-documented.^{5,6} Our analysis is also subject to other uncertainties and biases. For pathogens with a small number of outbreaks, outbreaks with a very large illness count can have substantial influence on the attribution point estimate. Further, this analysis only included 36% of reported outbreaks caused by the four priority pathogens (1,043 of 2,919 outbreaks in which the confirmed or suspected implicated food could be assigned to a single food category), which may not be representative of all outbreaks from these pathogens. Finally, nearly 10% of illnesses in our analysis occurred among institutionalized populations, such as those in prisons, hospitals, and schools; these populations are easier to identify and collect complete data from, have fewer food options, and are not representative of the general population.

These estimates should not be interpreted as suggesting that all foods in a category are equally likely to transmit pathogens. Caution should also be exercised when comparing estimates across years, as a decrease in a percentage may result, not from a decrease in the number of illnesses attributed to that food, but from an increase in illnesses attributed to another food. The analyses show relative changes in percentage, not absolute changes in attribution to a specific food. Therefore, we advise using these results with other scientific data for decision-making.

Conclusions

IFSAC's work to provide a harmonized analytic approach for estimating foodborne illness source attribution from outbreak data can provide consistency in the use and interpretation of estimates across agencies. As more data become available and methods evolve, attribution estimates may improve. Annual updates to these estimates will enhance IFSAC's efforts to inform and engage stakeholders, and further their ability to assess whether prevention measures are working.

IFSAC continues to enhance attribution efforts by working on projects that address limitations identified in this report. For more information on IFSAC's completed and ongoing projects, visit <http://www.cdc.gov/foodsafety/ifsac/index.html>.

Reference List

1. Scallan E, Hoekstra RM, Angulo FJ, Tauxe RV, Widdowson M-A, Roy SL, Jones JL, Griffin PM. Foodborne illness acquired in the United States—major pathogens. *Emerg Infect Dis*. 2011 Jan;17(1):7-15.
2. Richardson LC, Bazaco M, Parker CC, Dewey-Mattia D, Golden N, Jones K, Klontz K, Travis C, Zablotsky-Kufel J, Cole D. An updated scheme for categorizing foods implicated in foodborne disease outbreaks: a tri-agency collaboration. *Foodborne Pathog Dis*. (in press).
3. Interagency Food Safety Analytics Collaboration (IFSAC) Working Group. Foodborne illness source attribution estimates for *Salmonella*, *Escherichia coli* O157 (*E. coli* O157), *Listeria monocytogenes* (*Lm*), and *Campylobacter* using outbreak surveillance data. February 2015. Available at: <https://www.cdc.gov/foodsafety/pdfs/ifsac-project-report-508c.pdf>.
4. Interagency Food Safety Analytics Collaboration (IFSAC). Activities and Events; 2015. 2017 [cited 2017 August 7]; Available at <https://www.cdc.gov/foodsafety/ifsac/events.html>.
5. Gould LH, Walsh KA, Vieira AR, et al. Surveillance for foodborne disease outbreaks — United States, 1998–2008. *MMWR Surveill Summ*. 2013 Jun;62(2): 1-34.
6. US Department of Health and Human Services, Centers for Disease Control and Prevention. Surveillance for Foodborne Disease Outbreaks, United States, 2015, Annual Report. Atlanta, Georgia 2017.
7. Ebel ED, Williams MS, Cole D, Travis CC, Klontz KC, Golden NJ, Hoekstra RM. Comparing characteristics of sporadic and outbreak-associated foodborne illnesses, United States, 2004–2011. *Emerg Infect Dis*. 2016 Jul;22(7): 1193-1200.
8. Centers for Disease Control and Prevention (CDC). Multistate Outbreak of Shiga toxin-producing *Escherichia coli* O157:H7 Infections Linked to I.M. Healthy Brand SoyNut Butter (Final Update). May 4, 2017 [cited 2017 August 7]; Available at <https://www.cdc.gov/ecoli/2017/o157h7-03-17/index.html>.
9. Swaminathan B, Gerner-Smidt P. The epidemiology of human listeriosis. *Microbes Infect*. 2007 Aug;9(10):1236-43.
10. Domingues AR, Pires SM, Halasa T, Hald T. Source attribution of human salmonellosis using a meta-analysis of case-control studies of sporadic infections. *Epidemiol Infect*. 2012 Jun;140(6):970-81.
11. Friedman CR, Hoekstra RM, Samuel M, Marcus R, Bender J, Shiferaw B, Reddy S, Ahuja S, Helfrick DL, Hardnett F, Carter M, Anderson B, Tauxe RV. Risk factors for sporadic *Campylobacter* infection in the United States: A case-control study in FoodNet sites. *Clin Infect Dis*. 2004 Apr 15;38 Suppl 3:S285–96.
12. Hoffmann SA, Fischbeck P, Krupnick A, McWilliams M. Using expert elicitation to link foodborne illnesses in the United States to foods. *J Food Prot*. 2007 May;70(5):1220–9.
13. Havelaar AH, Galindo AV, Kurowicka D, Cooke RM. Attribution of foodborne pathogens using structured expert elicitation. *Foodborne Pathog Dis*. 2008 Oct;5(5):649–659.
14. Davidson V.J, Ravel A, Nguyen TN, Fazil A, Ruzante AJ. Food-specific attribution of selected gastrointestinal illnesses: estimates from a Canadian expert elicitation survey. *Foodborne Pathog Dis*. 2011 Sept;8(9):983–995.
15. Butler AJ, Pintar KD, Thomas KM. Estimating the relative role of various subcategories of food, water, and animal contact transmission of 28 enteric diseases in Canada. *Foodborne Pathog Dis*. 2016 Feb;13(2):57–64.
16. Tam CC, Larose T, O'Brien SJ. Costed extension to the Second Study of Infectious Intestinal Disease (IID) in the Community: Identifying the proportion of foodborne disease in the UK and attributing foodborne disease by food commodity. London: Food Standards Agency Project B18021 (FS231043); 2014. Available at <https://www.food.gov.uk/science/research/foodborneillness/b14programme/b14projlist/fs231043ext>