




Contents lists available at ScienceDirect

Food Microbiology

journal homepage: www.elsevier.com/locate/fm

Examining the persistence of human Coronavirus 229E on fresh produce

Madeleine Blondin-Brosseau^a, Jennifer Harlow^a, Tanushka Doctor^a, Neda Nasheri^{a,b,*}^a National Food Virology Reference Centre, Bureau of Microbial Hazards, Health Canada, Ottawa, ON, Canada^b Department of Biochemistry, Microbiology and Immunology, Faculty of Medicine, University of Ottawa, ON, Canada

ARTICLE INFO

Keywords:

Human coronavirus
Persistence
Fecal-oral transmission
Foodborne transmission
Plaque assay
Droplet-digital RT-PCR
Decay rate

ABSTRACT

Human coronaviruses (HCoVs) are mainly associated with respiratory infections. However, there is evidence that highly pathogenic HCoVs, including severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and Middle East Respiratory Syndrome (MERS-CoV), infect the gastrointestinal (GI) tract and are shed in the fecal matter of the infected individuals. These observations have raised questions regarding the possibility of fecal-oral route as well as foodborne transmission of SARS-CoV-2 and MERS-CoV. Studies regarding the survival of HCoVs on inanimate surfaces demonstrate that these viruses can remain infectious for hours to days, however, there is limited data regarding the viral survival on fresh produce, which is usually consumed raw or with minimal heat processing. To address this knowledge gap, we examined the persistence of HCoV-229E, as a surrogate for highly pathogenic HCoVs, on the surface of commonly consumed fresh produce, including: apples, tomatoes, cucumbers and lettuce. Herein, we demonstrated that viral infectivity declines within a few hours post-inoculation (p.i) on apples and tomatoes, and no infectious virus was detected at 24h p.i, while the virus persists in infectious form for 72h p.i on cucumbers and lettuce. The stability of viral RNA was examined by droplet-digital RT-PCR (ddRT-PCR), and it was observed that there is no considerable reduction in viral RNA within 72h p.i.

1. Introduction

Coronaviruses that infect humans (HCoV) belong to alpha and beta genera of the *Coronaviridae* family. Four common HCoVs (229E, OC43, HKU1, and NL63) are responsible for 10–30% of common cold symptoms that can be mild to moderate (Perlman and Netland, 2009). SARS-CoV-2, which is responsible for the coronavirus disease 2019 (COVID-19) pandemic, is a betacoronavirus that uses angiotensin conversion enzyme 2 (ACE-2) for entry into the host cells. ACE-2 is abundantly expressed in the epithelium of the respiratory tract as well as the oral cavity, intestine and colon (Lamers et al., 2020; Qian et al., 2020). It is evident now that a considerable proportion of COVID-19 patients demonstrate gastrointestinal symptoms including nausea, vomiting, diarrhea, and abdominal pain (Cheung et al., 2020; Zhou et al., 2020; Scalfaferrri et al., 2020). SARS-CoV-2 RNA has been detected in more than 50% of patients' stool specimens (Wolfel et al., 2020; Wang et al., 2020; Huang et al., 2020; Cha et al., 2020), and several studies have confirmed that the virus detected in stool is infectious (Xiao et al., 2020; Zhou et al., 2020). Moreover, persistent fecal viral shedding has been observed in pediatric patients (Xu et al., 2020a,b) and there is evidence that SARS-CoV-2 can replicate productively in human enteroids and

enterocytes (Lamers et al., 2020; Zhou et al., 2020). More recently, it was demonstrated that multi-route mucosal inoculation (including oral inoculation) of African green monkeys with SARS-CoV-2 results in infection in both the respiratory and gastrointestinal tract (Hartman et al., 2020), and orally inoculated golden Syrian hamsters develop respiratory and intestinal infection (Chak-Yiu Lee, et al. 2020). Collectively, these observations suggest that fecal-oral transmission of SARS-CoV-2 is possible.

Although the primary route of transmission for HCoVs is inhalation of contaminated respiratory droplets and possible direct contact with contaminated fomites, there is concern that food could also act as a vehicle of transmission if contaminated with HCoVs. Food may become contaminated with HCoVs by contact with body secretions or fluids or by contact with soiled hands. Also, HCoVs may become aerosolized via talking, sneezing, or coughing of food handlers and then be deposited on food surfaces. Food not only may act as a fomite, but can also transport the virus to the potentially susceptible oral cavity and the GI tract (Xu et al., 2020a,b). There is evidence that certain HCoVs including HCoV-229E and MERS can survive GI conditions including low pH, digestive enzymes and bile (Zhou et al., 2017). If this is the case for SARS-CoV-2, the relatively high viral titre in stool and rectal swabs of

* Corresponding author. Department of Biochemistry, Microbiology and Immunology, Faculty of Medicine, University of Ottawa, ON, Canada.
E-mail address: neda.nasheri@canada.ca (N. Nasheri).

<https://doi.org/10.1016/j.fm.2021.103780>

Received 4 December 2020; Received in revised form 24 February 2021; Accepted 24 February 2021

Available online 26 February 2021

0740-0020/Crown Copyright © 2021 Published by Elsevier Ltd. All rights reserved.

the infected individuals could be explained by active viral replication in the GI tract. Furthermore, fecal-oral is the main route of transmission for enteric coronaviruses such as swine coronaviruses (Wang et al., 2019), canine coronaviruses (Decaro and Buonavoglia, 2011), and equine coronavirus (Pusterla et al., 2018), demonstrating that these viruses are not sensitive to the GI fluids.

Although the results should be cautiously interpreted, China has reported the finding of SARS-CoV-2 in imported frozen food commodities (Roxanne Liu, 2020; Yusha, 2020), and it was shown that the isolated virus from imported frozen cod is infectious in tissue culture (Liu et al., 2020). More recently, genetic evidence was provided that would link COVID-19 resurgence in Beijing to cold-chain food contamination (Pang et al., 2020). It was also demonstrated that this virus is stable for weeks in cold storage ($-80\text{ }^{\circ}\text{C}$ to $+4\text{ }^{\circ}\text{C}$) on artificially contaminated pork, chicken and salmon (Fisher et al., 2020). However, there is limited data on HCoVs survival on fresh produce.

Contamination of fresh produce may result in the transmission of not only the enteric viruses that are traditionally considered foodborne pathogens, but also possibly respiratory viruses such as adenoviruses, coronaviruses, and influenza viruses that can infect via contact with mucosal membranes (O'Brien et al., 2020). This is of particular concern for uncooked fruits and vegetables. Additionally, food handlers infected with respiratory viruses could potentially contaminate "cold foods" such as salads and sandwiches (Yepiz-Gomez et al., 2013), and spread the infection through various routes such as close contact and fomites. Thus, it is imperative to examine the viral behaviour and inactivation on fresh produce.

Since working with SARS-CoV-2 requires biosafety level 3 (BSL-3) laboratory containment conditions, the use of surrogate HCoVs have been suggested to expand the current knowledge on coronavirus survival and inactivation under various conditions (Guillier et al., 2020). For this reason, we chose HCoV-229E as a surrogate virus, since it has similar physicochemical properties to the more virulent HCoVs responsible for MERS and SARS (Warnes et al., 2015). In this study, we examined the ability of HCoV-229E to retain infectivity on the surface of select fruits and vegetables, and thus obtained representative survival data that can be used to conduct risk assessments of SARS-CoV-2 transmission via food.

2. Materials and methods

2.1. Cells and viruses

HCoV-229E and human embryonic lung cell line MRC-5 were obtained from the American Type Culture Collection (CCL-171 and VR-740, respectively). Cells were grown at $37\text{ }^{\circ}\text{C}$ and 5% CO_2 in culture media composed of Eagle's minimal essential medium, supplemented with 0.23% (w/v) sodium bicarbonate, $500\text{ }\mu\text{g/mL}$ Penicillin-Streptomycin (ThermoFisher scientific), 1% Glutamax-1, 1% non-essential amino acids, and foetal bovine serum (FBS) 5% (v/v).

2.2. -Sample preparation

Four different produce types, all purchased from local grocery stores in Ottawa, Ontario, were tested: Royal Gala apples, Traditional Series tomatoes, English cucumbers, and Romaine lettuce (PLU code 4173, 4799, 4593, and 4640, respectively). Ten time points were selected, in triplicates: 0h, 0.5h, 1h, 2h, 4h, 6h, 16h, 24h, 48h and 72h. For Romaine lettuce, only 4 time points were tested: 0h, 24, 48h and 72h. Each of the produce items was rinsed with water, dried with Kimwipes and disinfected with 70% ethanol. On the surface of each produce item, a 5 cm by 5 cm square was delimited using tape. For Romaine lettuce, the square was carefully drawn on each leaf to include both the rib and the leafy part. This area was inoculated with $100\text{ }\mu\text{L}$ of HCoV-229E (ATCC VR-740, 5×10^5 PFU/mL). The liquid was spread using the tip of the pipette, then allowed to fully dry for 1h. After the appropriate time lapse

at ambient conditions ($22\text{ }^{\circ}\text{C}$; relative humidity, 30% – 40%), the surface was sampled with a cotton swab, according to the ISO/TS 15216–1:2017 (ISO, 2017) method, which was then placed into the MRC-5 culture media previously described (Nasheri et al., 2020). Samples were processed immediately after swabbing.

2.3. Viral quantification

- plaque assay:

Viral quantification and survival time were determined by plaque assay using MRC-5 cells. Cells were grown at $37\text{ }^{\circ}\text{C}$ and 5% CO_2 in the culture medium previously described for up to three days, before being seeded, transferred into 12-well plates at a targeted concentration of 5×10^5 cells/mL and incubated to reach a confluency of 80% – 90% . Samples were diluted in culture medium and $100\text{ }\mu\text{L}$ of at least two dilutions were used in duplicate to infect the prepared plates for 90 min at $35\text{ }^{\circ}\text{C}$ and 5% CO_2 . Plates were manually rocked every 10 min during the infection phase. Cells were then washed with phosphate buffered saline (PBS) and covered with 2 mL of overlay media, composed of a $50/50$ mix of $2 \times$ culture medium previously described and 0.5% agarose. Plates were incubated at $35\text{ }^{\circ}\text{C}$ and 5% CO_2 for 3–4 days. Cell monolayers were fixed using 3.7% paraformaldehyde for 4–16 h, freed from overlay plugs by running under tap water and stained with 0.1% crystal violet for 20 min. Plaques were counted for each dilution to determine the viral titre.

- Determining limit of detection

Each produce item was artificially inoculated with a serial dilution of the viral stock in triplicate. At T_0 , the virus was extracted and assayed by plaque assay as described above. The plaques were counted for each dilution and results were analyzed to determine the highest dilutions (lowest titre) for which plaques were still obtained in triplicate experiments.

- Recovery rate calculation

The recovery efficiency was determined by calculating the ratio between the viral titre recovered at T_0 and the viral titre that was used to inoculate the sample.

$$\text{Recoveryrate}(\%) : \frac{\text{obtained viral titre (PFU/mL)}}{\text{inoculated viral titre (PFU/mL)}} \times 100$$

- Estimating the decay rate:

Viral decay rate was calculated as described previously (Long and Short, 2016). Briefly, linear regressions of the natural logarithm of virus abundance versus time (in hours) was calculated. The slope of the regressions represent the decay rate and when multiplied by 100, represent percentage of infectivity lost per hour. Viral half-life was calculated by dividing $\ln(2)$ by the slope.

- ddRT-PCR

For each produce item, except lettuce, all triplicates of 10 time points were tested. Viral RNA was isolated using a QIAamp viral RNA kit (QIAGEN) and diluted in sterile molecular biology grade water (Corning). The QX200 ddPCR system (Bio-Rad) was used for quantification and all PCR reactions were prepared using the One-Step RT-ddPCR Advanced Kit for Probes (Bio-Rad Cat# 1864022). Primers used were previously described in (Vijgen et al., 2005): Forward primer 229E-FP (5-TTCCGACGTGCTCGAAGCTTT-3; GenBank accession no. M33560; nt 474 to 493) and reverse primer 229E-RP (5-CCAACACGGTTGTGACAGTGA-3; nt 523 to 543). A new probe that would complement the primers and be compatible with TaqMan qPCR requirements (ABI 7700

Users Manual) was designed by using Integrated DNA Technologies (IDT) OligoAnalyzer tool. The new probe had the appropriate dissociation temperature and a minimal likelihood for duplex or hairpin formation: 229E-PR (5'-/56-FAM/TGCATTGAC/ZEN/CTCAGGATTCCAT GCCC/3IABkFQ/-3'). Each PCR reaction contained 5 μ L of RNA, 1000 nmol/L of each primer, and 280 nmol/L of each probe. All samples were tested in duplicate. Droplets were generated using the QX200 droplet generator (Bio-Rad) according to the manufacturer's protocols, and PCR was performed using the following cycling conditions: an initial reverse transcription at 48 °C for 30 min, followed by PCR activation at 95 °C for 10 min and 45 cycles of amplification (15 s at 95 °C and 1 min at 60 °C). Droplets were detected in the QX200 droplet reader and analyzed using the Quantasoft version 1.7.4.0917 (Bio-Rad) software.

3. Results

3.1. Recovery efficiency from produce

As shown in Table 1, the recovery efficiency for infectious HCoV-229E from all the tested commodities is well above 1%, with the highest recovery rate (10.8%) from tomatoes and the lowest (4.1%) from cucumbers. The limit of detection (LOD) for each commodity is determined as the lowest spiking concentration that produced plaques for all three replicates. As indicated in Table 2, the LOD was approximately 125 PFU for tomatoes and apples, and 50 PFU for cucumbers.

3.2. Persistence of infectivity

We artificially inoculated the surface of apples, tomatoes and cucumbers with 5×10^4 PFU of HCoV-229E, which is consistent with the amount of virus that is typically exhaled by an infected individual (Ma et al., 2020). Fig. 1 shows the persistence in infectivity of HCoV-229E at RT within 72 h p.i. The change in infectious viral titre is similar in apples and tomatoes with a progressive decline in infectivity up to 16h p.i. (Fig. 1, Table 3). No infectious viral particles were isolated from tomatoes and apples at 24 h p.i., which demonstrates that viral infectivity is reduced below the LOD (i.e. >3 log reduction). However, infectious viral particles were detected on cucumbers up to 72 h p.i. Within the first 4 h p.i., viral infectivity reduces over 1 log on tomatoes and apples (1.18 and 1.27 log, respectively), while the reduction on cucumbers is only 0.75 log (Table 3). The reduction in infectivity is less than 2 log at 24 h p.i. on cucumbers and by 72 h p.i. reaches approximately 2.5 log. No infectious viral particles were detected on cucumbers at 96 h p.i.

The median decay rate of HCoV-229E on apples and tomatoes was similar at 30%/h and 34%/h respectively, while the median decay rate on cucumbers was considerably lower at 7.7%/h. The median half-life of the virus on apples and tomatoes was 2.3h and 2.05h respectively and the median half-life on cucumbers was 9.05h (Table 4).

Many factors might have contributed to the difference in viral survival on apples and tomatoes compared to cucumbers, but we hypothesized that the difference in surface pH between the examined produce could partly explained this observation. The surface pH of cucumbers (5.7) is considerably higher than the surface pH of tomatoes and apples (4.2 and 3.9, respectively) (McGlynn, 2016). For this reason, we examined viral survival on the surface of Romaine lettuce, which has a surface pH close to cucumbers (5.8) (McGlynn, 2016). As shown in Fig. 2, similar to what has been observed for cucumbers, infectious

Table 1

Recovered viral titre at T_0 and recovery rate in percentage for each produce type. The results are the mean of 3 independent experiments.

Produce	Titer at T_0 (PFU/mL)	Recovery rate(%)
Apple	1.45E+03	5.81
Tomato	2.69E+03	10.77
Cucumber	1.20E+03	4.09

Table 2

Detection of HCoV-229E on the surface of different produce. Samples were inoculated with 10^4 to 10^1 PFU of HCoV-229E and examined by plaque assay at T_0 . ND is not detected.

Produce	Viral Inoculum (PFU)						
	10,000	1000	500	250	125	50	10
Apple	3/3	3/3	3/3	3/3	3/3	ND	ND
Tomato	3/3	3/3	3/3	3/3	3/3	2/3	ND
Cucumber	3/3	3/3	3/3	3/3	3/3	3/3	ND

HCoV-229E was consistently isolated at 24, 48, and 74 h p.i from the surface of lettuce. The gradual pattern of infectivity loss on lettuce resembles to what has been observed on cucumbers (Fig. 2). This might indicate that the surface pH might play a role in viral survival at ambient temperature (see Fig. 2).

3.3. Persistence of viral RNA

We next set out to investigate the persistence of viral RNA on the examined produce over 72 h.p.i. at ambient temperature. As demonstrated in Fig. 3, no drastic reduction in viral RNA titre was observed over a 72h p.i. period. On apples, tomatoes, and cucumbers, viral RNA decreased by approximately 0.7 log, 0.5 log, and 0.3 log, respectively compared to T_0 . Altogether, these observations demonstrate that viral RNA is more resistant to degradation compared to viral infectivity on the surface of produce.

4. Discussion

Currently, SARS-CoV-2 is not considered a foodborne virus, and to date, there is no conclusive evidence of foodborne transmission of SARS-CoV-2. However, the traditional epidemiological foodborne investigation is unlikely to be employed with COVID-19 patients. For example, it is unlikely that infected people are asked to recall foods that they may have consumed during the period when they became infected. Without this information, any association between SARS-CoV-2 and foods cannot be made, and understanding the role of foodborne transmission remains elusive. Obtaining this epidemiological information would be helpful for efficient contact-tracing and source-tracking as about half of COVID-19 patients can not recall how and where they contracted the virus (Tenforde et al., 2020).

In this study, we used the ISO/TS 15216–1:2017 (ISO, 2017) method for the recovery of HCoV-229E from the examined surfaces, and we assessed the recovery rates by plaque assay, which indicate that the recovered viruses were infectious. The recovery range that we obtained was from 4.09% to 10.77%, which is significantly higher than 1% recovery rate that is considered acceptable by the method. However, we speculate that the recovery efficiency would be higher if the genetic materials were assessed instead of viral infectivity.

Environmental persistence of HCoVs has been examined by different groups, who have obtained contradictory results (Aboubakr et al., 2020). One study has shown that the stability of SARS-CoV-2 and SARS-CoV-1 on dry surfaces at RT is similar, with no infectious virus being retrieved after 72h p.i. (van Doremalen et al., 2020), while, Chin et al. recovered infectious SARS-CoV-2 from plastic and stainless steel up to 7 days p.i. (Chin et al., 2020). Keevil and coworkers reported that HCoV-229E remains infectious for 5 days at RT on a range of surface materials including glass and PVC, while it is rapidly inactivated on the surface of copper alloys (Warnes et al., 2015). In another study, more relevant to this work, it was shown that the infectivity of HCoV-229E is completely abolished within 4 days p.i. on lettuce at 4 °C (Yepiz-Gomez et al., 2013). Recently, it was demonstrated that SARS-CoV-2 remains infectious on salmon at RT for 2 days (Manman, 2020). Herein, we only examined viral survival at ambient temperature and we have shown the infectivity of HCoV-229E is reduced to below LOD followed by 24h




Fig. 1. Persistence of infectious HCoV-229E on commonly consumed fruits and vegetables. Approximately 5×10^4 PFU HCoV-229E (100 μ L viral stock) was applied to the tested surface and incubated at ambient conditions (22 °C; relative humidity, 30%–40%). Virus was extracted and assayed for infectivity at various time points as described in the text. The data represent the average of three independent experiments. Error bars represent standard deviation.

Table 3
Log reduction in viral titre compared to T_0 . The results are the mean of 3 independent experiments \pm Standard Deviation.

Time point	Apples	Tomatoes	Cucumbers
0.5h	0.09 \pm 0.01	0.09 \pm 0.05	0.10 \pm 0.01
1h	0.23 \pm 0.06	0.14 \pm 0.04	0.33 \pm 0.11
2h	0.90 \pm 0.12	0.68 \pm 0.05	0.38 \pm 0.11
4h	1.08 \pm 0.18	1.05 \pm 0.02	0.76 \pm 0.01
6h	1.27 \pm 0.08	1.18 \pm 0.06	0.79 \pm 0.04
16h	2.40 \pm 0.33	2.37 \pm 0.09	1.26 \pm 0.06
24h	3.16	3.43	1.92 \pm 0.15
48h	3.16	3.43	2.09 \pm 0.16
72h	3.16	3.43	2.48 \pm 0.035

Table 4
Decay rate (DR) in percentage and viral half-life (HL) in hours (h) on each produce type. The results are the median of 3 independent experiments \pm Standard Deviation.

	DR (%)	HL (h)
Apple	30 \pm 0.25	2.3 \pm 0.02
Tomato	34 \pm 0.1	2.05 \pm 0.06
Cucumber	7.7 \pm 0.6	9.05 \pm 0.75

incubation on tomatoes and apples, and 96h on cucumbers, and lettuce.

At this point, we speculate that the longer survival on cucumbers, and lettuce compared to apples and tomatoes could be partly explained by the difference in surface pH of these commodities. The influence of pH on the stability of several coronaviruses has been studied and it has been shown that in general, coronaviruses are more stable at near neutral pH as compared to acidic or alkaline pH (Aboubakr et al., 2020). As such, the near neutral surface pH of cucumbers, and lettuce (5.7, and 5.8, respectively), compared to the more acidic surface pH of tomatoes and apples (4.2 and 3.9, respectively), could be more suitable for the survival of HCoV-229E (McGlynn, 2016). It should also be noted that the LOD on cucumbers was lower compared to apples and tomatoes (50 PFU compared with 125 PFUs, respectively). Thus, it is possible that HCoV-229E remained infectious by 24 h p.i. on apples and tomatoes but the titre was below the LOD. However, the decay rate on cucumbers is considerably slower compared to apples and tomatoes (Fig. 1 and Table 4), and the viral half-life on cucumbers is very close to the viral half-life on plastic (van Doremalen et al., 2020) (9.05h and 9.04h, respectively). Further investigation is needed to determine whether the surface of apples and tomatoes has some virucidal properties that may




Fig. 2. Persistence of infectious HCoV-229E on Romaine lettuce. Approximately 5×10^4 PFU HCoV-229E (100 μ L viral stock) was applied to the tested surface and incubated at ambient conditions (22 °C; relative humidity, 30%–40%). Virus was extracted and assayed for infectivity at various time points as described in the text. The data represent the average of three independent experiments. Error bars represent standard deviation.

lead to a more rapid viral inactivation. Thus, our results are in accordance with the previous findings that HCoVs lose their infectivity within a few days on inanimate surfaces at RT (Sizun et al., 2000). Therefore, if produce becomes contaminated with HCoVs through irrigation or contaminated hands during pre- or post-harvest, while being stored at ambient temperature, the risk will be considerably reduced by the time it reaches the consumers. However, if the contamination occurs at the end of the food processing chain, for example by infected personnel in a restaurant setting, where the prepared food is consumed within a few minutes, there is a potential risk for infection (Zelner et al., 2020; de Wit et al., 2007). Although, to date there is no evidence that ingestion of SARS-CoV-2 could lead to infection.

The persistence of viral RNA on the studied produce for several days despite the loss of infectivity (Fig. 3), can be explained by the high environmental resilience of the coronavirus shell, which protects the viral genome (Goh et al., 2020).

It should be noted that our study involved experimental inoculation of fresh produce with HCoV-229E, and thus may not be fully representative of potential natural contamination. However, the infectious titre of virus used for inoculation of samples in the current study is representative of a worst-case scenario, if virus was found to be present on




Fig. 3. Persistence of viral RNA on commonly consumed fruits and vegetables. Approximately 2×10^8 RNA copies of HCoV-229E (100 μ L of viral stock) was applied to the tested surface and incubated at ambient conditions (22 °C; relative humidity, 30%–40%). Virus was extracted at indicated time points and viral RNA was quantified by ddRT-PCR. The data represent the average of three independent experiments. Error bars represent standard deviation.

fresh produce. Herein, we attempted to address an important knowledge gap regarding the survival of human coronaviruses on fresh produce at ambient temperature, although to date, there is no conclusive evidence that food could be a vehicle for SARS-CoV-2 transmission. Potential foodborne transmission poses important public health implications and may partly explain the possible recurrence of the disease and its persistent transmission. Thus, our results could support more robust decision-making concerning risk assessment for foodborne transmission of human coronaviruses.

Declaration of competing interest

The authors declare no conflict of interest.

Acknowledgements

The authors would like to thank Dr. Brent Dixon and Dr. Franco Pagotto from the Bureau of Microbial Hazards for kindly reviewing the manuscript and providing insightful comments. This study is financially supported by the Bureau of Microbial Hazards, Health Canada.

References

- Aboubakr, H.A., Sharafeldin, T.A., Goyal, S.M., 2020. Stability of SARS-CoV-2 and other coronaviruses in the environment and on common touch surfaces and the influence of climatic conditions: a review. *Transbound Emerg. Dis.* <https://doi.org/10.1111/tbed.13707>.
- Cha, M.H., Regueiro, M., Sandhu, D.S., 2020. Gastrointestinal and hepatic manifestations of COVID-19: a comprehensive review. *World J. Gastroenterol.* 26, 2323–2332.
- Chak-Yiu Lee, A., Zhang, A.J., Fuk-Woo Chan, J., Li, C., Fan, Z., Liu, F., Chen, Y., Liang, R., Sridhar, S., Cai, J.P., Kwok-Man Poon, V., Chung-Sing Chan, C., Kai-Wang To, K., Yuan, S., Zhou, J., Chu, H., Yuen, K.Y., 2020. Oral SARS-CoV-2 inoculation establishes subclinical respiratory infection with virus shedding in golden Syrian hamsters. *Cell. Rep. Med.* 100121.
- Cheung, K.S., Hung, I.F., Chan, P.P., Lung, K.C., Tso, E., Liu, R., Ng, Y.Y., Chu, M.Y., Chung, T.W., Tam, A.R., Yip, C.C., Leung, K.H., Yim-Fong Fung, A., Zhang, R.R., Lin, Y., Cheng, H.M., Zhang, A.J., To, K.K., Chan, K.H., Yuen, K.Y., Leung, W.K., 2020. Gastrointestinal manifestations of SARS-CoV-2 infection and virus load in fecal samples from the Hong Kong cohort and systematic review and meta-analysis. *Gastroenterology* 59 (1), 81–95.
- Chin, A.W.H., Chu, J.T.S., Perera, M.R.A., Hui, K.P.Y., Yen, H.L., Chan, M.C.W., Peiris, M., Poon, L.L.M., 2020. Stability of SARS-CoV-2 in different environmental conditions. *Lancet Microbe* 1, e10–5247.
- de Wit, M.A., Widdowson, M.A., Vennema, H., de Bruin, E., Fernandes, T., Koopmans, M., 2007. Large outbreak of norovirus: the baker who should have known better. *J. Infect.* 55, 188–193.
- Decaro, N., Buonavoglia, C., 2011. Canine coronavirus: not only an enteric pathogen. *Vet. Clin. North Am. Small Anim. Pract.* 41, 1121–1132.
- Fisher, D., Reilly, A., Kang Eng Zheng, A., Cook, A.R., Anderson, D.E., 2020. Seeding of outbreaks of COVID-19 by contaminated fresh and frozen food. *BioRxiv*.
- Goh, G.K., Dunker, A.K., Foster, J.A., Uversky, V.N., 2020. Shell disorder analysis predicts greater resilience of the SARS-CoV-2 (COVID-19) outside the body and in body fluids. *Microb. Pathog.* 144, 104177.
- Guillier, L., Martin-Latil, S., Chaix, E., Thebault, A., Pavio, N., Le Poder, S., Batejat, C., Biot, F., Koch, L., Schaffner, D., Sanaa, M., Covid-19 Emergency Collective Expert Appraisal Group, 2020. Modelling the inactivation of viruses from the Coronaviridae family in response to temperature and relative humidity in suspensions or surfaces. *Appl. Environ. Microbiol.* 01244–20–20. <https://doi.org/10.1128/AEM.01244-20>.
- Hartman, A.L., Nambulli, S., McMillen, C.M., White, A.G., Tilston-Lunel, N.L., Albe, J.R., Cottle, E., Dunn, M.D., Frye, L.J., Gilliland, T.H., Olsen, E.L., O'Malley, K.J., Schwarz, M.M., Tomko, J.A., Walker, R.C., Xia, M., Hartman, M.S., Klein, E., Scanga, C.A., Flynn, J.L., Klimstra, W.B., McElroy, A.K., Reed, D.S., Duprex, W.P., 2020. SARS-CoV-2 infection of African green monkeys results in mild respiratory disease discernible by PET/CT imaging and shedding of infectious virus from both respiratory and gastrointestinal tracts. *PLoS Pathog.* 16, e1008903 <https://doi.org/10.1371/journal.ppat.1008903>.
- Huang, C., Wang, Y., Li, X., Ren, L., Zhao, J., Hu, Y., Zhang, L., Fan, G., Xu, J., Gu, X., Cheng, Z., Yu, T., Xia, J., Wei, Y., Wu, W., Xie, X., Yin, W., Li, H., Liu, M., Xiao, Y., Gao, H., Guo, L., Xie, J., Wang, G., Jiang, R., Gao, Z., Jin, Q., Wang, J., Cao, B., 2020. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 395, 497–506.
- ISO, 2017. EN ISO 15216-1:2017. Microbiology of the food chain—horizontal method for determination of hepatitis A virus and norovirus using real-time RT-PCR- Part1: method for quantification. International Organization for Standardization.
- Lamers, M.M., Beumer, J., van der Vaart, J., Kooops, K., Puschhof, J., Breugem, T.I., Ravelli, R.B.G., Paul van Schayck, J., Mykityn, A.Z., Duimel, H.Q., van Donselaar, E., Riesebosch, S., Kuijpers, H.J.H., Schippers, D., van de Wetering, W.J., de Graaf, M., Koopmans, M., Cuppen, E., Peters, P.J., Haagmans, B.L., Clevers, H., 2020. SARS-CoV-2 productively infects human gut enterocytes. *Science* 369 (6499), 50–54.
- Liu, P., Yang, M., Zhao, X., Guo, Y., Wang, L., Zhang, J., Lei, W., Han, W., Jiang, F., Liu, W.J., Gao, G.F., Wu, G., 2020. Cold-chain transportation in the frozen food industry may have caused a recurrence of COVID-19 cases in destination: successful isolation of SARS-CoV-2 virus from the imported frozen cod package surface. *Biosaf Health* 2 (4), 199–201.
- Long, A.M., Short, S.M., 2016. Seasonal determinations of algal virus decay rates reveal overwintering in a temperate freshwater pond. *ISME J.* 10, 1602–1612.
- Ma, J., Qi, X., Chen, H., Li, X., Zhang, Z., Wang, H., Sun, L., Zhang, L., Guo, J., Morawska, L., Grinshpun, S.A., Biswas, P., Flagan, R.C., Yao, M., 2020. COVID-19 patients in earlier stages exhaled millions of SARS-CoV-2 per hour. *Clin. Infect. Dis.* <https://doi.org/10.1093/cid/ciaa1283>.
- Manman, D.e. a., 2020. Long-term survival of salmon-attached SARS-CoV-2 at 4 °C as a potential source of transmission in seafood markets. *BioRxiv*.
- McGlynn, W., 2016. The Importance of Food pH in Commercial Canning Operations. Food Technology Fact Sheet.
- Nasheri, N., Harlow, J., Chen, A., Corneau, N., Bidawid, S., 2020. Evaluation of bead-based assays in the isolation of foodborne viruses from low-moisture foods. *J. Food Protect.* 83, 388–396.
- O'Brien, B., Goodridge, L., Ronholm, J., Nasheri, N., 2020. Exploring the potential of foodborne transmission of respiratory viruses. *Food Microbiol.* 103709.
- Pang, X., Ren, L., Wu, S., Ma, W., Yang, J., Di, L., Li, J., Xiao, Y., Kang, L., Du, S., Du, J., Wang, J., Li, G., Zhai, S., Chen, L., Zhou, W., Lai, S., Gao, L., Pan, Y., Wang, Q., Li, M., Wang, J., Huang, Y., Wang, J., COVID-19 Field Response Group and COVID-19 Laboratory Testing Group, 2020. Cold-chain food contamination as the possible origin of Covid-19 resurgence in Beijing. *Natl Sci Rev* nwa264.
- Perlman, S., Netland, J., 2009. Coronaviruses post-SARS: update on replication and pathogenesis. *Nat. Rev. Microbiol.* 7, 439–450.

- Pusterla, N., Vin, R., Leutenegger, C.M., Mittel, L.D., Divers, T.J., 2018. Enteric coronavirus infection in adult horses. *Vet. J.* 231, 13–18.
- Qian, Q., Fan, L., Liu, W., Li, J., Yue, J., Wang, M., Ke, X., Yin, Y., Chen, Q., Jiang, C., 2020. Direct evidence of active SARS-CoV-2 replication in the intestine. *Clin. Infect. Dis.* ciaa925.
- Roxanne Liu, D.S., 2020. Traces of coronavirus found in frozen chicken wings, shrimp packaging in China, Global News. Available at: <https://globalnews.ca/news/7271588/coronavirus-traces-frozen-chicken-shrimp-packaging/>.
- Scalaferrri, F., Ianiro, G., Privitera, G., Lopetuso, L.R., Vetrone, L.M., Petito, V., Pugliese, D., Neri, M., Cammarota, G., Ringel, Y., Costamagna, G., Gasbarrini, A., Boskoski, I., Armuzzi, A., 2020. The thrilling journey of SARS-CoV-2 into the intestine: from pathogenesis to future clinical implications. *Inflamm. Bowel Dis.* <https://doi.org/10.1093/ibd/izaa181>.
- Sizun, J., Yu, M.W., Talbot, P.J., 2000. Survival of human coronaviruses 229E and OC43 in suspension and after drying on surfaces: a possible source of hospital-acquired infections. *J. Hosp. Infect.* 46, 55–60.
- Tenforde, M.W., Billig Rose, E., Lindsell, C.J., Shapiro, N.I., Files, D.C., Gibbs, K.W., Prekker, M.E., Steingrub, J.S., Smithline, H.A., Gong, M.N., Aboodi, M.S., Exline, M. C., Henning, D.J., Wilson, J.G., Khan, A., Qadir, N., Stubblefield, W.B., Patel, M.M., Self, W.H., Feldstein, L.R., CDC COVID-19 Response Team, 2020. Characteristics of adult outpatients and inpatients with COVID-19 - 11 academic medical centers, United States, march-may 2020. *MMWR Morb. Mortal. Wkly. Rep.* 69, 841–846.
- van Doremalen, N., Bushmaker, T., Morris, D.H., Holbrook, M.G., Gamble, A., Williamson, B.N., Tamin, A., Harcourt, J.L., Thornburg, N.J., Gerber, S.I., Lloyd-Smith, J.O., de Wit, E., Munster, V.J., 2020. Aerosol and surface stability of SARS-CoV-2 as compared with SARS-CoV-1. *N. Engl. J. Med.* 382, 1564–1567.
- Vijgen, L., Keyaerts, E., Moes, E., Maes, P., Duson, G., Van Ranst, M., 2005. Development of one-step, real-time, quantitative reverse transcriptase PCR assays for absolute quantitation of human coronaviruses OC43 and 229E. *J. Clin. Microbiol.* 43, 5452–5456.
- Wang, Q., Vlasova, A.N., Kenney, S.P., Saif, L.J., 2019. Emerging and re-emerging coronaviruses in pigs. *Curr. Opin. Virol.* 34, 39–49.
- Wang, W., Xu, Y., Gao, R., Lu, R., Han, K., Wu, G., Tan, W., 2020. Detection of SARS-CoV-2 in different types of clinical specimens. *J. Am. Med. Assoc.* <https://doi.org/10.1001/jama.2020.3786>.
- Warnes, S.L., Little, Z.R., Keevil, C.W., 2015. Human coronavirus 229E remains infectious on common touch surface materials. *mBio* 6, e01697–15.
- Wolfel, R., Corman, V.M., Guggemos, W., Seilmaier, M., Zange, S., Müller, M.A., Niemeyer, D., Jones, T.C., Vollmar, P., Rothe, C., Hoelscher, M., Bleicker, T., Brunink, S., Schneider, J., Ehmann, R., Zwirgmaier, K., Drosten, C., Wendtner, C., 2020. Virological assessment of hospitalized patients with COVID-2019. *Nature* 581, 465–469.
- Xiao, F., Sun, J., Xu, Y., Li, F., Huang, X., Li, H., Zhao, J., Huang, J., Zhao, J., 2020. Infectious SARS-CoV-2 in feces of patient with severe COVID-19. *Emerg. Infect. Dis.* 26, 1920–1922.
- Xu, H., Zhong, L., Deng, J., Peng, J., Dan, H., Zeng, X., Li, T., Chen, Q., 2020a. High expression of ACE2 receptor of 2019-nCoV on the epithelial cells of oral mucosa. *Int. J. Oral Sci.* 12, 8–20.
- Xu, Y., Li, X., Zhu, B., Liang, H., Fang, C., Gong, Y., Guo, Q., Sun, X., Zhao, D., Shen, J., Zhang, H., Liu, H., Xia, H., Tang, J., Zhang, K., Gong, S., 2020b. Characteristics of pediatric SARS-CoV-2 infection and potential evidence for persistent fecal viral shedding. *Nat. Med.* 26, 502–505.
- Yepiz-Gomez, M.S., Gerba, C.P., Bright, K.R., 2013. Survival of respiratory viruses on fresh produce. *Food Environ. Virol.* 5 (3), 150–156.
- Yusha, Z., 2020. China's CDC experts investigate Xinfadi market three times, announce groundbreaking virus tracing discovery, Global Times. Available at: <https://www.globaltimes.cn/content/1192146.shtml>.
- Zelner, J., Adams, C., Havumaki, J., Lopman, B., 2020. Understanding the importance of contact heterogeneity and variable infectiousness in the dynamics of a large norovirus outbreak. *Clin. Infect. Dis.* 70, 493–500.
- Zhou, J., Li, C., Liu, X., Chiu, M.C., Zhao, X., Wang, D., Wei, Y., Lee, A., Zhang, A.J., Chu, H., Cai, J.P., Yip, C.C., Chan, I.H., Wong, K.K., Tsang, O.T., Chan, K.H., Chan, J. F., To, K.K., Chen, H., Yuen, K.Y., 2020. Infection of bat and human intestinal organoids by SARS-CoV-2. *Nat. Med.* 26 (7), 1077–1083.
- Zhou, J., Li, C., Zhao, G., Chu, H., Wang, D., Yan, H.H., Poon, V.K., Wen, L., Wong, B.H., Zhao, X., Chiu, M.C., Yang, D., Wang, Y., Au-Yeung, R.K.H., Chan, I.H., Sun, S., Chan, J.F., To, K.K., Memish, Z.A., Corman, V.M., Drosten, C., Hung, I.F., Zhou, Y., Leung, S.Y., Yuen, K.Y., 2017. Human intestinal tract serves as an alternative infection route for Middle East respiratory syndrome coronavirus. *Sci. Adv.* 3, eaao4966.